

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 6 : <b>C12N 15/12, C07K 14/47, C12N 15/10, 15/66, C12Q 1/68, G01N 33/50, C07K 16/18, G01N 33/53, A61K 48/00, 38/17</b></p>	<p><b>A2</b></p>	<p>(11) International Publication Number: <b>WO 99/06549</b> (43) International Publication Date: <b>11 February 1999 (11.02.99)</b></p>
<p>(21) International Application Number: <b>PCT/IB98/01231</b> (22) International Filing Date: <b>31 July 1998 (31.07.98)</b> (30) Priority Data: <b>08/905,279 1 August 1997 (01.08.97) US</b> (71) Applicant (for all designated States except US): <b>GENSET [FR/FR]; 24, rue Royale, F-75008 Paris (FR).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>DUMAS MILNE ED- WARDS, Jean-Baptiste [FR/FR]; 8, rue Grégoire de Tours, F-75006 Paris (FR). DUCLERT, Aymeric [FR/FR]; 6 ter, rue Victorine, F-94100 Saint-Maur (FR). LACROIX, Bruno [FR/FR]; 93, route de Vourles, F-69230 Saint-Genis Laval (FR).</b> (74) Agents: <b>MARTIN, Jean-Jacques et al.; Cabinet Regimbeau, 26, avenue Kléber, F-75116 Paris (FR).</b></p>		<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: <b>5' ESTs FOR SECRETED PROTEINS EXPRESSED IN TESTIS AND OTHER TISSUES</b></p> <p>(57) Abstract</p> <p>The sequences of 5' ESTs derived from mRNAs encoding secreted proteins are disclosed. The 5' ESTs may be to obtain cDNAs and genomic DNAs corresponding to the 5' ESTs. The 5' ESTs may also be used in diagnostic, forensic, gene therapy, and chromosome mapping procedures. Upstream regulatory sequences may also be obtained using the 5' ESTs. The 5' ESTs may also be used to design expression vectors and secretion vectors.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



## 5' ESTs FOR SECRETED PROTEINS EXPRESSED IN TESTIS AND OTHER TISSUES

### Background of the Invention

The estimated 50,000-100,000 genes scattered along the human chromosomes offer  
5 tremendous promise for the understanding, diagnosis, and treatment of human diseases. In  
addition, probes capable of specifically hybridizing to loci distributed throughout the human  
genome find applications in the construction of high resolution chromosome maps and in the  
identification of individuals.

In the past, the characterization of even a single human gene was a painstaking  
10 process, requiring years of effort. Recent developments in the areas of cloning vectors, DNA  
sequencing, and computer technology have merged to greatly accelerate the rate at which  
human genes can be isolated, sequenced, mapped, and characterized. Cloning vectors such as  
yeast artificial chromosomes (YACs) and bacterial artificial chromosomes (BACs) are able to  
accept DNA inserts ranging from 300 to 1000 kilobases (kb) or 100-400 kb in length  
15 respectively, thereby facilitating the manipulation and ordering of DNA sequences distributed  
over great distances on the human chromosomes. Automated DNA sequencing machines  
permit the rapid sequencing of human genes. Bioinformatics software enables the  
comparison of nucleic acid and protein sequences, thereby assisting in the characterization of  
human gene products.

20 Currently, two different approaches are being pursued for identifying and  
characterizing the genes distributed along the human genome. In one approach, large  
fragments of genomic DNA are isolated, cloned, and sequenced. Potential open reading  
frames in these genomic sequences are identified using bioinformatics software. However,  
this approach entails sequencing large stretches of human DNA which do not encode proteins  
25 in order to find the protein encoding sequences scattered throughout the genome. In addition  
to requiring extensive sequencing, the bioinformatics software may mischaracterize the  
genomic sequences obtained. Thus, the software may produce false positives in which non-  
coding DNA is mischaracterized as coding DNA or false negatives in which coding DNA is  
misabeled as non-coding DNA.

30 An alternative approach takes a more direct route to identifying and characterizing  
human genes. In this approach, complementary DNAs (cDNAs) are synthesized from

isolated messenger RNAs (mRNAs) which encode human proteins. Using this approach, sequencing is only performed on DNA which is derived from protein coding portions of the genome. Often, only short stretches of the cDNAs are sequenced to obtain sequences called expressed sequence tags (ESTs). The ESTs may then be used to isolate or purify extended  
5 cDNAs which include sequences adjacent to the EST sequences. The extended cDNAs may contain all of the sequence of the EST which was used to obtain them or only a portion of the sequence of the EST which was used to obtain them. In addition, the extended cDNAs may contain the full coding sequence of the gene from which the EST was derived or, alternatively, the extended cDNAs may include portions of the coding sequence of the  
10 gene from which the EST was derived. It will be appreciated that there may be several extended cDNAs which include the EST sequence as a result of alternate splicing or the activity of alternative promoters.

In the past, these short EST sequences were often obtained from oligo-dT primed cDNA libraries. Accordingly, they mainly corresponded to the 3' untranslated region of the  
15 mRNA. In part, the prevalence of EST sequences derived from the 3' end of the mRNA is a result of the fact that typical techniques for obtaining cDNAs are not well suited for isolating cDNA sequences derived from the 5' ends of mRNAs. (Adams *et al.*, *Nature* 377:3-174, 1996; Hillier *et al.*, *Genome Res.* 6:807-828, 1996).

In addition, in those reported instances where longer cDNA sequences have been  
20 obtained, the reported sequences typically correspond to coding sequences and do not include the full 5' untranslated region of the mRNA from which the cDNA is derived. Such incomplete sequences may not include the first exon of the mRNA, particularly in situations where the first exon is short. Furthermore, they may not include some exons, often short ones, which are located upstream of splicing sites. Thus, there is a need to obtain sequences  
25 derived from the 5' ends of mRNAs.

While many sequences derived from human chromosomes have practical applications, approaches based on the identification and characterization of those chromosomal sequences which encode a protein product are particularly relevant to diagnostic and therapeutic uses. Of the 50,000-100,000 protein coding genes, those genes encoding proteins which are  
30 secreted from the cell in which they are synthesized, as well as the secreted proteins themselves, are particularly valuable as potential therapeutic agents. Such proteins are often

involved in cell to cell communication and may be responsible for producing a clinically relevant response in their target cells.

In fact, several secretory proteins, including tissue plasminogen activator, G-CSF, GM-CSF, erythropoietin, human growth hormone, insulin, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$ , and interleukin-2, are currently in clinical use. These proteins are used to treat a wide range of conditions, including acute myocardial infarction, acute ischemic stroke, anemia, diabetes, growth hormone deficiency, hepatitis, kidney carcinoma, chemotherapy induced neutropenia and multiple sclerosis. For these reasons, extended cDNAs encoding secreted proteins or portions thereof represent a particularly valuable source of therapeutic agents. Thus, there is a need for the identification and characterization of secreted proteins and the nucleic acids encoding them.

In addition to being therapeutically useful themselves, secretory proteins include short peptides, called signal peptides, at their amino termini which direct their secretion. These signal peptides are encoded by the signal sequences located at the 5' ends of the coding sequences of genes encoding secreted proteins. Because these signal peptides will direct the extracellular secretion of any protein to which they are operably linked, the signal sequences may be exploited to direct the efficient secretion of any protein by operably linking the signal sequences to a gene encoding the protein for which secretion is desired. In addition, portions of signal sequences may also be used to direct the intracellular import of a peptide or protein of interest. This may prove beneficial in gene therapy strategies in which it is desired to deliver a particular gene product to cells other than the cell in which it is produced. Signal sequences encoding signal peptides also find application in simplifying protein purification techniques. In such applications, the extracellular secretion of the desired protein greatly facilitates purification by reducing the number of undesired proteins from which the desired protein must be selected. Thus, there exists a need to identify and characterize the 5' portions of the genes for secretory proteins which encode signal peptides.

Public information on the number of human genes for which the promoters and upstream regulatory regions have been identified and characterized is quite limited. In part this may be due to the difficulty of isolating such regulatory sequences. Upstream regulatory sequences such as transcription factor binding sites are typically too short to be utilized as probes for isolating promoters from human genomic libraries. Recently, some approaches

have been developed to isolate human promoters. One of them consists of making a CpG island library (Cross, *et al.*, *Nature Genetics* 6: 236-244, 1994). The second consists of isolating human genomic DNA sequences containing SpeI binding sites by the use of SpeI binding protein. (Mortlock *et al.*, *Genome Res.* 6:327-335, 1996). Both of these approaches  
5 have their limits due to a lack of specificity or of comprehensiveness.

The present 5' ESTs may be used to efficiently identify and isolate upstream regulatory regions which control the location, developmental stage, rate, and quantity of protein synthesis, as well as the stability of the mRNA. (Theil, *BioFactors* 4:87-93, 1993). Once identified and characterized, these regulatory regions may be utilized in gene therapy or  
10 protein purification schemes to obtain the desired amount and locations of protein synthesis or to inhibit, reduce, or prevent the synthesis of undesirable gene products.

In addition, ESTs containing the 5' ends of secretory protein genes may include sequences useful as probes for chromosome mapping and the identification of individuals. Thus, there is a need to identify and characterize the sequences upstream of the 5' coding  
15 sequences of genes encoding secretory proteins.

#### Summary of the Invention

The present invention relates to purified, isolated, or recombinant ESTs which include sequences derived from the authentic 5' ends of their corresponding mRNAs. The term  
20 "corresponding mRNA" refers to the mRNA which was the template for the cDNA synthesis which produced the 5' EST. These sequences will be referred to hereinafter as "5' ESTs." As used herein, the term "purified" does not require absolute purity; rather, it is intended as a relative definition. Individual 5' EST clones isolated from a cDNA library have been conventionally purified to electrophoretic homogeneity. The sequences obtained from these  
25 clones could not be obtained directly either from the library or from total human DNA. The cDNA clones are not naturally occurring as such, but rather are obtained via manipulation of a partially purified naturally occurring substance (messenger RNA). The conversion of mRNA into a cDNA library involves the creation of a synthetic substance (cDNA) and pure individual cDNA clones can be isolated from the synthetic library by clonal selection. Thus,  
30 creating a cDNA library from messenger RNA and subsequently isolating individual clones from that library results in an approximately  $10^4$ - $10^6$  fold purification of the native message.

Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

As used herein, the term "isolated" requires that the material be removed from its original environment (c.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide present in a living animal is not isolated, but the same polynucleotide, separated from some or all of the coexisting materials in the natural system, is isolated.

As used herein, the term "recombinant" means that the 5' EST is adjacent to "backbone" nucleic acid to which it is not adjacent in its natural environment. Additionally, to be "enriched" the 5' ESTs will represent 5% or more of the number of nucleic acid inserts in a population of nucleic acid backbone molecules. Backbone molecules according to the present invention include nucleic acids such as expression vectors, self-replicating nucleic acids, viruses, integrating nucleic acids, and other vectors or nucleic acids used to maintain or manipulate a nucleic acid insert of interest. Preferably, the enriched 5' ESTs represent 15% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. More preferably, the enriched 5' ESTs represent 50% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules. In a highly preferred embodiment, the enriched 5' ESTs represent 90% or more of the number of nucleic acid inserts in the population of recombinant backbone molecules.

"Stringent", moderate, and "low" hybridization conditions are as defined in Example 29.

Unless otherwise indicated, a "complementary" sequence is fully complementary.

Thus, 5' ESTs in cDNA libraries in which one or more 5' ESTs make up 5% or more of the number of nucleic acid inserts in the backbone molecules are "enriched recombinant 5' ESTs" as defined herein. Likewise, 5' ESTs in a population of plasmids in which one or more 5' EST of the present invention have been inserted such that they represent 5% or more of the number of inserts in the plasmid backbone are "enriched recombinant 5' ESTs" as defined herein. However, 5' ESTs in cDNA libraries in which 5' ESTs constitute less than 5% of the number of nucleic acid inserts in the population of backbone molecules, such as libraries in

which backbone molecules having a 5' EST insert are extremely rare, are not "enriched recombinant 5' ESTs".

In particular, the present invention relates to 5' ESTs which are derived from genes encoding secreted proteins. As used herein, a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal peptides in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g. soluble proteins), or partially (e.g. receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins which are transported across the membrane of the endoplasmic reticulum.

Such 5' ESTs include nucleic acid sequences, called signal sequences, which encode signal peptides which direct the extracellular secretion of the proteins encoded by the genes from which the 5' ESTs are derived. Generally, the signal peptides are located at the amino termini of secreted proteins.

Secreted proteins are translated by ribosomes associated with the "rough" endoplasmic reticulum. Generally, secreted proteins are co-translationally transferred to the membrane of the endoplasmic reticulum. Association of the ribosome with the endoplasmic reticulum during translation of secreted proteins is mediated by the signal peptide. The signal peptide is typically cleaved following its co-translational entry into the endoplasmic reticulum. After delivery to the endoplasmic reticulum, secreted proteins may proceed through the Golgi apparatus. In the Golgi apparatus, the proteins may undergo post-translational modification before entering secretory vesicles which transport them across the cell membrane.

The 5' ESTs of the present invention have several important applications. For example, they may be used to obtain and express cDNA clones which include the full protein coding sequences of the corresponding gene products, including the authentic translation start sites derived from the 5' ends of the coding sequences of the mRNAs from which the 5' ESTs are derived. These cDNAs will be referred to hereinafter as "full length cDNAs." These cDNAs may also include DNA derived from mRNA sequences upstream of the translation start site. The full length cDNA sequences may be used to express the proteins corresponding to the 5' ESTs. As discussed above, secreted proteins are therapeutically important. Thus, the proteins expressed from the cDNAs may be useful in treating or

controlling a variety of human conditions. The 5' ESTs may also be used to obtain the corresponding genomic DNA. The term "corresponding genomic DNA" refers to the genomic DNA which encodes the mRNA from which the 5' EST was derived.

Alternatively, the 5' ESTs may be used to obtain and express extended cDNAs  
5 encoding portions of the secreted protein. The portions may comprise the signal peptides of the secreted proteins or the mature proteins generated when the signal peptide is cleaved off. The portions may also comprise polypeptides having at least 10 consecutive amino acids encoded by the extended cDNAs or full length cDNAs. Alternatively, the portions may  
10 comprise at least 15 consecutive amino acids encoded by the extended cDNAs or full length cDNAs. In some embodiments, the portions may comprise at least 25 consecutive amino acids encoded by the extended cDNAs or full length cDNAs. In other embodiments, the portions may comprise at least 40 amino acids encoded by the extended cDNAs or full length cDNAs.

Antibodies which specifically recognize the entire secreted proteins encoded by the  
15 extended cDNAs, full length cDNAs, or fragments thereof having at least 10 consecutive amino acids, at least 15 consecutive amino acids, at least 25 consecutive amino acids, or at least 40 consecutive amino acids may also be obtained as described below. Antibodies which specifically recognize the mature protein generated when the signal peptide is cleaved may also be obtained as described below. Similarly, antibodies which specifically recognize the  
20 signal peptides encoded by the extended cDNAs or full length cDNAs may also be obtained.

In some embodiments, the extended cDNAs obtained using the 5' ESTs include the signal sequence. In other embodiments, the extended cDNAs obtained using the 5' ESTs may include the full coding sequence for the mature protein (*i.e.* the protein generated when the signal polypeptide is cleaved off). In addition, the extended cDNAs obtained using the 5'  
25 ESTs may include regulatory regions upstream of the translation start site or downstream of the stop codon which control the amount, location, or developmental stage of gene expression.

As discussed above, secreted proteins are therapeutically important. Thus, the proteins expressed from the extended cDNAs or full length cDNAs obtained using the 5'  
30 ESTs may be useful in treating or controlling a variety of human conditions.

The 5' ESTs (or cDNAs or genomic DNAs obtained therefrom) may be used in forensic procedures to identify individuals or in diagnostic procedures to identify individuals having genetic diseases resulting from abnormal expression of the genes corresponding to the 5' ESTs. In addition, the present invention is useful for constructing a high resolution map of the human chromosomes.

The present invention also relates to secretion vectors capable of directing the secretion of a protein of interest. Such vectors may be used in gene therapy strategies in which it is desired to produce a gene product in one cell which is to be delivered to another location in the body. Secretion vectors may also facilitate the purification of desired proteins.

The present invention also relates to expression vectors capable of directing the expression of an inserted gene in a desired spatial or temporal manner or at a desired level. Such vectors may include sequences upstream of the 5' ESTs, such as promoters or upstream regulatory sequences.

Finally, the present invention may also be used for gene therapy to control or treat genetic diseases. Signal peptides may also be fused to heterologous proteins to direct their extracellular secretion.

Bacterial clones containing Bluescript plasmids having inserts containing the 5' ESTs of the present invention (SEQ ID NOs: 38-270 are presently stored at 80°C in 4% (v/v) glycerol in the inventor's laboratories under the designations listed next to the SEQ ID NOs in

II). The inserts may be recovered from the deposited materials by growing the appropriate clones on a suitable medium. The Bluescript DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a cesium chloride gradient, size exclusion chromatography, or anion exchange chromatography. The plasmid DNA obtained using these procedures may then be manipulated using standard cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the EST insertion. The PCR product which corresponds to the 5' EST can then be manipulated using standard cloning techniques familiar to those skilled in the art.



One aspect of the present invention is a purified or isolated nucleic acid having the sequence of one of SEQ ID NOs: 38-270 or having a sequence complementary thereto. In one embodiment, the nucleic acid is recombinant.

Another aspect of the present invention is a purified or isolated nucleic acid  
5 comprising at least 10 consecutive bases of the sequence of one of SEQ ID NOs: 38-270 or one of the sequences complementary thereto.

Yet another aspect of the present invention is a purified or isolated nucleic acid comprising at least 15 consecutive bases of one of the sequences of SEQ ID NOs: 38-270 or one of the sequences complementary thereto. In one embodiment, the nucleic acid is  
10 recombinant.

A further aspect of the present invention is a purified or isolated nucleic acid of at least 15 bases capable of hybridizing under stringent conditions to the sequence of one of SEQ ID NOs: 38-270 or one of the sequences complementary to the sequences of SEQ ID NOs: 38-270. In one embodiment, the nucleic acid is recombinant.

15 Another aspect of the present invention is a purified or isolated nucleic acid encoding a human gene product, said human gene product having a sequence partially encoded by one of the sequences of SEQ ID NO: 38-270.

Still another aspect of the present invention is a method of making a cDNA encoding a human secretory protein, said human secretory protein being partially encoded by one of  
20 SEQ ID NOs 38-270, comprising the steps of contacting a collection of mRNA molecules from human cells with a primer comprising at least 15 consecutive nucleotides of a sequence complementary to one of SEQ ID NOs: 38-270; hybridizing said primer to an mRNA in said collection that encodes said protein; reverse transcribing said hybridized primer to make a first cDNA strand from said mRNA; making a second cDNA strand complementary to said first  
25 cDNA strand; and isolating the resulting cDNA encoding said protein comprising said first cDNA strand and said second cDNA strand.

Another aspect of the invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270 or a fragment thereof of at least 10 amino acids, said cDNA being  
30 obtainable by the method described in the preceding paragraph. In one embodiment, the

cDNA comprises the full protein coding sequence of said protein which sequence is partially included in one of the sequences of SEQ ID NOs: 38-270.

Another aspect of the present invention is a method of making a cDNA encoding a human secretory protein that is partially encoded by one of SEQ ID NOs 38-270, comprising  
5 the steps of obtaining a cDNA comprising one of the sequences of SEQ ID NOs: 38-270; contacting said cDNA with a detectable probe comprising at least 15 consecutive nucleotides of said sequence of SEQ ID NO: 38-270 or a sequence complementary thereto under conditions which permit said probe to hybridize to said cDNA; identifying a cDNA which hybridizes to said detectable probe; and isolating said cDNA which hybridizes to said probe.

10 Another aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the  
15 sequences of SEQ ID NOs: 38-270.

Another aspect of the present invention is a method of making a cDNA comprising one of the sequence of SEQ ID NOs: 38-270, comprising the steps of contacting a collection of mRNA molecules from human cells with a first primer capable of hybridizing to the polyA  
20 tail of said mRNA; hybridizing said first primer to said polyA tail; reverse transcribing said mRNA to make a first cDNA strand; making a second cDNA strand complementary to said first cDNA strand using at least one primer comprising at least 15 nucleotides of one of the sequences of SEQ ID NOs 38-270; and isolating the resulting cDNA comprising said first cDNA strand and said second cDNA strand.

Another aspect of the present invention is an isolated or purified cDNA encoding a  
25 human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

30 In one embodiment of the method described in the two paragraphs above, the second cDNA strand is made by contacting said first cDNA strand with a first pair of primers, said

first pair of primers comprising a second primer comprising at least 15 consecutive nucleotides of one of the sequences of SEQ ID NOs 38-270 and a third primer having a sequence therein which is included within the sequence of said first primer; performing a first polymerase chain reaction with said first pair of nested primers to generate a first PCR product; contacting said first PCR product with a second pair of primers, said second pair of primers comprising a fourth primer, said fourth primer comprising at least 15 consecutive nucleotides of said sequence of one of SEQ ID NOs: 38-270, and a fifth primer, said fourth and fifth primers being capable of hybridizing to sequences within said first PCR product; and performing a second polymerase chain reaction, thereby generating a second PCR product.

One aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270, or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

Another aspect of the present invention is the method described four paragraphs above in which the second cDNA strand is made by contacting said first cDNA strand with a second primer comprising at least 15 consecutive nucleotides of the sequences of SEQ ID NOs: 38-270; hybridizing said second primer to said first strand cDNA; and extending said hybridized second primer to generate said second cDNA strand.

Another aspect of the present invention is an isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein partially encoded by one of SEQ ID NOs 38-270 or comprising a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method described in the preceding paragraph. In one embodiment, the cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

Another aspect of the present invention is a method of making a protein comprising one of the sequences of SEQ ID NOs: 271-503, comprising the steps of obtaining a cDNA encoding the full protein sequence partially included in one of the sequences of sequence of SEQ ID NOs: 38-270; inserting said cDNA in an expression vector such that said cDNA is

operably linked to a promoter; introducing said expression vector into a host cell whereby said host cell produces the protein encoded by said cDNA; and isolating said protein.

Another aspect of the present invention is an isolated protein obtainable by the method described in the preceding paragraph.

5 Another aspect of the present invention is a method of obtaining a promoter DNA comprising the steps of obtaining DNAs located upstream of the nucleic acids of SEQ ID NOs: 38-270 or the sequences complementary thereto; screening said upstream DNAs to identify a promoter capable of directing transcription initiation; and isolating said DNA comprising said identified promoter. In one embodiment, the obtaining step comprises  
10 chromosome walking from said nucleic acids of SEQ ID NOs: 38-270 or sequences complementary thereto. In another embodiment, the screening step comprises inserting said upstream sequences into a promoter reporter vector. In another embodiment, the screening step comprises identifying motifs in said upstream DNAs which are transcription factor binding sites or transcription start sites.

15 Another aspect of the present invention is an isolated promoter obtainable by the method described above.

Another aspect of the present invention is an isolated or purified protein comprising one of the sequences of SEQ ID NOs: 271-503.

20 Another aspect of the present invention is the inclusion of at least one of the sequences of SEQ ID NOs: 38-270, or one of the sequences complementary to the sequences of SEQ ID NOs: 38-270, or a fragment thereof of at least 15 consecutive nucleotides in an array of discrete ESTs or fragments thereof of at least 15 nucleotides in length. In one embodiment, the array includes at least two of the sequences of SEQ ID NOs: 38-270, the sequences complementary to the sequences of SEQ ID NOs: 38-270, or fragments thereof of  
25 at least 15 consecutive nucleotides. In another embodiment, the array includes at least five of the sequences of SEQ ID NOs: 38-270, the sequences complementary to the sequences of SEQ ID NOs: 38-270, or fragments thereof of at least 15 consecutive nucleotides.

Another aspect of the present invention is a promoter having a sequence selected from the group consisting of SEQ ID NOs 31, 34, and 37.

### Brief Description of the Drawings

Figure 1 is a summary of a procedure for obtaining cDNAs which have been selected to include the 5' ends of the mRNAs from which they derived.

Figure 2 shows the distribution of Von Heijne scores for 5' ESTs in each of the categories described herein and the probability that these 5' ESTs encode a signal peptide.

Figure 3 summarizes a general method used to clone and sequence extended cDNAs containing sequences adjacent to 5' ESTs.

Figure 4 (description of promoters structure isolated from SignalTag 5' ESTs) provides a schematic description of promoters isolated and the way they are assembled with the corresponding 5' tags.

### Detailed Description of the Preferred Embodiment

Table IV is an analysis of the 43 amino acids located at the N terminus of all human SwissProt proteins to determine the frequency of false positives and false negatives using the techniques for signal peptide identification described herein.

Table V shows the distribution of 5' ESTs in each category described herein and the number of 5' ESTs in each category having a given minimum Von Heijne's score.

Table VI shows the distribution of 5' ESTs in each category described herein with respect to the tissue from which the 5' ESTs of the corresponding mRNA were obtained.

Table VII describes the transcription factor binding sites present in each of these promoters.

### I. General Methods for Obtaining 5' ESTs derived from mRNAs with intact 5' ends

In order to obtain the 5' ESTs of the present invention, mRNAs with intact 5' ends must be obtained. Currently, there are two approaches for obtaining such mRNAs with intact 5' ends as described below: either chemical (1) or enzymatic (2).

#### 1. Chemical Methods for Obtaining mRNAs having Intact 5' Ends

One of these approaches is a chemical modification method involving derivatization of the 5' ends of the mRNAs and selection of the derivatized mRNAs. The 5' ends of eukaryotic mRNAs possess a structure referred to as a "cap" which comprises a guanosine

5 methylated at the 7 position. The cap is joined to the first transcribed base of the mRNA by a 5', 5'-triphosphate bond. In some instances, the 5' guanosine is methylated in both the 2 and 7 positions. Rarely, the 5' guanosine is trimethylated at the 2, 7 and 7 positions. In the chemical method for obtaining mRNAs having intact 5' ends, the 5' cap is specifically derivatized and coupled to a reactive group on an immobilizing substrate. This specific derivatization is based on the fact that only the ribose linked to the methylated guanosine at the 5' end of the mRNA and the ribose linked to the base at the 3' terminus of the mRNA, possess 2', 3'-cis diols.

10 Optionally, the 2', 3'-cis diol of the 3' terminal ribose may be chemically modified, substituted, converted, or eliminated, leaving only the ribose linked to the methylated guanosine at the 5' end of the mRNA with a 2', 3'-cis diol. A variety of techniques are available for eliminating the 2', 3'-cis diol on the 3' terminal ribose. For example, controlled alkaline hydrolysis may be used to generate mRNA fragments in which the 3' terminal ribose is a 3'-phosphate, 2'-phosphate or (2', 3')-cyclophosphate. 15 Thereafter, the fragment which includes the original 3' ribose may be eliminated from the mixture through chromatography on an oligodT column. Alternatively, a base which lacks the 2', 3'-cis diol may be added to the 3' end of the mRNA using an RNA ligase such as T4 RNA ligase. Example 1 below describes a method for ligation of a nucleoside diphosphate to the 3' end of messenger RNA.

20

### EXAMPLE 1

#### Ligation of the Nucleoside Diphosphate pCp to the 3' End of mRNA.

One  $\mu$ g of RNA was incubated in a final reaction medium of 10  $\mu$ l in the presence of 5 U of T<sub>4</sub> phage RNA ligase in the buffer provided by the manufacturer (Gibco - 25 BRL), 40 U of the RNase inhibitor RNasin (Promega) and, 2  $\mu$ l of <sup>32</sup>pCp (Amersham #PB 10208). The incubation was performed at 37°C for 2 hours or overnight at 7-8°C.

Following modification or elimination of the 2', 3'-cis diol at the 3' ribose, the 2', 3'-cis diol present at the 5' end of the mRNA may be oxidized using reagents such as 30 NaBH<sub>4</sub>, NaBH<sub>3</sub>CN, or sodium periodate, thereby converting the 2', 3'-cis diol to a

dialdehyde. Example 2 describes the oxidation of the 2', 3'-cis diol at the 5' end of the mRNA with sodium periodate.

## EXAMPLE 2

### 5      Oxidation of 2', 3'-cis diol at the 5' End of the mRNA with Sodium Periodate

0.1 OD unit of either a capped oligoribonucleotide of 47 nucleotides (including the cap) or an uncapped oligoribonucleotide of 46 nucleotides were treated as follows. The oligoribonucleotides were produced by *in vitro* transcription using the transcription kit "AmpliScribe T7" (Epicentre Technologies). As indicated below, the DNA template for the  
10 RNA transcript contained a single cytosine. To synthesize the uncapped RNA, all four NTPs were included in the *in vitro* transcription reaction. To obtain the capped RNA, GTP was replaced by an analogue of the cap, m7G(5')ppp(5')G. This compound, recognized by the polymerase, was incorporated into the 5' end of the nascent transcript during the initiation of transcription but was not incorporated during the extension step. Consequently, the resulting  
15 RNA contained a cap at its 5' end. The sequences of the oligoribonucleotides produced by the *in vitro* transcription reaction were:

+Cap:

5'm7GpppGCAUCCUACUCCCAUCCAAUUCCACCCUAACUCCUCCCAUCUCCAC-  
3' (SEQ ID NO:1)

20    -Cap:

5'-pppGCAUCCUACUCCCAUCCAAUUCCACCCUAACUCCUCCCAUCUCCAC-3'  
(SEQ ID NO:2)

The oligoribonucleotides were dissolved in 9 µl of acetate buffer (0.1 M sodium acetate, pH 5.2) and 3 µl of freshly prepared 0.1 M sodium periodate solution. The mixture  
25 was incubated for 1 hour in the dark at 4°C or room temperature. Thereafter, the reaction was stopped by adding 4 µl of 10% ethylene glycol. The product was ethanol precipitated, resuspended in at least 10 µl of water or appropriate buffer and dialyzed against water.

The resulting aldehyde groups may then be coupled to molecules having a reactive  
30 amine group, such as hydrazine, carbazide, thiocarbazide or semicarbazide groups, in order to facilitate enrichment of the 5' ends of the mRNAs. Molecules having reactive amine groups

which are suitable for use in selecting mRNAs having intact 5' ends include avidin, proteins, antibodies, vitamins, ligands capable of specifically binding to receptor molecules, or oligonucleotides. Example 3 below describes the coupling of the resulting dialdehyde to biotin.


5

### EXAMPLE 3

#### Coupling of the Dialdehyde at the 5' End of Transcripts with Biotin

The oxidation product obtained in Example 2 was dissolved in 50  $\mu$ l of sodium acetate at a pH between 5 and 5.2 and 50  $\mu$ l of freshly prepared 0.02 M solution of biotin hydrazide in a methoxyethanol/water mixture (1:1) of formula:

10



In the compound used in these experiments,  $n=5$ . However, it will be appreciated that other commercially available hydrazides may also be used, such as molecules of the above formula in which  $n$  varies from 0 to 5. The mixture was then incubated for 2 hours at 37°C, precipitated with ethanol and dialyzed against distilled water. Example 4 demonstrates the specificity of the biotinylation reaction.

15

### EXAMPLE 4

#### Specificity of Biotinylation of Capped Transcripts

The specificity of the biotinylation for capped mRNAs was evaluated by gel electrophoresis of the following samples:

20

Sample 1. The 46 nucleotide uncapped *in vitro* transcript prepared as in Example 2 and labeled with  $^{32}\text{pCp}$  as described in Example 1.

Sample 2. The 46 nucleotide uncapped *in vitro* transcript prepared as in Example 2, labeled with  $^{32}\text{pCp}$  as described in Example 1, treated with the oxidation reaction of Example 2, and subjected to the biotinylation conditions of Example 3.

25



Sample 3. The 47 nucleotide capped *in vitro* transcript prepared as in Example 2 and labeled with <sup>32</sup>pCp as described in Example 1.

Sample 4. The 47 nucleotide capped *in vitro* transcript prepared as in Example 2, labeled with <sup>32</sup>pCp as described in Example 1, treated with the oxidation reaction of Example 2, and subjected to the biotinylation conditions of Example 3.

Samples 1 and 2 had identical migration rates, demonstrating that the uncapped RNAs were not oxidized and biotinylated. Sample 3 migrated more slowly than Samples 1 and 2, while Sample 4 exhibited the slowest migration. The difference in migration of the RNAs in Samples 3 and 4 demonstrates that the capped RNAs were specifically biotinylated.

10

In some cases, mRNAs having intact 5' ends may be enriched by binding the molecule containing a reactive amine group to a suitable solid phase substrate such as the inside of the vessel containing the mRNAs, magnetic beads, chromatography matrices, or nylon or nitrocellulose membranes. For example, where the molecule having a reactive amine group is biotin, the solid phase substrate may be coupled to avidin or streptavidin. Alternatively, where the molecule having the reactive amine group is an antibody or receptor ligand, the solid phase substrate may be coupled to the cognate antigen or receptor. Finally, where the molecule having a reactive amine group comprises an oligonucleotide, the solid phase substrate may comprise a complementary oligonucleotide.

The mRNAs having intact 5' ends may be released from the solid phase following the enrichment procedure. For example, where the dialdehyde is coupled to biotin hydrazide and the solid phase comprises streptavidin, the mRNAs may be released from the solid phase by simply heating to 95 degrees Celsius in 2% SDS. In some methods, the molecule having a reactive amine group may also be cleaved from the mRNAs having intact 5' ends following enrichment. Example 5 describes the capture of biotinylated mRNAs with streptavidin coated beads and the release of the biotinylated mRNAs from the beads following enrichment.

25

### EXAMPLE 5

#### Capture and Release of Biotinylated mRNAs Using Streptavidin Coated Beads

The streptavidin coated magnetic beads were prepared according to the manufacturer's instructions (CPG Inc., USA) The biotinylated mRNAs were added to a

30

hybridization buffer (1.5 M NaCl, pH 5 - 6). After incubating for 30 minutes, the unbound and nonbiotinylated material was removed. The beads were then washed several times in water with 1% SDS. The beads thus obtained were incubated for 15 minutes at 95°C in water containing 2% SDS.

- 5        Example 6 demonstrates the efficiency with which biotinylated mRNAs were recovered from the streptavidin coated beads.

### EXAMPLE 6

#### Efficiency of Recovery of Biotinylated mRNAs

- 10        The efficiency of the recovery procedure was evaluated as follows. Capped RNAs were labeled with <sup>32</sup>pCp, oxidized, biotinylated and bound to streptavidin coated beads as described above. Subsequently, the bound RNAs were incubated for 5, 15 or 30 minutes at 95°C in the presence of 2% SDS.

- 15        The products of the reaction were analyzed by electrophoresis on 12% polyacrylamide gels under denaturing conditions (7 M urea). The gels were subjected to autoradiography. During this manipulation, the hydrazone bonds were not reduced.

Increasing amounts of nucleic acids were recovered as incubation times in 2% SDS increased, demonstrating that biotinylated mRNAs were efficiently recovered.

- 20        In an alternative method for obtaining mRNAs having intact 5' ends, an oligonucleotide which has been derivatized to contain a reactive amine group is specifically coupled to mRNAs having an intact cap. Preferably, the 3' end of the mRNA is blocked prior to the step in which the aldehyde groups are joined to the derivatized oligonucleotide, as described above, so as to prevent the derivatized oligonucleotide from being joined to the 3' end of the mRNA. For example, pCp may be attached to the 3' end of the mRNA using T4 RNA ligase as described in example 1. However, as discussed above, blocking the 3' end of the mRNA is an optional step. Derivatized oligonucleotides may be prepared as described in Example 7.
- 25

**EXAMPLE 7****Derivatization of Oligonucleotides**

An oligonucleotide phosphorylated at its 3' end was converted to a 3' hydrazide in 3' by treatment with an aqueous solution of hydrazine or of dihydrazide of the formula  
5  $H_2N(R_1)NH_2$  at about 1 to 3 M, and at pH 4.5 at a temperature of 8°C overnight. This incubation was performed in the presence of a carbodiimide type agent soluble in water such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide at a final concentration of 0.3 M.

The derivatized oligonucleotide was then separated from the other agents and products using a standard technique for isolating oligonucleotides.

10 As discussed above, the mRNAs to be enriched may be treated to eliminate the 3' OH groups which may be present thereon. This may be accomplished by enzymatic ligation of sequences lacking a 3' OH, such as pCp, as described in Example 1. Alternatively, the 3' OH groups may be eliminated by alkaline hydrolysis as described in Example 8 below.

15

**EXAMPLE 8****Elimination of 3' OH Groups of mRNA Using Alkaline Hydrolysis**

In a total volume of 100  $\mu$ l of 0.1 N sodium hydroxide, 1.5  $\mu$ g mRNA is incubated for 40 to 60 minutes at 4°C. The solution is neutralized with acetic acid and precipitated with ethanol.

20 Following the optional elimination of the 3' OH groups, the diol groups at the 5' ends of the mRNAs are oxidized as described below in Example 9.

**EXAMPLE 9****Oxidation of Diols of mRNA**

25 Up to 1 OD unit of RNA was dissolved in 9  $\mu$ l of buffer (0.1 M sodium acetate, pH 6-7) or water and 3  $\mu$ l of freshly prepared 0.1 M sodium periodate solution. The reaction was incubated for 1 h in the dark at 4°C or room temperature. Following the incubation, the reaction was stopped by adding 4  $\mu$ l of 10% ethylene glycol. Thereafter the mixture was incubated at room temperature for 15 minutes. After ethanol precipitation, the product was  
30 resuspended in at least 10  $\mu$ l of water or appropriate buffer and dialyzed against water.

Following oxidation of the diol groups at the 5' ends of the mRNAs, the derivatized oligonucleotide was joined to the resulting aldehydes as described in Example 10.

#### EXAMPLE 10

##### 5        Ligature of Aldehydes of mRNA to Derivatized Oligonucleotides

The oxidized mRNA was dissolved in an acidic medium such as 50  $\mu$ l of sodium acetate pH 4-6. Fifty  $\mu$ l of a solution of the derivatized oligonucleotide were added in order to obtain an mRNA:derivatized oligonucleotide ratio of 1:20. The mixture was reduced with a borohydride and incubated for 2 h at 37°C or overnight (14 h) at 10°C. The mixture was  
10 then ethanol precipitated, resuspended in 10  $\mu$ l or more of water or appropriate buffer and dialyzed against distilled water. If desired, the resulting product may be analyzed using acrylamide gel electrophoresis, HPLC analysis, or other conventional techniques.

Following the attachment of the derivatized oligonucleotide to the mRNAs, a  
15 reverse transcription reaction may be performed as described in Example 11 below.

#### EXAMPLE 11

##### Reverse Transcription of mRNAs Ligatured to Derivatized Oligonucleotides

An oligodeoxyribonucleotide was derivatized as follows. Three OD units of an  
20 oligodeoxyribonucleotide of sequence 5'ATCAAGAATTCGCACGAGACCATTAA3' (SEQ ID NO:3) having 5'-OH and 3'-P ends were dissolved in 70  $\mu$ l of a 1.5 M hydroxybenzotriazole solution, pH 5.3, prepared in dimethylformamide/water (75:25) containing 2  $\mu$ g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide. The mixture was incubated for 2 h 30 min at 22°C and then precipitated twice in LiClO<sub>4</sub>/acetone. The pellet  
25 was resuspended in 200  $\mu$ l of 0.25 M hydrazine and incubated at 8°C from 3 to 14 h. Following the hydrazine reaction, the mixture was precipitated twice in LiClO<sub>4</sub>/acetone.

The messenger RNAs to be reverse transcribed were extracted from blocks of placenta having sides of 2 cm which had been stored at -80°C. The total RNA was extracted using conventional acidic phenol techniques. Oligo-dT chromatography was used to purify  
30 the mRNAs. The integrity of the mRNAs was checked by Northern-blotting.

The diol groups on 7  $\mu$ g of the placental mRNAs were oxidized as described above in Example 9. The derivatized oligonucleotide was joined to the mRNAs as described in Example 10 above except that the precipitation step was replaced by an exclusion chromatography step to remove derivatized oligodeoxyribonucleotides which were not joined to mRNAs. Exclusion chromatography was performed as follows:

Ten ml of Ultrogel AcA34 (BioSeptra#230151) gel, a mix of agarose and acrylamide, were equilibrated in 50 ml of a solution of 10 mM Tris pH 8.0, 300 mM NaCl, 1 mM EDTA, and 0.05% SDS. The mixture was allowed to sediment. The supernatant was eliminated and the gel was resuspended in 50 ml of buffer. This procedure was repeated 2 or 3 times.

10 A glass bead (diameter 3 mm) was introduced into a 2 ml disposable pipette (length 25 cm). The pipette was filled with the gel suspension until the height of the gel stabilized at 1 cm from the top of the pipette. The column was then equilibrated with 20 ml of equilibration buffer (10 mM Tris HCl pH 7.4, 20 mM NaCl).

Ten  $\mu$ l of the mRNA which had reacted with the derivatized oligonucleotide were mixed in 39  $\mu$ l of 10 mM urea and 2  $\mu$ l of blue-glycerol buffer, which had been prepared by dissolving 5 mg of bromophenol blue in 60% glycerol (v/v), and passing the mixture through a 0.45  $\mu$ m diameter filter.

The column was then loaded with the mRNAs coupled to the oligonucleotide. As soon as the sample had penetrated, equilibration buffer was added. Hundred  $\mu$ l fractions were then collected. Derivatized oligonucleotide which had not been attached to mRNA appeared in fraction 16 and later fractions. Thus, fractions 3 to 15 were combined and precipitated with ethanol.

To determine whether the derivatized oligonucleotide was actually linked to mRNA, one tenth of the combined fractions were spotted twice on a nylon membrane and hybridized to a radioactive probe using conventional techniques. The  $^{32}$ P labeled probe used in these hybridizations was an oligodeoxyribonucleotide of sequence 5'TAATGGTCTCGTGCGAATTCTTGAT3' (SEQ ID NO:4) anticomplementary to the derivatized oligonucleotide. A signal observed after autoradiography, indicated that the derivatized oligonucleotide had been truly joined to the mRNA.

30 The remaining nine tenths of the mRNAs which had reacted with the derivatized oligonucleotide was reverse transcribed as follows. A reverse transcription reaction was

carried out with reverse transcriptase following the manufacturer's instructions and 50 pmol of nonamers with random sequence as primers.

To ensure that reverse transcription had been carried out through the cap structure, two types of experiments were performed.

5 In the first approach, after elimination of RNA of the cDNA:RNA heteroduplexes obtained from the reverse transcription reaction by an alkaline hydrolysis, a portion of the resulting single stranded cDNAs was spotted on a positively charged membrane and hybridized, using conventional methods, to a <sup>32</sup>P labeled probe having a sequence identical to that of the derivatized oligonucleotide. Control spots containing, 1 pmol, 100 fmol, 50 fmol,  
10 10 fmol and 1 fmol of a control oligodeoxyribonucleotide of sequence identical to that of the derivatized oligonucleotide were included. The signal observed in the spots containing the cDNA indicated that approximately 15 fmol of the derivatized oligonucleotide had been reverse transcribed. These results demonstrate that the reverse transcription can be performed through the cap and, in particular, that reverse transcriptase crosses the 5'-P-P-P-  
15 5' bond of the cap of eukaryotic messenger RNAs.

In the second type of experiment, the single stranded cDNAs obtained from the above first strand synthesis were used as template for PCR reactions. Two types of reactions were carried out. First, specific amplification of the mRNAs for alpha globin, dehydrogenase, pp15 and elongation factor E4 were carried out using the following pairs of  
20 oligodeoxyribonucleotide primers.

alpha-globin

GLO-S: 5'CCG ACA AGA CCA ACG TCA AGG CCG C3' (SEQ ID NO:5)

GLO-As: 5'TCA CCA GCA GGC AGT GGC TTA GGA G 3' (SEQ ID NO:6)

25

dehydrogenase

3 DH-S: 5'AGT GAT TCC TGC TAC TTT GGA TGG C3' (SEQ ID NO:7)

3 DH-As: 5'GCT TGG TCT TGT TCT GGA GTT TAG A3' (SEQ ID NO:8)

30

pp15

PP15-S: 5'TCC AGA ATG GGA GAC AAG CCA ATT T3' (SEQ ID NO:9)

PP15-As: 5'AGG GAG GAG GAA ACA GCG TGA GTC C3' (SEQ ID NO:10)

Elongation factor E4

EFA1-S: 5'ATG GGA AAG GAA AAG ACT CAT ATC A3' (SEQ ID NO:11)

5 EF1A-As: 5'AGC AGC AAC AAT CAG GAC AGC ACA G3' (SEQ ID NO:12)

Second, non specific amplifications were also carried out with the antisense oligodeoxyribonucleotides of the pairs described above and with a primer derived from the sequence of the derivatized oligodeoxyribonucleotide  
10 (5'ATCAAGAATTCGCACGAGACCATT3') (SEQ ID NO:13).

One twentieth of the following RT-PCR product samples were run on a 1.5% agarose gel and stained with ethidium bromide.

Sample 1: The products of a PCR reaction using the globin primers of SEQ ID NOs 5 and 6 in the presence of cDNA.

15 Sample 2: The products of a PCR reaction using the globin primers of SEQ ID NOs 5 and 6 in the absence of added cDNA.

Sample 3: The products of a PCR reaction using the dehydrogenase primers of SEQ ID NOs 7 and 8 in the presence of cDNA.

20 Sample 4: The products of a PCR reaction using the dehydrogenase primers of SEQ ID NOs 7 and 8 in the absence of added cDNA.

Sample 5: The products of a PCR reaction using the pp15 primers of SEQ ID NOs 9 and 10 in the presence of cDNA.

Sample 6: The products of a PCR reaction using the pp15 primers of SEQ ID NOs 9 and 10 in the absence of added cDNA.

25 Sample 7: The products of a PCR reaction using the EIF4 primers of SEQ ID NOs 11 and 12 in the presence of added cDNA.

Sample 8: The products of a PCR reaction using the EIF4 primers of SEQ ID NOs 11 and 12 in the absence of added cDNA.

30 A band of the size expected for the PCR product was observed only in samples 1, 3, 5 and 7, thus indicating the presence of the corresponding sequence in the cDNA population

PCR reactions were also carried out with the antisense oligonucleotides of the globin and dehydrogenase primers (SEQ ID NOs 6 and 8) and an oligonucleotide whose sequence corresponds to that of the derivatized oligonucleotide. The presence of PCR products of the expected size in the samples equivalent to above samples 1 and 3 indicated that the  
5 derivatized oligonucleotide had been linked to mRNA.

The above examples summarize the chemical procedure for enriching mRNAs for those having intact 5' ends as illustrated in Figure 1. Further detail regarding the chemical approaches for obtaining such mRNAs are disclosed in International Application No.  
10 WO96/34981, published November 7, 1996, which is incorporated herein by reference. Strategies based on the above chemical modifications to the 5' cap structure may be utilized to generate cDNAs selected to include the 5' ends of the mRNAs from which they derived. In one version of such procedures, the 5' ends of the mRNAs are modified as described above. Thereafter, a reverse transcription reaction is conducted to extend a primer  
15 complementary to the 5' end of the mRNA. Single stranded RNAs are eliminated to obtain a population of cDNA/mRNA heteroduplexes in which the mRNA includes an intact 5' end. The resulting heteroduplexes may be captured on a solid phase coated with a molecule capable of interacting with the molecule used to derivatize the 5' end of the mRNA. Thereafter, the strands of the heteroduplexes are separated to recover single stranded first  
20 cDNA strands which include the 5' end of the mRNA. Second strand cDNA synthesis may then proceed using conventional techniques. For example, the procedures disclosed in WO 96/34981 or in Caminci. *et al.*, *Genomics* 37:327-336, 1996, the disclosures of which are incorporated herein by reference, may be employed to select cDNAs which include the sequence derived from the 5' end of the coding sequence of the mRNA.

25 Following ligation of the oligonucleotide tag to the 5' cap of the mRNA, a reverse transcription reaction is conducted to extend a primer complementary to the mRNA to the 5' end of the mRNA. Following elimination of the RNA component of the resulting heteroduplex using standard techniques, second strand cDNA synthesis is conducted with a primer complementary to the oligonucleotide tag.



## 2. Enzymatic Methods for Obtaining mRNAs having Intact 5' Ends

Other techniques for selecting cDNAs extending to the 5' end of the mRNA from which they are derived are fully enzymatic. Some versions of these techniques are disclosed in Dumas Milne Edwards J.B. (Doctoral Thesis of Paris VI University, Le clonage des ADNc complets: difficultes et perspectives nouvelles. Apports pour l'etude de la regulation de l'expression de la tryptophane hydroxylase de rat, 20 Dec. 1993), EP0 625572 and Kato *et al.*, *Gene* 150:243-250, 1994, the disclosures of which are incorporated herein by reference.

Briefly, in such approaches, isolated mRNA is treated with alkaline phosphatase to remove the phosphate groups present on the 5' ends of uncapped incomplete mRNAs.

Following this procedure, the cap present on full length mRNAs is enzymatically removed with a decapping enzyme such as T4 polynucleotide kinase or tobacco acid pyrophosphatase. An oligonucleotide, which may be either a DNA oligonucleotide or a DNA-RNA hybrid oligonucleotide having RNA at its 3' end, is then ligated to the phosphate present at the 5' end of the decapped mRNA using T4 RNA ligase. The oligonucleotide may include a restriction site to facilitate cloning of the cDNAs following their synthesis. Example 12 below describes one enzymatic method based on the doctoral thesis of Dumas.

### EXAMPLE 12

#### Enzymatic Approach for Obtaining 5' ESTs

Twenty micrograms of PolyA+ RNA were dephosphorylated using Calf Intestinal Phosphatase (Biolabs). After a phenol chloroform extraction, the cap structure of mRNA was hydrolysed using the Tobacco Acid Pyrophosphatase (purified as described by Shinshi *et al.*, *Biochemistry* 15: 2185-2190, 1976) and a hemi 5'DNA/RNA-3' oligonucleotide having an unphosphorylated 5' end, a stretch of adenosine ribophosphate at the 3' end, and an EcoRI site near the 5' end was ligated to the 5'P ends of mRNA using the T4 RNA ligase (Biolabs). Oligonucleotides suitable for use in this procedure are preferably 30 to 50 bases in length. Oligonucleotides having an unphosphorylated 5' end may be synthesized by adding a fluorochrome at the 5' end. The inclusion of a stretch of adenosine ribophosphates at the 3' end of the oligonucleotide increases ligation efficiency. It will be appreciated that the oligonucleotide may contain cloning sites other than EcoRI.

Following ligation of the oligonucleotide to the phosphate present at the 5' end of the decapped mRNA, first and second strand cDNA synthesis is carried out using conventional methods or those specified in EP0 625,572 and Kato *et al. supra*, and Dumas Milne Edwards, *supra*, the disclosures of which are incorporated herein by reference. The resulting cDNA may then be ligated into vectors such as those disclosed

5 in Kato *et al. supra* or other nucleic acid vectors known to those skilled in the art using techniques such as those described in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual 2d Ed., Cold Spring Harbor Laboratory Press, 1989, the disclosure of which is incorporated herein by reference.

10

## II. Obtention and Characterization of the 5' ESTs of the Present Invention

The 5' ESTs of the present invention were obtained using the aforementioned chemical and enzymatic approaches for enriching mRNAs for those having intact 5' ends as described below.

15

### 1. Obtention of 5' ESTS Using mRNAs with Intact 5' Ends

First, mRNAs were prepared as described in Example 13 below.

## EXAMPLE 13

20

### Preparation of mRNA With Intact 5' Ends

Total human RNAs or polyA<sup>+</sup> RNAs derived from 29 different tissues were respectively purchased from LABIMO and CLONTECH and used to generate 44 cDNA libraries as follows. The purchased RNA had been isolated from cells or tissues using acid guanidium thiocyanate-phenol-chloroform extraction (Chomczynski and Sacchi, *Analytical Biochemistry* 162:156-159, 1987). PolyA<sup>+</sup> RNA was isolated from total RNA (LABIMO)

25 by two passes of oligo dT chromatography, as described by Aviv and Leder, *Proc. Natl. Acad. Sci. USA* 69:1408-1412, 1972 in order to eliminate ribosomal RNA.

30

The quality and the integrity of the polyA<sup>+</sup> RNAs were checked. Northern blots hybridized with a globin probe were used to confirm that the mRNAs were not degraded. Contamination of the polyA<sup>+</sup> mRNAs by ribosomal sequences was checked using Northern blots and a probe derived from the sequence of the 28S rRNA. Preparations of mRNAs with

less than 5% of rRNAs were used in library construction. To avoid constructing libraries with RNAs contaminated by exogenous sequences (prokaryotic or fungal), the presence of bacterial 16S ribosomal sequences or of two highly expressed fungal mRNAs was examined using PCR.

5        Following preparation of the mRNAs, the above described chemical and/or the enzymatic procedures for enriching mRNAs for those having intact 5' ends were employed to obtain 5' ESTs from various tissues. In both approaches, an oligonucleotide tag was attached to the 5' ends of the mRNAs. The oligonucleotide tag had an EcoRI site therein to facilitate later cloning procedures. To facilitate the processing of single stranded and double  
10        stranded cDNA obtained in the construction of the libraries, the same nucleotidic sequence was used to design the ligated oligonucleotide in both chemical and enzymatic approaches. Nevertheless, in the chemical procedure, the tag used was an oligodeoxyribonucleotide which was linked to the cap of the mRNA whereas in the enzymatic ligation, the tag was a chimeric  
15        hemi 5'DNA/RNA3' oligonucleotide which was ligated to the 5' end of decapped mRNA as described in example 12.

      Following attachment of the oligonucleotide tag to the mRNA by either the chemical or enzymatic methods, the integrity of the mRNA was examined by performing a Northern blot with 200 to 500 ng of mRNA using a probe complementary to the oligonucleotide tag before performing the first strand synthesis as described in example 14.

20

#### EXAMPLE 14

##### cDNA Synthesis Using mRNA Templates Having Intact 5' Ends

      For the mRNAs joined to oligonucleotide tags using both the chemical and enzymatic methods, first strand cDNA synthesis was performed using the Superscript II (Gibco BRL) or  
25        the Rnase H Minus M-MLV (Promega) reverse transcriptase with random nonamers as primers. In order to protect internal EcoRI sites in the cDNA from digestion at later steps in the procedure, methylated dCTP was used for first strand synthesis. After removal of RNA by an alkaline hydrolysis, the first strand of cDNA was precipitated using isopropanol in order to eliminate residual primers.

30        For both the chemical and the enzymatic methods, the second strand of the cDNA was synthesized with a Klenow fragment using a primer corresponding to the 5' end of the

ligated oligonucleotide described in Example 12. Preferably, the primer is 20-25 bases in length. Methylated dCTP was also used for second strand synthesis in order to protect internal EcoRI sites in the cDNA from digestion during the cloning process.

Following cDNA synthesis, the cDNAs were cloned into pBlueScript as described in  
5 Example 15 below.

### EXAMPLE 15

#### Cloning of cDNAs derived from mRNA with intact 5' ends into BlueScript

Following second strand synthesis, the ends of the cDNA were blunted with T4 DNA  
10 polymerase (Biolabs) and the cDNA was digested with EcoRI. Since methylated dCTP was used during cDNA synthesis, the EcoRI site present in the tag was the only hemi-methylated site, hence the only site susceptible to EcoRI digestion. The cDNA was then size fractionated using exclusion chromatography (AcA, Biosepra) and fractions corresponding to cDNAs of  
15 more than 150 bp were pooled and ethanol precipitated. The cDNA was directionally cloned into the SmaI and EcoRI ends of the phagemid pBlueScript vector (Stratagene). The ligation mixture was electroporated into bacteria and propagated under appropriate antibiotic selection.

Clones containing the oligonucleotide tag attached were then selected as described in  
Example 16 below.

20

### EXAMPLE 16

#### Selection of Clones Having the Oligonucleotide Tag Attached Thereto

The plasmid DNAs containing 5' EST libraries made as described above were purified (Qiagen). A positive selection of the tagged clones was performed as follows  
25 Briefly, in this selection procedure, the plasmid DNA was converted to single stranded DNA using gene II endonuclease of the phage F1 in combination with an exonuclease (Chang *et al.* *Gene* 127:95-8, 1993) such as exonuclease III or T7 gene 6 exonuclease. The resulting single stranded DNA was then purified using paramagnetic beads as described by Fry *et al.* *Biotechniques*, 13: 124-131, 1992. In this procedure, the single stranded DNA was  
30 hybridized with a biotinylated oligonucleotide having a sequence corresponding to the 3' end of the oligonucleotide described in Example 13. Preferably, the primer has a length of 20-

bases. Clones including a sequence complementary to the biotinylated oligonucleotide were captured by incubation with streptavidin coated magnetic beads followed by magnetic selection. After capture of the positive clones, the plasmid DNA was released from the magnetic beads and converted into double stranded DNA using a DNA polymerase such as the ThermoSequenase obtained from Amersham Pharmacia Biotech. Alternatively, protocols such as the one described in the Gene Trapper kit available from Gibco BRL may be used. The double stranded DNA was then electroporated into bacteria. The percentage of positive clones having the 5' tag oligonucleotide was estimated to typically rank between 90 and 98% using dot blot analysis.

Following electroporation, the libraries were ordered in 384-microtiter plates (MTP). A copy of the MTP was stored for future needs. Then the libraries were transferred into 96 MTP and sequenced as described below.

#### EXAMPLE 17

##### Sequencing of Inserts in Selected Clones

Plasmid inserts were first amplified by PCR on PE 9600 thermocyclers (Perkin-Elmer, Applied Biosystems Division, Foster City, CA), using standard SETA-A and SETA-B primers (Genset SA), AmpliTaqGold (Perkin-Elmer), dNTPs (Boehringer), buffer and cycling conditions as recommended by the Perkin-Elmer Corporation.

PCR products were then sequenced using automatic ABI Prism 377 sequencers (Perkin Elmer). Sequencing reactions were performed using PE 9600 thermocyclers with standard dye-primer chemistry and ThermoSequenase (Amersham Pharmacia Biotech). The primers used were either T7 or 21M13 (available from Genset SA) as appropriate. The primers were labeled with the JOE, FAM, ROX and TAMRA dyes. The dNTPs and ddNTPs used in the sequencing reactions were purchased from Boehringer. Sequencing buffer, reagent concentrations and cycling conditions were as recommended by Amersham.

Following the sequencing reaction, the samples were precipitated with ethanol, resuspended in formamide loading buffer, and loaded on a standard 4% acrylamide gel. Electrophoresis was performed for 2.5 hours at 3000V on an ABI 377 sequencer, and the sequence data were collected and analyzed using the ABI Prism DNA Sequencing Analysis Software, version 2.1.2.

## 2. Computer analysis of the Obtained 5' ESTs: Construction of NetGene and SignalTag databases

The sequence data from the 44 cDNA libraries made as described above were transferred to a proprietary database, where quality control and validation steps were performed. A proprietary base-caller, working using a Unix system, automatically flagged suspect peaks, taking into account the shape of the peaks, the inter-peak resolution, and the noise level. The proprietary base-caller also performed an automatic trimming. Any stretch of 25 or fewer bases having more than 4 suspect peaks was considered unreliable and was discarded. Sequences corresponding to cloning vector or ligation oligonucleotides were automatically removed from the EST sequences. However, the resulting EST sequences may contain 1 to 5 bases belonging to the above mentioned sequences at their 5' end. If needed, these can easily be removed on a case to case basis.

Following sequencing as described above, the sequences of the 5' ESTs were entered in NetGene™, a proprietary database called for storage and manipulation as described below. It will be appreciated by those skilled in the art that the data could be stored and manipulated on any medium which can be read and accessed by a computer. Computer readable media include magnetically, optically, or electronically readable media. For example, the computer readable media may be a hard disc, a floppy disc, a magnetic tape, CD-ROM, RAM, or ROM as well as other types of other media known to those skilled in the art.

In addition, the sequence data may be stored and manipulated in a variety of data processor programs in a diversity of formats. For instance, the sequence data may be stored as text in a word processing file, such as Microsoft WORD or WORDPERFECT or as an ASCII file in a variety of database programs familiar to those of skill in the art, such as DB2, SYBASE, or ORACLE.

The computer readable media on which the sequence information is stored may be in a personal computer, a network, a server or other computer systems known to those skilled in the art. The computer or other system preferably includes the storage media described above, and a processor for accessing and manipulating the sequence data. Once the sequence data has been stored, it may be manipulated and searched to locate those stored sequences which contain a desired nucleic acid sequence or which encode a protein having a particular functional domain. For example, the stored sequence information may be compared to other

known sequences to identify homologies, motifs implicated in biological function, or structural motifs.

Programs which may be used to search or compare the stored sequences include the MacPattern (EMBL), BLAST, and BLAST2 program series (NCBI), basic local alignment  
5 search tool programs for nucleotide (BLASTN) and peptide (BLASTX) comparisons (Altschul *et al*, *J. Mol. Biol.* 215: 403, 1990) and FASTA (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85: 2444, 1988). The BLAST programs then extend the alignments on the basis of defined match and mismatch criteria.

Motifs which may be detected using the above programs and those described in  
10 Example 28 include sequences encoding leucine zippers, helix-turn-helix motifs, glycosylation sites, ubiquitination sites, alpha helices, and beta sheets, signal sequences encoding signal peptides which direct the secretion of the encoded proteins, sequences implicated in transcription regulation such as homeoboxes, acidic stretches, enzymatic active sites, substrate binding sites, and enzymatic cleavage sites.

15 Before searching the cDNAs in the NetGene™ database for sequence motifs of interest, cDNAs derived from mRNAs which were not of interest were identified and eliminated from further consideration as described in Example 18 below.

### EXAMPLE 18

#### 20 Elimination of Undesired Sequences from Further Consideration

5' ESTs in the NetGene™ database which were derived from undesired sequences such as transfer RNAs, ribosomal RNAs, mitochondrial RNAs, prokaryotic RNAs, fungal RNAs, Alu sequences, L1 sequences, or repeat sequences were identified using the FASTA and BLASTN programs with the parameters listed in Table I.

25 To eliminate 5' ESTs encoding tRNAs from further consideration, the 5' EST sequences were compared to the sequences of 1190 known tRNAs obtained from EMBL release 38, of which 100 were human. The comparison was performed using FASTA on both strands of the 5' ESTs. Sequences having more than 80% homology over more than 60 nucleotides were identified as tRNA. Of the 144,341 sequences screened, 26 were identified  
30 as tRNAs and eliminated from further consideration.

To eliminate 5' ESTs encoding rRNAs from further consideration, the 5' EST sequences were compared to the sequences of 2497 known rRNAs obtained from EMBL release 38, of which 73 were human. The comparison was performed using BLASTN on both strands of the 5' ESTs with the parameter S=108. Sequences having more than 80%  
5 homology over stretches longer than 40 nucleotides were identified as rRNAs. Of the 144,341 sequences screened, 3,312 were identified as rRNAs and eliminated from further consideration.

To eliminate 5' ESTs encoding mtRNAs from further consideration, the 5' EST sequences were compared to the sequences of the two known mitochondrial genomes for  
10 which the entire genomic sequences are available and all sequences transcribed from these mitochondrial genomes including tRNAs, rRNAs, and mRNAs for a total of 38 sequences. The comparison was performed using BLASTN on both strands of the 5' ESTs with the parameter S=108. Sequences having more than 80% homology over stretches longer than 40 nucleotides were identified as mtRNAs. Of the 144,341 sequences screened, 6,110 were  
15 identified as mtRNAs and eliminated from further consideration.

Sequences which might have resulted from exogenous contaminants were eliminated from further consideration by comparing the 5' EST sequences to release 46 of the EMBL bacterial and fungal divisions using BLASTN with the parameter S=144. All sequences having more than 90% homology over at least 40 nucleotides were identified as exogenous  
20 contaminants. Of the 42 cDNA libraries examined, the average percentages of prokaryotic and fungal sequences contained therein were 0.2% and 0.5% respectively. Among these sequences, only one could be identified as a sequence specific to fungi. The others were either fungal or prokaryotic sequences having homologies with vertebrate sequences or including repeat sequences which had not been masked during the electronic comparison.

25 In addition, the 5' ESTs were compared to 6093 Alu sequences and 1115 L1 sequences to mask 5' ESTs containing such repeat sequences. 5' ESTs including THE and MER repeats, SSTR sequences or satellite, micro-satellite, or telomeric repeats were also eliminated from further consideration. On average, 11.5% of the sequences in the libraries contained repeat sequences. Of this 11.5%, 7% contained Alu repeats, 3.3% contained L1  
30 repeats and the remaining 1.2% were derived from the other screened types of repetitive sequences. These percentages are consistent with those found in cDNA libraries prepared by



other groups. For example, the cDNA libraries of Adams *et al.* contained between 0% and 7.4% Alu repeats depending on the source of the RNA which was used to prepare the cDNA library (Adams *et al.*, *Nature* 377:174, 1996).

- 5           The sequences of those 5' ESTs remaining after the elimination of undesirable sequences were compared with the sequences of known human mRNAs to determine the accuracy of the sequencing procedures described above.

#### EXAMPLE 19

10           Measurement of Sequencing Accuracy by Comparison to Known Sequences

To further determine the accuracy of the sequencing procedure described above, the sequences of 5' ESTs derived from known sequences were identified and compared to the original known sequences. First, a FASTA analysis with overhangs shorter than 5 bp on both ends was conducted on the 5' ESTs to identify those matching an entry in the public human mRNA database. The 6655 5' ESTs which matched a known human mRNA were then  
15           realigned with their cognate mRNA and dynamic programming was used to include substitutions, insertions, and deletions in the list of "errors" which would be recognized. Errors occurring in the last 10 bases of the 5' EST sequences were ignored to avoid the inclusion of spurious cloning sites in the analysis of sequencing accuracy.

- 20           This analysis revealed that the sequences incorporated in the NetGene™ database had an accuracy of more than 99.5%.

To determine the efficiency with which the above selection procedures select cDNAs which include the 5' ends of their corresponding mRNAs, the following analysis  
25           was performed.

#### EXAMPLE 20

Determination of Efficiency of 5' EST Selection

- To determine the efficiency at which the above selection procedures isolated 5' ESTs  
30           which included sequences close to the 5' end of the mRNAs from which they derived, the sequences of the ends of the 5' ESTs derived from the elongation factor 1 subunit  $\alpha$  are:

ferritin heavy chain genes were compared to the known cDNA sequences of these genes. Since the transcription start sites of both genes are well characterized, they may be used to determine the percentage of derived 5' ESTs which included the authentic transcription start sites.

5 For both genes, more than 95% of the obtained 5' ESTs actually included sequences close to or upstream of the 5' end of the corresponding mRNAs.

To extend the analysis of the reliability of the procedures for isolating 5' ESTs from ESTs in the NetGene™ database, a similar analysis was conducted using a database composed of human mRNA sequences extracted from GenBank database release 97 for  
10 comparison. The 5' ends of more than 85% of 5' ESTs derived from mRNAs included in the GeneBank database were located close to the 5' ends of the known sequence. As some of the mRNA sequences available in the GenBank database are deduced from genomic sequences, a 5' end matching with these sequences will be counted as an internal match. Thus, the method used here underestimates the yield of ESTs including the authentic 5' ends  
15 of their corresponding mRNAs.

The EST libraries made above included multiple 5' ESTs derived from the same mRNA. The sequences of such 5' ESTs were compared to one another and the longest 5' ESTs for each mRNA were identified. Overlapping cDNAs were assembled into  
20 continuous sequences (contigs). The resulting continuous sequences were then compared to public databases to gauge their similarity to known sequences, as described in Example 21 below.

#### EXAMPLE 21

##### 25 Clustering of the 5' ESTs and Calculation of Novelty Indices for cDNA Libraries

For each sequenced EST library, the sequences were clustered by the 5' end. Each sequence in the library was compared to the others with BLASTN2 (direct strand, parameters S=107). ESTs with High Scoring Segment Pairs (HSPs) at least 25 bp long, having 95% identical bases and beginning closer than 10 bp from each EST 5' end were grouped. The  
30 longest sequence found in the cluster was used as representative of the group. A global clustering between libraries was then performed leading to the definition of super-contigs.

To assess the yield of new sequences within the EST libraries, a novelty rate (NR) was defined as:  $NR = 100 \times (\text{Number of new unique sequences found in the library} / \text{Total number of sequences from the library})$ . Typically, novelty rating ranged between 10% and 41% depending on the tissue from which the EST library was obtained. For most of the  
5 libraries, the random sequencing of 5' EST libraries was pursued until the novelty rate reached 20%.

Following characterization as described above, the collection of 5' ESTs in NetGene™ was screened to identify those 5' ESTs bearing potential signal sequences as  
10 described in Example 22 below.

## EXAMPLE 22

### Identification of Potential Signal Sequences in 5' ESTs

The 5' ESTs in the NetGene™ database were screened to identify those having an  
15 uninterrupted open reading frame (ORF) longer than 45 nucleotides beginning with an ATG codon and extending to the end of the EST. Approximately half of the cDNA sequences in NetGene™ contained such an ORF. The ORFs of these 5' ESTs were then searched to identify potential signal motifs using slight modifications of the procedures disclosed in Von Heijne, *Nucleic Acids Res.* 14:4683-4690, 1986, the disclosure of which is incorporated  
20 herein by reference. Those 5' EST sequences encoding a stretch of at least 15 amino acid long with a score of at least 3.5 in the Von Heijne signal peptide identification matrix were considered to possess a signal sequence. Those 5' ESTs which matched a known human mRNA or EST sequence and had a 5' end more than 20 nucleotides downstream of the  
25 known 5' end were excluded from further analysis. The remaining cDNAs having signal sequences therein were included in a database called SignalTag™.

To confirm the accuracy of the above method for identifying signal sequences, the analysis of Example 23 was performed.

**EXAMPLE 23**Confirmation of Accuracy of Identification of Potential Signal Sequences in 5' ESTs

The accuracy of the above procedure for identifying signal sequences encoding signal peptides was evaluated by applying the method to the 43 amino acids located at the N terminus of all human SwissProt proteins. The computed Von Heijne score for each protein was compared with the known characterization of the protein as being a secreted protein or a non-secreted protein. In this manner, the number of non-secreted proteins having a score higher than 3.5 (false positives) and the number of secreted proteins having a score lower than 3.5 (false negatives) could be calculated.

Using the results of the above analysis, the probability that a peptide encoded by the 5' region of the mRNA is in fact a genuine signal peptide based on its Von Heijne's score was calculated based on either the assumption that 10% of human proteins are secreted or the assumption that 20% of human proteins are secreted. The results of this analysis are shown in Figure 2 and table IV.

Using the above method of identification of secretory proteins, 5' ESTs of the following polypeptides known to be secreted were obtained: human glucagon, gamma interferon induced monokine precursor, secreted cyclophilin-like protein, human pleiotropin, and human biotinidase precursor. Thus, the above method successfully identified those 5' ESTs which encode a signal peptide.

To confirm that the signal peptide encoded by the 5' ESTs actually functions as a signal peptide, the signal sequences from the 5' ESTs may be cloned into a vector designed for the identification of signal peptides. Such vectors are designed to confer the ability to grow in selective medium only to host cells containing a vector with an operably linked signal sequence. For example, to confirm that a 5' EST encodes a genuine signal peptide, the signal sequence of the 5' EST may be inserted upstream and in frame with a non-secreted form of the yeast invertase gene in signal peptide selection vectors such as those described in U.S. Patent No. 5,536,637, the disclosure of which is incorporated herein by reference. Growth of host cells containing signal sequence selection vectors with the correctly inserted 5' EST signal sequence confirms that the 5' EST encodes a genuine signal peptide.

Alternatively, the presence of a signal peptide may be confirmed by cloning the extended cDNAs obtained using the ESTs into expression vectors such as pXT1 (as described below in example 30), or by constructing promoter-signal sequence-reporter gene vectors which encode fusion proteins between the signal peptide and an assayable reporter protein. After introduction of these vectors into a suitable host cell, such as COS cells or NIH 3T3 cells, the growth medium may be harvested and analyzed for the presence of the secreted protein. The medium from these cells is compared to the medium from control cells containing vectors lacking the signal sequence or extended cDNA insert to identify vectors which encode a functional signal peptide or an authentic secreted protein.

Those 5' ESTs which encoded a signal peptide, as determined by the method of Example 22 above, were further grouped into four categories based on their homology to known sequences as described in Example 24 below.

#### EXAMPLE 24

##### Categorization of 5' ESTs Encoding a Signal Peptide

Those 5' ESTs having a sequence not matching any known vertebrate sequence nor any publicly available EST sequence were designated "new." Of the sequences in the SignalTag™ database, 947 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those 5' ESTs having a sequence not matching any vertebrate sequence but matching a publicly known EST were designated "EST-ext", provided that the known EST sequence was extended by at least 40 nucleotides in the 5' direction. Of the sequences in the SignalTag™ database, 150 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those ESTs not matching any vertebrate sequence but matching a publicly known EST without extending the known EST by at least 40 nucleotides in the 5' direction were designated "EST." Of the sequences in the SignalTag™ database, 599 of the 5' ESTs having a Von Heijne's score of at least 3.5 fell into this category.

Those 5' ESTs matching a human mRNA sequence but extending the known sequence by at least 40 nucleotides in the 5' direction were designated "VERT-ext." Of the sequences in the SignalTag™ database, 23 of the 5' ESTs having a Von Heijne's score of at

least 3.5 fell into this category. Included in this category was a 5' EST which extended the known sequence of the human translocase mRNA by more than 200 bases in the 5' direction. A 5' EST which extended the sequence of a human tumor suppressor gene in the 5' direction was also identified.

5 Table V shows the distribution of 5' ESTs in each category and the number of 5' ESTs in each category having a given minimum von Heijne's score.

### 3. Evaluation of Spatial and Temporal Expression of mRNAs Corresponding to the 5'ESTs or Extended cDNAs

10

Each of the 5' ESTs was also categorized based on the tissue from which its corresponding mRNA was obtained, as described below in Example 25.

#### **EXAMPLE 25**

15

##### Categorization of Expression Patterns

Table VI shows the distribution of 5' ESTs in each of the above defined category with respect to the tissue from which the 5'ESTs of the corresponding mRNA were obtained

20 Table II provides the sequence identification numbers of 5' EST sequences derived from testis and other tissues, the categories in which these sequences fall, and the von Heijne's score of the signal peptides which they encode. The 5' EST sequences and the amino acid sequences they encode are provided in the appended sequence listings. Table III provides the sequence ID numbers of the 5' ESTs and the sequences of the signal peptides which they encode. The sequences of the 5' ESTs and the polypeptides they encode are provided in the sequence listing appended hereto.

25 The sequences of DNA SEQ ID NOs: 38-270 can readily be screened for any errors therein and any sequence ambiguities can be resolved by resequencing a fragment containing such errors or ambiguities on both strands. Such fragments may be obtained from the plasmids stored in the inventors' laboratory or can be isolated using the techniques described herein. Resolution of any such ambiguities or errors may be facilitated by using primers  
30 which hybridize to sequences located close to the ambiguous or erroneous sequences. For example, the primers may hybridize to sequences within 50-75 bases of the ambiguity or

error. Upon resolution of an error or ambiguity, the corresponding corrections can be made in the protein sequences encoded by the DNA containing the error or ambiguity.

In addition to categorizing the 5' ESTs with respect to their tissue of origin, the spatial and temporal expression patterns of the mRNAs corresponding to the 5' ESTs, as well as their expression levels, may be determined as described in Example 26 below. Characterization of the spatial and temporal expression patterns and expression levels of these mRNAs is useful for constructing expression vectors capable of producing a desired level of gene product in a desired spatial or temporal manner, as will be discussed in more detail below.

Furthermore, 5' ESTs whose corresponding mRNAs are associated with disease states may also be identified. For example, a particular disease may result from the lack of expression, over expression, or under expression of an mRNA corresponding to a 5' EST. By comparing mRNA expression patterns and quantities in samples taken from healthy individuals with those from individuals suffering from a particular disease, 5' ESTs responsible for the disease may be identified.

It will be appreciated that the results of the above characterization procedures for 5' ESTs also apply to extended cDNAs (obtainable as described below) which contain sequences adjacent to the 5' ESTs. It will also be appreciated that if desired, characterization may be delayed until extended cDNAs have been obtained rather than characterizing the ESTs themselves.

## EXAMPLE 26

### Evaluation of Expression Levels and Patterns of mRNAs

#### Corresponding to 5' ESTs or Extended cDNAs

Expression levels and patterns of mRNAs corresponding to 5' ESTs or extended cDNAs (obtainable as described below in example 27) may be analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277, the entire contents of which are hereby incorporated by reference. Briefly, a 5' EST, extended cDNA, or fragment thereof corresponding to the gene encoding the mRNA to be characterized is inserted at a cloning site immediately downstream of a bacteriophage (T3,

T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the 5' EST or extended cDNA has 100 or more nucleotides. The plasmid is linearized and transcribed in the presence of ribonucleotides comprising modified ribonucleotides (*i.e.* biotin-UTP and DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated  
5 from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50°C for 16 hours in an 80% formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (*i.e.* RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence  
10 of the DIG modification enables the hybrid to be detected and quantified by ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

The 5' ESTs, extended cDNAs, or fragments thereof may also be tagged with nucleotide sequences for the serial analysis of gene expression (SAGE) as disclosed in UK Patent Application No. 2 305 241 A, the entire contents of which are incorporated by  
15 reference. In this method, cDNAs are prepared from a cell, tissue, organism or other source of nucleic acid for which gene expression patterns must be determined. The resulting cDNAs are separated into two pools. The cDNAs in each pool are cleaved with a first restriction endonuclease, called an anchoring enzyme, having a recognition site which is likely to be present at least once in most cDNAs. The fragments which contain the 5' or 3' most region  
20 of the cleaved cDNA are isolated by binding to a capture medium such as streptavidin coated beads. A first oligonucleotide linker having a first sequence for hybridization of an amplification primer and an internal restriction site for a so-called tagging endonuclease is ligated to the digested cDNAs in the first pool. Digestion with the second endonuclease produces short tag fragments from the cDNAs.

25 A second oligonucleotide having a second sequence for hybridization of an amplification primer and an internal restriction site is ligated to the digested cDNAs in the second pool. The cDNA fragments in the second pool are also digested with the tagging endonuclease to generate short tag fragments derived from the cDNAs in the second pool. The tags resulting from digestion of the first and second pools with the anchoring enzyme and  
30 the tagging endonuclease are ligated to one another to produce so-called ditags. In some embodiments, the ditags are concatamerized to produce ligation products containing from 2



to 200 ditags. The tag sequences are then determined and compared to the sequences of the 5' ESTs or extended cDNAs to determine which 5' ESTs or extended cDNAs are expressed in the cell, tissue, organism, or other source of nucleic acids from which the tags were derived. In this way, the expression pattern of the 5' ESTs or extended cDNAs in the cell,  
5 tissue, organism, or other source of nucleic acids is obtained.

Quantitative analysis of gene expression may also be performed using arrays. As used herein, the term array means a one dimensional, two dimensional, or multidimensional arrangement of full length cDNAs (*i.e.* extended cDNAs which include the coding sequence for the signal peptide, the coding sequence for the mature protein, and a stop codon),  
10 extended cDNAs, 5' ESTs or fragments thereof of sufficient length to permit specific detection of gene expression. Preferably, the fragments are at least 15 nucleotides in length. More preferably, the fragments are at least 100 nucleotide long. More preferably, the fragments are more than 100 nucleotides in length. In some embodiments, the fragments may be more than 500 nucleotide long.

For example, quantitative analysis of gene expression may be performed with full length cDNAs as defined below, extended cDNAs, 5' ESTs, or fragments thereof in a complementary DNA microarray as described by Schena *et al.* (*Science* 270:467-470, 1995; *Proc. Natl. Acad. Sci. U.S.A.* 93:10614-10619, 1996). Full length cDNAs, extended cDNAs, 5' ESTs or fragments thereof are amplified by PCR and arrayed from 96-well microtiter  
20 plates onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2 min at 95°C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25°C.

Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm<sup>2</sup> microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60°C. Arrays are washed for 5 min at 25°C in low stringency wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a  
30 fluorescence laser scanning device fitted with a custom filter set. Accurate differential

expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

Quantitative analysis of the expression of genes may also be performed with full length cDNAs, extended cDNAs, 5' ESTs, or fragments thereof in complementary DNA  
5 arrays as described by Pietu *et al.* (*Genome Research* 6:492-503, 1996). The full length cDNAs, extended cDNAs, 5' ESTs or fragments thereof are PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate  
10 experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis of the 5' ESTs or extended cDNAs can be done through high density nucleotide arrays as described by Lockhart *et al.* (*Nature Biotechnology* 14: 1675-1680, 1996) and Sosnowsky *et al.* (*Proc. Natl. Acad. Sci.* 94:1119-1123, 1997).  
15 Oligonucleotides of 15-50 nucleotides corresponding to sequences of the 5' ESTs or extended cDNAs are synthesized directly on the chip (Lockhart *et al.*, *supra*) or synthesized and then addressed to the chip (Sosnowsky *et al.*, *supra*). Preferably, the oligonucleotides are about 20 nucleotides in length.

cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin  
20 or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the chip. After washing as described in Lockhart *et al.*, *supra* and application of different electric fields (Sonowsky *et al.*, *supra.*), the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed.  
25 Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of the mRNA corresponding to the 5' EST or extended cDNA from which the oligonucleotide sequence has been designed.

### III. Use of 5' ESTs to Clone Extended cDNAs and to Clone the Corresponding Genomic DNAs

Once 5' ESTs which include the 5' end of the corresponding mRNAs have been selected using the procedures described above, they can be utilized to isolate extended cDNAs which contain sequences adjacent to the 5' ESTs. The extended cDNAs may include the entire coding sequence of the protein encoded by the corresponding mRNA, including the authentic translation start site, the signal sequence, and the sequence encoding the mature protein remaining after cleavage of the signal peptide. Such extended cDNAs are referred to herein as "full length cDNAs." Alternatively, the extended cDNAs may include only the sequence encoding the mature protein remaining after cleavage of the signal peptide, or only the sequence encoding the signal peptide.

Example 27 below describes a general method for obtaining extended cDNAs using 5' ESTs. Example 28 below provides experimental results, using the method explained in example 27, describing several extended cDNAs including the entire coding sequence and authentic 5' end of the corresponding mRNA for several secreted proteins.

The methods of Examples 27, 28, and 29 can also be used to obtain extended cDNAs which encode less than the entire coding sequence of the secreted proteins encoded by the genes corresponding to the 5' ESTs. In some embodiments, the extended cDNAs isolated using these methods encode at least 10 amino acids of one of the proteins encoded by the sequences of SEQ ID NOs: 38-270. In further embodiments, the extended cDNAs encode at least 20 amino acids of the proteins encoded by the sequences of SEQ ID NOs: 38-270. In further embodiments, the extended cDNAs encode at least 30 amino acids of the sequences of SEQ ID NOs: 38-270. In a preferred embodiment, the extended cDNAs encode a full length protein sequence, which includes the protein coding sequences of SEQ ID NOs: 38-270.

#### EXAMPLE 27

##### General Method for Using 5' ESTs to Clone and Sequence cDNAs which Include the Entire Coding Region and the Authentic 5' End of the Corresponding mRNA

The following general method has been used to quickly and efficiently isolate extended cDNAs having the authentic 5' ends of their corresponding mRNAs as well as

the full protein coding sequence and including sequence adjacent to the sequences of the 5' ESTs used to obtain them. This method may be applied to obtain extended cDNAs for any 5' EST in the NetGene™ database, including those 5' ESTs encoding polypeptides belonging to secreted proteins. The method is summarized in figure 3.

5

#### 1. Obtention of Extended cDNAs

##### *a) First strand synthesis*

The method takes advantage of the known 5' sequence of the mRNA. A reverse transcription reaction is conducted on purified mRNA with a poly 14dT primer containing a 49 nucleotide sequence at its 5' end allowing the addition of a known sequence at the end of the cDNA which corresponds to the 3' end of the mRNA. For example, the primer may have the following sequence: 5'-ATC GTT GAG ACT CGT ACC AGC AGA GTC ACG AGA GAG ACT ACA CGG TAC TGG TTT TTT TTT TTT TTVN -3' (SEQ ID NO:14). Those skilled in the art will appreciate that other sequences may also be added to the poly dT sequence and used to prime the first strand synthesis. Using this primer and a reverse transcriptase such as the Superscript II (Gibco BRL) or Rnase H Minus M-MLV (Promega) enzyme, a reverse transcript anchored at the 3' polyA site of the RNAs is generated.

15

20

After removal of the mRNA hybridized to the first cDNA strand by alkaline hydrolysis, the products of the alkaline hydrolysis and the residual poly dT primer are eliminated with an exclusion column such as an AcA34 (Biosepra) matrix as explained in Example 11.

##### *b) Second strand synthesis*

A pair of nested primers on each end is designed based on the known 5' sequence from the 5' EST and the known 3' end added by the poly dT primer used in the first strand synthesis. Softwares used to design primers are either based on GC content and melting temperatures of oligonucleotides, such as OSP (Illier and Green, *PCR Meth. Appl.* 1:124-128, 1991), or based on the octamer frequency disparity method (Griffais *et al.*, *Nucleic Acids Res.* 19: 3887-3891, 1991) such as PC-Rare (<http://bioinformatics.weizmann.ac.il/software/PC-Rare/doc/manuel.html>).

25

Preferably, the nested primers at the 5' end are separated from one another by four to nine bases. The 5' primer sequences may be selected to have melting temperatures and specificities suitable for use in PCR.

Preferably, the nested primers at the 3' end are separated from one another by four to nine bases. For example, the nested 3' primers may have the following sequences: (5'- CCA GCA GAG TCA CGA GAG AGA CTA CAC GG -3'(SEQ ID NO:15), and 5'- CAC GAG AGA GAC TAC ACG GTA CTG G -3' (SEQ ID NO:16). These primers were selected because they have melting temperatures and specificities compatible with their use in PCR. However, those skilled in the art will appreciate that other sequences may also be used as primers.

The first PCR run of 25 cycles is performed using the Advantage Tth Polymerase Mix (Clontech) and the outer primer from each of the nested pairs. A second 20 cycle PCR using the same enzyme and the inner primer from each of the nested pairs is then performed on 1/2500 of the first PCR product. Thereafter, the primers and nucleotides are removed.

## 2. Sequencing of Full Length Extended cDNAs or Fragments Thereof

Due to the lack of position constraints on the design of 5' nested primers compatible for PCR use using the OSP software, amplicons of two types are obtained. Preferably, the second 5' primer is located upstream of the translation initiation codon thus yielding a nested PCR product containing the whole coding sequence. Such a full length extended cDNA undergoes a direct cloning procedure as described in section a. However, in some cases, the second 5' primer is located downstream of the translation initiation codon, thereby yielding a PCR product containing only part of the ORF. Such incomplete PCR products are submitted to a modified procedure described in section b.

### *a) Nested PCR products containing complete ORFs*

When the resulting nested PCR product contains the complete coding sequence, as predicted from the 5'EST sequence, it is cloned in an appropriate vector such as pED6dpc2, as described in section 3.

*h) Nested PCR products containing incomplete ORFs*

When the amplicon does not contain the complete coding sequence, intermediate steps are necessary to obtain both the complete coding sequence and a PCR product containing the full coding sequence. The complete coding sequence can be assembled  
5 from several partial sequences determined directly from different PCR products as described in the following section.

Once the full coding sequence has been completely determined, new primers compatible for PCR use are designed to obtain amplicons containing the whole coding region. However, in such cases, 3' primers compatible for PCR use are located inside the  
10 3' UTR of the corresponding mRNA, thus yielding amplicons which lack part of this region, *i.e.* the polyA tract and sometimes the polyadenylation signal, as illustrated in figure 3. Such full length extended cDNAs are then cloned into an appropriate vector as described in section 3.

*c) Sequencing extended cDNAs*

15 Sequencing of extended cDNAs is performed using a Die Terminator approach with the AmpliTaq DNA polymerase FS kit available from Perkin Elmer.

In order to sequence PCR fragments, primer walking is performed using software such as OSP to choose primers and automated computer software such as ASMG (Sutton *et al.*, *Genome Science Technol.* 1: 9-19, 1995) to construct contigs of walking sequences  
20 including the initial 5' tag using minimum overlaps of 32 nucleotides. Preferably, primer walking is performed until the sequences of full length cDNAs are obtained.

Completion of the sequencing of a given extended cDNA fragment is assessed as follows. Since sequences located after a polyA tract are difficult to determine precisely in the case of uncloned products, sequencing and primer walking processes for PCR products are  
25 interrupted when a polyA tract is identified in extended cDNAs obtained as described in case b. The sequence length is compared to the size of the nested PCR product obtained as described above. Due to the limited accuracy of the determination of the PCR product size by gel electrophoresis, a sequence is considered complete if the size of the obtained sequence is at least 70 % the size of the first nested PCR product. If the length of the sequence  
30 determined from the computer analysis is not at least 70% of the length of the nested PCR product, these PCR products are cloned and the sequence of the insertion is determined.

When Northern blot data are available, the size of the mRNA detected for a given PCR product is used to finally assess that the sequence is complete. Sequences which do not fulfill the above criteria are discarded and will undergo a new isolation procedure.

Sequence data of all extended cDNAs are then transferred to a proprietary  
5 database, where quality controls and validation steps are carried out as described in example 15.

### 3. Cloning of Full Length Extended cDNAs

The PCR product containing the full coding sequence is then cloned in an appropriate  
10 vector. For example, the extended cDNAs can be cloned into the expression vector pED6dpc2 (DiscoverEase, Genetics Institute, Cambridge, MA) as follows. pED6dpc2 vector DNA is prepared with blunt ends by performing an EcoRI digestion followed by a fill in reaction. The blunt ended vector is dephosphorylated. After removal of PCR primers and ethanol precipitation, the PCR product containing the full coding sequence or the extended  
15 cDNA obtained as described above is phosphorylated with a kinase subsequently removed by phenol-Sevag extraction and precipitation. The double stranded extended cDNA is then ligated to the vector and the resulting expression plasmid introduced into appropriate host cells.

Since the PCR products obtained as described above are blunt ended molecules that  
20 can be cloned in either direction, the orientation of several clones for each PCR product is determined. Then, 4 to 10 clones are ordered in microtiter plates and subjected to a PCR reaction using a first primer located in the vector close to the cloning site and a second primer located in the portion of the extended cDNA corresponding to the 3' end of the mRNA. This second primer may be the antisense primer used in anchored PCR in the case of direct cloning  
25 (case a) or the antisense primer located inside the 3'UTR in the case of indirect cloning (case b). Clones in which the start codon of the extended cDNA is operably linked to the promoter in the vector so as to permit expression of the protein encoded by the extended cDNA are conserved and sequenced. In addition to the ends of cDNA inserts, approximately 50 bp of vector DNA on each side of the cDNA insert are also sequenced.

30 The cloned PCR products are then entirely sequenced according to the aforementioned procedure. In this case, contiguation of long fragments is then performed

on walking sequences that have already contigated for uncloned PCR products during primer walking. Sequencing of cloned amplicons is complete when the resulting contigs include the whole coding region as well as overlapping sequences with vector DNA on both ends.

5

#### 4. Computer analysis of Full Length Extended cDNA

Sequences of all full length extended cDNAs are then submitted to further analysis as described below. Before searching the extended full length cDNAs for sequences of interest, extended cDNAs which are not of interest (vector RNAs, transfer RNAs, ribosomal RNAs, mitochondrial RNAs, prokaryotic RNAs and fungal RNAs) are discarded using methods essentially similar to those described for 5'ESTs in Example 18.

##### *a) Identification of structural features*

Structural features, e.g. polyA tail and polyadenylation signal, of the sequences of full length extended cDNAs are subsequently determined as follows.

15 A polyA tail is defined as a homopolymeric stretch of at least 11 A with at most one alternative base within it. The polyA tail search is restricted to the last 100 nt of the sequence and limited to stretches of 11 consecutive A's because sequencing reactions are often not readable after such a polyA stretch. Stretches having more than 90% homology over 8 nucleotides are identified as polyA tails using BLAST2N.

20 To search for a polyadenylation signal, the polyA tail is clipped from the full-length sequence. The 50 bp preceding the polyA tail are first searched for the canonic polyadenylation AAUAAA signal and, if the canonic signal is not detected, for the alternative AUUAAA signal (Sheets *et al.*, *Nuc. Acids Res.* 18: 5799-5805, 1990). If neither of these consensus polyadenylation signals is found, the canonic motif is searched again allowing one mismatch to account for possible sequencing errors. More than 85 % of identified polyadenylation signals of either type actually ends 10 to 30 bp from the polyA tail. Alternative AUUAAA signals represents approximately 15 % of the total number of identified polyadenylation signals.

##### *b) Identification of functional features*

30 Functional features, e.g. ORFs and signal sequences, of the sequences of full length extended cDNAs were subsequently determined as follows.



The 3 upper strand frames of extended cDNAs are searched for ORFs defined as the maximum length fragments beginning with a translation initiation codon and ending with a stop codon. ORFs encoding at least 20 amino acids are preferred.

Each found ORF is then scanned for the presence of a signal peptide in the first  
5 50 amino-acids or, where appropriate, within shorter regions down to 20 amino acids or less in the ORF, using the matrix method of von Heijne (*Nuc. Acids Res.* 14: 4683-4690, 1986), the disclosure of which is incorporated herein by reference as described in Example 22.

*c) Homology to either nucleotidic or proteic sequences*

10 Categorization of full-length sequences may be achieved using procedures essentially similar to those described for 5'ESTs in Example 24.

Extended cDNAs prepared as described above may be subsequently engineered to obtain nucleic acids which include desired portions of the extended cDNA using conventional  
15 techniques such as subcloning, PCR, or *in vitro* oligonucleotide synthesis. For example, nucleic acids which include only the full coding sequences (*i.e.* the sequences encoding the signal peptide and the mature protein remaining after the signal peptide is cleaved off) may be obtained using techniques known to those skilled in the art. Alternatively, conventional techniques may be applied to obtain nucleic acids which contain only the coding sequences  
20 for the mature protein remaining after the signal peptide is cleaved off or nucleic acids which contain only the coding sequences for the signal peptides.

Similarly, nucleic acids containing any other desired portion of the coding sequences for the secreted protein may be obtained. For example, the nucleic acid may contain at least 10 consecutive bases of an extended cDNA such as one of the extended cDNAs described  
25 below. In another embodiment, the nucleic acid may contain at least 15 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. Alternatively, the nucleic acid may contain at least 20 consecutive bases of an extended cDNA such as one of the extended cDNAs described below. In another embodiment, the nucleic acid may contain at least 25 consecutive bases of an extended cDNA such as one of the extended cDNAs  
30 described below. In yet another embodiment, the nucleic acid may contain at least 40

consecutive bases of an extended cDNA such as one of the extended cDNAs described below.

Once an extended cDNA has been obtained, it can be sequenced to determine the amino acid sequence it encodes. Once the encoded amino acid sequence has been  
5 determined, one can create and identify any of the many conceivable cDNAs that will encode that protein by simply using the degeneracy of the genetic code. For example, allelic variants or other homologous nucleic acids can be identified as described below. Alternatively, nucleic acids encoding the desired amino acid sequence can be synthesized *in vitro*.

In a preferred embodiment, the coding sequence may be selected using the known  
10 codon or codon pair preferences for the host organism in which the cDNA is to be expressed.

The extended cDNAs derived from the 5' ESTS of the present invention were obtained as described in Example 28 below.

#### EXAMPLE 28

##### 15 Characterization of cloned extended cDNAs obtained using 5' ESTs

The procedure described in Example 27 above was used to obtain the extended cDNAs derived from the 5' ESTs of the present invention in a variety of tissues. The following list provides a few examples of thus obtained extended cDNAs.

Using this approach, the full length cDNA of SEQ ID NO:17 (internal identification  
20 number 48-19-3-G1-FL1) was obtained. This cDNA falls into the "EST-ext" category described above and encodes the signal peptide MKKVLLLITAILAVAVG (SEQ ID NO: 18) having a von Heijne score of 8.2.

The full length cDNA of SEQ ID NO:19 (internal identification number 58-34-2-E7-  
FL2) was also obtained using this procedure. This cDNA falls into the "EST-ext" category  
25 described above and encodes the signal peptide MWWFQQGLSFLPSALVTWTS (SEQ ID NO:20) having a von Heijne score of 5.5.

Another full length cDNA obtained using the procedure described above has the sequence of SEQ ID NO:21 (internal identification number 51-27-1-E8-FL1). This cDNA, falls into the "EST-ext" category described above and encodes the signal peptide  
30 MVLTTLPANSANSPPVNMPTTGPNLSYASSALSPCLT (SEQ ID NO:22) having a von Heijne score of 5.9.

The above procedure was also used to obtain a full length cDNA having the sequence of SEQ ID NO:23 (internal identification number 76-4-1-G5-FL1). This cDNA falls into the "EST-ext" category described above and encodes the signal peptide ILSTVTALTFAXA (SEQ ID NO:24) having a von Heijne score of 5.5.

5       The full length cDNA of SEQ ID NO:25 (internal identification number 51-3-3-B10-FL3) was also obtained using this procedure. This cDNA falls into the "new" category described above and encodes a signal peptide LVLTLCTLPLAVA (SEQ ID NO:26) having a von Heijne score of 10.1.

10       The full length cDNA of SEQ ID NO:27 (internal identification number 58-35-2-F10-FL2) was also obtained using this procedure. This cDNA falls into the "new" category described above and encodes a signal peptide LWLLFFLVTAIHA (SEQ ID NO:28) having a von Heijne score of 10.7.

Bacterial clones containing plasmids containing the full length cDNAs described above are presently stored in the inventor's laboratories under the internal identification numbers provided above. The inserts may be recovered from the stored materials by growing  
15       an aliquot of the appropriate bacterial clone in the appropriate medium. The plasmid DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a cesium chloride gradient,  
20       size exclusion chromatography, or anion exchange chromatography. The plasmid DNA obtained using these procedures may then be manipulated using standard cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the cDNA insertion. The PCR product which corresponds to the cDNA can then be manipulated using standard cloning techniques familiar to those skilled in the art.

25       The polypeptides encoded by the extended cDNAs may be screened for the presence of known structural or functional motifs or for the presence of signatures, small amino acid sequences which are well conserved amongst the members of a protein family. The conserved regions have been used to derive consensus patterns or matrices included in the PROSITE data bank, in particular in the file prosite.dat (Release 13.0 of November 1995,  
30       located at <http://expasy.hcuge.ch/sprot/prosite.html>. Prosite\_convert and prosite\_scan

programs ([http://ulrec3.unil.ch/ftpserveur/prosite\\_scan](http://ulrec3.unil.ch/ftpserveur/prosite_scan)) may be used to find signatures on the extended cDNAs.

For each pattern obtained with the `prosite_convert` program from the `prosite.dat` file, the accuracy of the detection on a new protein sequence may be assessed by evaluating the frequency of irrelevant hits on the population of human secreted proteins included in the data bank SWISSPROT. The ratio between the number of hits on shuffled proteins (with a window size of 20 amino acids) and the number of hits on native (unshuffled) proteins may be used as an index. Every pattern for which the ratio is greater than 20% (one hit on shuffled proteins for 5 hits on native proteins) may be skipped during the search with `prosite_scan`. The program used to shuffle protein sequences (`db_shuffled`) and the program used to determine the statistics for each pattern in the protein data banks (`prosite_statistics`) are available on the ftp site [http://ulrec3.unil.ch/ftpserveur/prosite\\_scan](http://ulrec3.unil.ch/ftpserveur/prosite_scan).

In addition to PCR based methods for obtaining extended cDNAs, traditional hybridization based methods may also be employed. These methods may also be used to obtain the genomic DNAs which encode the mRNAs from which the 5' ESTs were derived, mRNAs corresponding to the extended cDNAs, or nucleic acids which are homologous to extended cDNAs or 5' ESTs. Example 29 below provides examples of such methods.

#### EXAMPLE 29

##### Methods for Obtaining cDNAs which include the Entire Coding Region and the Authentic 5' End of the Corresponding mRNA

A full length cDNA library can be made using the strategies described in Examples 13, 14, 15, and 16 above by replacing the random nonamer used in Example 14 with an oligo-dT primer. For instance, the oligonucleotide of SEQ ID NO: 14 may be used.

Alternatively, a cDNA library or genomic DNA library may be obtained from a commercial source or made using techniques familiar to those skilled in the art. Such cDNA or genomic DNA libraries may be used to isolate extended cDNAs obtained from 5' EST or nucleic acids homologous to extended cDNAs or 5' EST as follows. The cDNA library or genomic DNA library is hybridized to a detectable probe comprising at least 10 consecutive

nucleotides from the 5' EST or extended cDNA using conventional techniques. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST or extended cDNA. More preferably, the probe comprises at least 20 to 30 consecutive nucleotides from the 5' EST or extended cDNA. In some embodiments, the probe comprises  
5 more than 30 nucleotides from the 5' EST or extended cDNA.

Techniques for identifying cDNA clones in a cDNA library which hybridize to a given probe sequence are disclosed in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual 2d Ed.*, Cold Spring Harbor Laboratory Press, 1989, the disclosure of which is incorporated  
10 herein by reference. The same techniques may be used to isolate genomic DNAs.

Briefly, cDNA or genomic DNA clones which hybridize to the detectable probe are identified and isolated for further manipulation as follows. A probe comprising at least 10 consecutive nucleotides from the 5' EST or extended cDNA is labeled with a detectable label such as a radioisotope or a fluorescent molecule. Preferably, the probe comprises at least 12,  
15 15, or 17 consecutive nucleotides from the 5' EST or extended cDNA. More preferably, the probe comprises 20 to 30 consecutive nucleotides from the 5' EST or extended cDNA. In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST or extended cDNA.

Techniques for labeling the probe are well known and include phosphorylation with  
20 polynucleotide kinase, nick translation, *in vitro* transcription, and non radioactive techniques. The cDNAs or genomic DNAs in the library are transferred to a nitrocellulose or nylon filter and denatured. After blocking of non specific sites, the filter is incubated with the labeled probe for an amount of time sufficient to allow binding of the probe to cDNAs or genomic DNAs containing a sequence capable of hybridizing thereto.

25 By varying the stringency of the hybridization conditions used to identify extended cDNAs or genomic DNAs which hybridize to the detectable probe, extended cDNAs having different levels of homology to the probe can be identified and isolated as described below.

1. Identification of Extended cDNA or Genomic cDNA Sequences Having a High Degree of Homology to the Labeled Probe

To identify extended cDNAs or genomic DNAs having a high degree of homology to the probe sequence, the melting temperature of the probe may be calculated using the following formulas:

For probes between 14 and 70 nucleotides in length the melting temperature ( $T_m$ ) is calculated using the formula:  $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (600/N)$  where N is the length of the probe.

If the hybridization is carried out in a solution containing formamide, the melting temperature may be calculated using the equation  $T_m = 81.5 + 16.6(\log [Na^+]) + 0.41(\text{fraction G+C}) - (0.63\% \text{ formamide}) - (600/N)$  where N is the length of the probe.

Prehybridization may be carried out in 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100  $\mu$ g denatured fragmented salmon sperm DNA or 6X SSC, 5X Denhardt's reagent, 0.5% SDS, 100  $\mu$ g denatured fragmented salmon sperm DNA, 50% formamide. The formulas for SSC and Denhardt's solutions are listed in Sambrook *et al.*, *supra*.

Hybridization is conducted by adding the detectable probe to the prehybridization solutions listed above. Where the probe comprises double stranded DNA, it is denatured before addition to the hybridization solution. The filter is contacted with the hybridization solution for a sufficient period of time to allow the probe to hybridize to extended cDNAs or genomic DNAs containing sequences complementary thereto or homologous thereto. For probes over 200 nucleotides in length, the hybridization may be carried out at 15-25°C below the  $T_m$ . For shorter probes, such as oligonucleotide probes, the hybridization may be conducted at 15-25°C below the  $T_m$ . Preferably, for hybridizations in 6X SSC, the hybridization is conducted at approximately 68°C. Preferably, for hybridizations in 50% formamide containing solutions, the hybridization is conducted at approximately 42°C.

All of the foregoing hybridizations would be considered to be under "stringent" conditions.

Following hybridization, the filter is washed in 2X SSC, 0.1% SDS at room temperature for 15 minutes. The filter is then washed with 0.1X SSC, 0.5% SDS at room temperature for 30 minutes to 1 hour. Thereafter, the solution is washed at the hybridization

temperature in 0.1X SSC, 0.5% SDS. A final wash is conducted in 0.1X SSC at room temperature.

Extended cDNAs, nucleic acids homologous to extended cDNAs or 5' ESTs, or genomic DNAs which have hybridized to the probe are identified by autoradiography or other conventional techniques.

## 2. Obtention of Extended cDNA or Genomic cDNA Sequences Having Lower Degrees of Homology to the Labeled Probe

The above procedure may be modified to identify extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs having decreasing levels of homology to the probe sequence. For example, to obtain extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs of decreasing homology to the detectable probe, less stringent conditions may be used. For example, the hybridization temperature may be decreased in increments of 5°C from 68°C to 42°C in a hybridization buffer having a sodium concentration of approximately 1M. Following hybridization, the filter may be washed with 2X SSC, 0.5% SDS at the temperature of hybridization. These conditions are considered to be "moderate" conditions above 50°C and "low" conditions below 50°C.

Alternatively, the hybridization may be carried out in buffers, such as 6X SSC, containing formamide at a temperature of 42°C. In this case, the concentration of formamide in the hybridization buffer may be reduced in 5% increments from 50% to 0% to identify clones having decreasing levels of homology to the probe. Following hybridization, the filter may be washed with 6X SSC, 0.5% SDS at 50°C. These conditions are considered to be "moderate" conditions above 25% formamide and "low" conditions below 25% formamide.

Extended cDNAs, nucleic acids homologous to extended cDNAs, or genomic DNAs which have hybridized to the probe are identified by autoradiography.

## 3. Determination of the Degree of Homology Between the Obtained Extended cDNAs and the Labeled Probe

If it is desired to obtain nucleic acids homologous to extended cDNAs, such as allelic variants thereof or nucleic acids encoding proteins related to the proteins encoded by the extended cDNAs, the level of homology between the hybridized nucleic acid and the

extended cDNA or 5' EST used as the probe may be further determined using BLAST2N; parameters may be adapted depending on the sequence length and degree of homology studied. To determine the level of homology between the hybridized nucleic acid and the extended cDNA or 5'EST from which the probe was derived, the nucleotide sequences of the hybridized nucleic acid and the extended cDNA or 5'EST from which the probe was derived are compared. For example, using the above methods, nucleic acids having at least 95% nucleic acid homology to the extended cDNA or 5'EST from which the probe was derived may be obtained and identified. Similarly, by using progressively less stringent hybridization conditions one can obtain and identify nucleic acids having at least 90%, at least 85%, at least 80% or at least 75% homology to the extended cDNA or 5'EST from which the probe was derived.

To determine whether a clone encodes a protein having a given amount of homology to the protein encoded by the extended cDNA or 5' EST, the amino acid sequence encoded by the extended cDNA or 5' EST is compared to the amino acid sequence encoded by the hybridizing nucleic acid. Homology is determined to exist when an amino acid sequence in the extended cDNA or 5' EST is closely related to an amino acid sequence in the hybridizing nucleic acid. A sequence is closely related when it is identical to that of the extended cDNA or 5' EST or when it contains one or more amino acid substitutions therein in which amino acids having similar characteristics have been substituted for one another. Using the above methods and algorithms such as FASTA with parameters depending on the sequence length and degree of homology studied, one can obtain nucleic acids encoding proteins having at least 95%, at least 90%, at least 85%, at least 80% or at least 75% homology to the proteins encoded by the extended cDNA or 5'EST from which the probe was derived.

In addition to the above described methods, other protocols are available to obtain extended cDNAs using 5' ESTs as outlined in the following paragraphs.

Extended cDNAs may be prepared by obtaining mRNA from the tissue, cell, or organism of interest using mRNA preparation procedures utilizing polyA selection procedures or other techniques known to those skilled in the art. A first primer capable of hybridizing to the polyA tail of the mRNA is hybridized to the mRNA and a reverse transcription reaction is performed to generate a first cDNA strand.



The first cDNA strand is hybridized to a second primer containing at least 10 consecutive nucleotides of the sequences of SEQ ID NOs 38-270. Preferably, the primer comprises at least 12, 15, or 17 consecutive nucleotides from the sequences of SEQ ID NOs 38-270. More preferably, the primer comprises 20 to 30 consecutive nucleotides from the sequences of SEQ ID NOs 38-270. In some embodiments, the primer comprises more than 30 nucleotides from the sequences of SEQ ID NOs 38-270. If it is desired to obtain extended cDNAs containing the full protein coding sequence, including the authentic translation initiation site, the second primer used contains sequences located upstream of the translation initiation site. The second primer is extended to generate a second cDNA strand complementary to the first cDNA strand. Alternatively, RT-PCR may be performed as described above using primers from both ends of the cDNA to be obtained.

Extended cDNAs containing 5' fragments of the mRNA may be prepared by hybridizing an mRNA comprising the sequence of the 5'EST for which an extended cDNA is desired with a primer comprising at least 10 consecutive nucleotides of the sequences complementary to the 5'EST and reverse transcribing the hybridized primer to make a first cDNA strand from the mRNAs. Preferably, the primer comprises at least 12, 15, or 17 consecutive nucleotides from the 5'EST. More preferably, the primer comprises 20 to 30 consecutive nucleotides from the 5'EST.

Thereafter, a second cDNA strand complementary to the first cDNA strand is synthesized. The second cDNA strand may be made by hybridizing a primer complementary to sequences in the first cDNA strand to the first cDNA strand and extending the primer to generate the second cDNA strand.

The double stranded extended cDNAs made using the methods described above are isolated and cloned. The extended cDNAs may be cloned into vectors such as plasmids or viral vectors capable of replicating in an appropriate host cell. For example, the host cell may be a bacterial, mammalian, avian, or insect cell.

Techniques for isolating mRNA, reverse transcribing a primer hybridized to mRNA to generate a first cDNA strand, extending a primer to make a second cDNA strand complementary to the first cDNA strand, isolating the double stranded cDNA and cloning the double stranded cDNA are well known to those skilled in the art and are described in *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc. 1997 and Sambrook *et al.*

*Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989, the entire disclosures of which are incorporated herein by reference.

Alternatively, procedures such as the one described in Example 29 may be used for obtaining full length cDNAs or extended cDNAs. In this approach, full length or extended  
5 cDNAs are prepared from mRNA and cloned into double stranded phagemids as follows. The cDNA library in the double stranded phagemids is then rendered single stranded by treatment with an endonuclease, such as the Gene II product of the phage F1, and an exonuclease (Chang *et al.*, *Gene* 127:95-8, 1993). A biotinylated oligonucleotide comprising the sequence of a 5' EST, or a fragment containing at least 10 nucleotides thereof, is  
10 hybridized to the single stranded phagemids. Preferably, the fragment comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST. More preferably, the fragment comprises 20-30 consecutive nucleotides from the 5' EST. In some procedures, the fragment may comprise more than 30 consecutive nucleotides from the 5' EST.

Hybrids between the biotinylated oligonucleotide and phagemids having inserts  
15 containing the 5' EST sequence are isolated by incubating the hybrids with streptavidin coated paramagnetic beads and retrieving the beads with a magnet (Fry *et al.*, *Biotechniques*, 13: 124-131, 1992). Thereafter, the resulting phagemids containing the 5' EST sequence are released from the beads and converted into double stranded DNA using a primer specific for the 5' EST sequence. Alternatively, protocols such as the Gene Trapper kit (Gibco BRL)  
20 may be used. The resulting double stranded DNA is transformed into bacteria. Extended cDNAs containing the 5' EST sequence are identified by colony PCR or colony hybridization

Using any of the above described methods in section III, a plurality of extended cDNAs containing full length protein coding sequences or sequences encoding only the  
25 mature protein remaining after the signal peptide is cleaved off may be provided as cDNA libraries for subsequent evaluation of the encoded proteins or use in diagnostic assays as described below.

#### IV. Expression of Proteins Encoded by Extended cDNAs Isolated Using 5' ESTs

30 Extended cDNAs containing the full protein coding sequences of their corresponding mRNAs or portions thereof, such as cDNAs encoding the mature protein, may be used to

express the encoded secreted proteins or portions thereof as described in Example 30 below. If desired, the extended cDNAs may contain the sequences encoding the signal peptide to facilitate secretion of the expressed protein. It will be appreciated that a plurality of extended cDNAs containing the full protein coding sequences or portions thereof may be simultaneously cloned into expression vectors to create an expression library for analysis of the encoded proteins as described below.

### EXAMPLE 30

#### Expression of the Proteins Encoded by the Genes Corresponding to 5' ESTs or Portions Thereof

To express the proteins encoded by the genes corresponding to 5' ESTs (or portions thereof), full length cDNAs containing the entire protein coding region or extended cDNAs containing sequences adjacent to the 5' ESTs (or portions thereof) are obtained as described in Examples 27-29 and cloned into a suitable expression vector. If desired, the nucleic acids may contain the sequences encoding the signal peptide to facilitate secretion of the expressed protein. The nucleic acids inserted into the expression vectors may also contain sequences upstream of the sequences encoding the signal peptide, such as sequences which regulate expression levels or sequences which confer tissue specific expression.

The nucleic acid encoding the protein or polypeptide to be expressed is operably linked to a promoter in an expression vector using conventional cloning technology. The expression vector may be any of the mammalian, yeast, insect or bacterial expression systems known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). If desired, to enhance expression and facilitate proper protein folding, the codon context and codon pairing of the sequence may be optimized for the particular expression organism in which the expression vector is introduced, as explained by Hatfield, *et al.*, U.S. Patent No. 5,082,767, incorporated herein by this reference.

The cDNA cloned into the expression vector may encode the entire protein (*i.e.* the signal peptide and the mature protein), the mature protein (*i.e.* the protein created by cleaving the signal peptide off), only the signal peptide or any other portion thereof.

The following is provided as one exemplary method to express the proteins encoded by the extended cDNAs corresponding to the 5' ESTs or the nucleic acids described above. First, the methionine initiation codon for the gene and the polyA signal of the gene are identified. If the nucleic acid encoding the polypeptide to be expressed lacks a methionine to serve as the initiation site, an initiating methionine can be introduced next to the first codon of the nucleic acid using conventional techniques. Similarly, if the extended cDNA lacks a polyA signal, this sequence can be added to the construct by, for example, splicing out the polyA signal from pSG5 (Stratagene) using BglII and SalI restriction endonuclease enzymes and incorporating it into the mammalian expression vector pXT1 (Stratagene). pXT1 contains the LTRs and a portion of the *gag* gene from Moloney Murine Leukemia Virus. The position of the LTRs in the construct allow efficient stable transfection. The vector includes the Herpes Simplex thymidine kinase promoter and the selectable neomycin gene. The extended cDNA or portion thereof encoding the polypeptide to be expressed is obtained by PCR from the bacterial vector using oligonucleotide primers complementary to the extended cDNA or portion thereof and containing restriction endonuclease sequences for Pst I incorporated into the 5' primer and BglII at the 5' end of the corresponding cDNA 3' primer, taking care to ensure that the extended cDNA is positioned with the poly A signal. The purified fragment obtained from the resulting PCR reaction is digested with PstI, blunt ended with an exonuclease, digested with Bgl II, purified and ligated to pXT1 containing a poly A signal and prepared for this ligation (blunt/BglII).

The ligated product is transfected into mouse NIH 3T3 cells using Lipofectin (Life Technologies, Inc., Grand Island, New York) under conditions outlined in the product specification. Positive transfectants are selected after growing the transfected cells in 600 µg/ml G418 (Sigma, St. Louis, Missouri). Preferably the expressed protein is released into the culture medium, thereby facilitating purification.

Alternatively, the extended cDNAs may be cloned into pED6dpc2 as described above. The resulting pED6dpc2 constructs may be transfected into a suitable host cell, such as COS 1 cells. Methotrexate resistant cells are selected and expanded. Preferably, the protein expressed from the extended cDNA is released into the culture medium thereby facilitating purification.

Proteins in the culture medium are separated by gel electrophoresis. If desired, the proteins may be ammonium sulfate precipitated or separated based on size or charge prior to electrophoresis.

As a control, the expression vector lacking a cDNA insert is introduced into host cells  
5 or organisms and the proteins in the medium are harvested. The secreted proteins present in the medium are detected using techniques familiar to those skilled in the art such as Coomassie blue or silver staining or using antibodies against the protein encoded by the extended cDNA

Antibodies capable of specifically recognizing the protein of interest may be generated  
10 using synthetic 15-mer peptides having a sequence encoded by the appropriate 5' EST, extended cDNA, or portion thereof. The synthetic peptides are injected into mice to generate antibody to the polypeptide encoded by the 5' EST, extended cDNA, or portion thereof.

Secreted proteins from the host cells or organisms containing an expression vector which contains the extended cDNA derived from a 5' EST or a portion thereof are compared  
15 to those from the control cells or organism. The presence of a band in the medium from the cells containing the expression vector which is absent in the medium from the control cells indicates that the extended cDNA encodes a secreted protein. Generally, the band corresponding to the protein encoded by the extended cDNA will have a mobility near that expected based on the number of amino acids in the open reading frame of the extended  
20 cDNA. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic cleavage.

Alternatively, if the protein expressed from the above expression vectors does not contain sequences directing its secretion, the proteins expressed from host cells containing an expression vector with an insert encoding a secreted protein or portion thereof can be  
25 compared to the proteins expressed in control host cells containing the expression vector without an insert. The presence of a band in samples from cells containing the expression vector with an insert which is absent in samples from cells containing the expression vector without an insert indicates that the desired protein or portion thereof is being expressed. Generally, the band will have the mobility expected for the secreted protein or portion  
30 thereof. However, the band may have a mobility different than that expected as a result of modifications such as glycosylation, ubiquitination, or enzymatic cleavage.

The protein encoded by the extended cDNA may be purified using standard immunochromatography techniques. In such procedures, a solution containing the secreted protein, such as the culture medium or a cell extract, is applied to a column having antibodies against the secreted protein attached to the chromatography matrix. The secreted protein is  
5 allowed to bind the immunochromatography column. Thereafter, the column is washed to remove non-specifically bound proteins. The specifically bound secreted protein is then released from the column and recovered using standard techniques.

If antibody production is not possible, the extended cDNA sequence or portion thereof may be incorporated into expression vectors designed for use in purification schemes  
10 employing chimeric polypeptides. In such strategies, the coding sequence of the extended cDNA or portion thereof is inserted in frame with the gene encoding the other half of the chimera. The other half of the chimera may be  $\beta$ -globin or a nickel binding polypeptide. A chromatography matrix having antibody to  $\beta$ -globin or nickel attached thereto is then used to purify the chimeric protein. Protease cleavage sites may be engineered between the  $\beta$ -globin  
15 gene or the nickel binding polypeptide and the extended cDNA or portion thereof. Thus, the two polypeptides of the chimera may be separated from one another by protease digestion.

One useful expression vector for generating  $\beta$ -globin chimerics is pSG5 (Stratagene), which encodes rabbit  $\beta$ -globin. Intron II of the rabbit  $\beta$ -globin gene facilitates splicing of the expressed transcript, and the polyadenylation signal incorporated into the construct increases  
20 the level of expression. These techniques as described are well known to those skilled in the art of molecular biology. Standard methods are published in methods texts such as Davis *et al.*, (*Basic Methods in Molecular Biology*, Davis, Dibner, and Battey, ed., Elsevier Press, NY, 1986) and many of the methods are available from Stratagene, Life Technologies, Inc., or Promega. Polypeptide may additionally be produced from the construct using *in vitro*  
25 translation systems such as the *In vitro* Express<sup>TM</sup> Translation Kit (Stratagene).

Following expression and purification of the secreted proteins encoded by the 5' ESTs, extended cDNAs, or fragments thereof, the purified proteins may be tested for the ability to bind to the surface of various cell types as described in Example 31 below. It will be  
30 appreciated that a plurality of proteins expressed from these cDNAs may be included in a

panel of proteins to be simultaneously evaluated for the activities specifically described below, as well as other biological roles for which assays for determining activity are available.

### EXAMPLE 31

#### 5      Analysis of Secreted Proteins to Determine Whether they Bind to the Cell Surface

The proteins encoded by the 5' ESTs, extended cDNAs, or fragments thereof are cloned into expression vectors such as those described in Example 30. The proteins are purified by size, charge, immunochromatography or other techniques familiar to those skilled in the art. Following purification, the proteins are labeled using techniques known to those skilled in the art. The labeled proteins are incubated with cells or cell lines derived from a variety of organs or tissues to allow the proteins to bind to any receptor present on the cell surface. Following the incubation, the cells are washed to remove non-specifically bound protein. The labeled proteins are detected by autoradiography. Alternatively, unlabeled proteins may be incubated with the cells and detected with antibodies having a detectable label, such as a fluorescent molecule, attached thereto.

Specificity of cell surface binding may be analyzed by conducting a competition analysis in which various amounts of unlabeled protein are incubated along with the labeled protein. The amount of labeled protein bound to the cell surface decreases as the amount of competitive unlabeled protein increases. As a control, various amounts of an unlabeled protein unrelated to the labeled protein is included in some binding reactions. The amount of labeled protein bound to the cell surface does not decrease in binding reactions containing increasing amounts of unrelated unlabeled protein, indicating that the protein encoded by the cDNA binds specifically to the cell surface.

As discussed above, secreted proteins have been shown to have a number of important physiological effects and, consequently, represent a valuable therapeutic resource. The secreted proteins encoded by the extended cDNAs or portions thereof made according to Examples 27-29 may be evaluated to determine their physiological activities as described below.

## EXAMPLE 32

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Cytokine,  
Cell Proliferation or Cell Differentiation Activity

As discussed above, secreted proteins may act as cytokines or may affect cellular proliferation or differentiation. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein encoded by the extended cDNAs is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M<sup>c</sup> (preB M<sup>c</sup>), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7c and CMK. The proteins encoded by the above extended cDNAs or portions thereof may be evaluated for their ability to regulate T cell or thymocyte proliferation in assays such as those described above or in the following references, which are incorporated herein by reference: *Current Protocols in Immunology*, Ed. by Coligan *et al.*, Greene Publishing Associates and Wiley-Interscience; Takai *et al.* *J. Immunol.* 137:3494-3500, 1986.; Bertagnolli *et al.*, *J. Immunol.* 145:1706-1712, 1990.; Bertagnolli *et al.*, *Cell. Immunol.* 133:327-341, 1991; Bertagnolli, *et al.*, *J. Immunol.* 149:3778-3783, 1992, Bowman *et al.*, *J. Immunol.* 152:1756-1761, 1994.

In addition, numerous assays for cytokine production and/or the proliferation of spleen cells, lymph node cells and thymocytes are known. These include the techniques disclosed in *Current Protocols in Immunology*, *supra* 1:3.12.1-3.12.14; and Schreiber In *Current Protocols in Immunology*, *supra* 1: 6.8.1-6.8.8.

The proteins encoded by the cDNAs may also be assayed for the ability to regulate the proliferation and differentiation of hematopoietic or lymphopoietic cells. Many assays for such activity are familiar to those skilled in the art, including the assays in the following references, which are incorporated herein by reference: Bottomly *et al.*, In *Current Protocols in Immunology*, *supra* 1: 6.3.1-6.3.12.; deVries *et al.*, *J. Exp. Med.* 173:1205-1211, 1991, Moreau *et al.*, *Nature* 36:690-692, 1988; Greenberger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983; Nordan, R., In *Current Protocols in Immunology*, *supra* 1: 6.6.1-6.6.5; Smith *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; Bennett *et al.*, in



*Current Protocols in Immunology supra* 1 : 6.15.1; Ciarletta *et al.*, In *Current Protocols in Immunology. supra* 1 : 6.13.1.

The proteins encoded by the cDNAs may also be assayed for their ability to regulate T-cell responses to antigens. Many assays for such activity are familiar to those skilled in the art, including the assays described in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function), Chapter 6 (Cytokines and Their Cellular Receptors) and Chapter 7, (Immunologic Studies in Humans) in *Current Protocols in Immunology supra*; Weinberger *et al.*, *Proc. Natl. Acad. Sci. USA* 77:6091-6095, 1980; Weinberger *et al.*, *Eur. J. Immun.* 11:405-411, 1981; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988.

Those proteins which exhibit cytokine, cell proliferation, or cell differentiation activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which induction of cell proliferation or differentiation is beneficial. Alternatively, as described in more detail below, genes encoding these proteins or nucleic acids regulating the expression of these proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

### EXAMPLE 33

#### Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Activity as Immune System Regulators

The proteins encoded by the cDNAs may also be evaluated for their effects as immune regulators. For example, the proteins may be evaluated for their activity to influence thymocyte or splenocyte cytotoxicity. Numerous assays for such activity are familiar to those skilled in the art including the assays described in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function 3.1-3.19) and Chapter 7 (Immunologic studies in Humans) in *Current Protocols in Immunology*, Coligan *et al.*, Eds, Greene Publishing Associates and Wiley-Interscience; Herrmann *et al.*, *Proc. Natl. Acad. Sci. USA* 78:2488-2492, 1981; Herrmann *et al.*, *J. Immunol.* 128:1968-1974, 1982; Handa *et al.*, *J. Immunol.* 135:1564-1572, 1985; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988;

Bowman *et al.*, *J. Virology* 61:1992-1998; Bertagnolli *et al.*, *Cell. Immunol.* 133:327-341, 1991; Brown *et al.*, *J. Immunol.* 153:3079-3092, 1994.

The proteins encoded by the cDNAs may also be evaluated for their effects on T-cell dependent immunoglobulin responses and isotype switching. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Maliszewski, *J. Immunol.* 144:3028-3033, 1990; Mond *et al.* in *Current Protocols in Immunology*, 1: 3.8.1-3.8.16, *supra*.

The proteins encoded by the cDNAs may also be evaluated for their effect on immune effector cells, including their effect on Th1 cells and cytotoxic lymphocytes. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Chapter 3 (*In Vitro* Assays for Mouse Lymphocyte Function 3.1-3.19) and Chapter 7 (Immunologic Studies in Humans) in *Current Protocols in Immunology*, *supra*; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; Takai *et al.*, *J. Immunol.* 140:508-512, 1988; Bertagnolli *et al.*, *J. Immunol.* 149:3778-3783, 1992.

The proteins encoded by the cDNAs may also be evaluated for their effect on dendritic cell mediated activation of naive T-cells. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Guery *et al.*, *J. Immunol.* 134:536-544, 1995; Inaba *et al.*, *J. Exp. Med.* 173:549-559, 1991; Macatonia *et al.*, *J. Immunol.* 154:5071-5079, 1995; Porgador *et al.*, *J. Exp. Med.* 182:255-260, 1995; Nair *et al.*, *J. Virol.* 67:4062-4069, 1993; Huang *et al.*, *Science* 264:961-965, 1994; Macatonia *et al.*, *J. Exp. Med.* 169:1255-1264, 1989; Bhardwaj *et al.*, *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba *et al.*, *J. Exp. Med.* 172:631-640, 1990.

The proteins encoded by the cDNAs may also be evaluated for their influence on the lifetime of lymphocytes. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Darzynkiewicz *et al.*, *Cytometry* 13:795-808, 1992; Gorczyca *et al.*, *Leukemia* 7:659-670, 1993; Gorczyca *et al.*, *Cancer Res.* 53:1945-1951, 1993; Itoh *et al.*, *Cell* 66:233-243, 1991; Zacharchuk, *J. Immunol.* 145:4037-4045, 1990; Zamai *et al.*, *Cytometry* 14:891-897, 1993; Gorczyca *et al.*, *Int. J. Oncol.* 1:639-648, 1992.

The proteins encoded by the cDNAs may also be evaluated for their influence on early steps of T-cell commitment and development. Numerous assays for such activity are familiar to those skilled in the art, including without limitation the assays disclosed in the following references, which are incorporated herein by references: Antica *et al.*, *Blood* 84:111-117, 1994; Fine *et al.*, *Cell. Immunol.* 155:111-122, 1994; Galy *et al.*, *Blood* 85:2770-2778, 1995; Toki *et al.*, *Proc. Nat. Acad. Sci. USA* 88:7548-7551, 1991.

Those proteins which exhibit activity as immune system regulators activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of immune activity is beneficial. For example, the protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, *Leishmania* spp., *Plasmodium* and various fungal infections such as candidiasis. Of course, in this regard, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the treatment of cancer.

Alternatively, proteins encoded by extended cDNAs derived from the 5' ESTs of the present invention may be used in treatment of autoimmune disorders including, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention.

Using the proteins of the invention it may also be possible to regulate immune responses either up or down.

Down regulation may involve inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of  
5 activated T-cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active non-antigen-specific process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T  
10 cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after the end of exposure to the tolerizing agent. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions, such as, for example, B7 costimulation), e.g.,  
15 preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the  
20 transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation, can lead to the binding of the molecule to the natural  
25 ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this manner prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may  
30 avoid the necessity of repeated administration of these blocking reagents. To achieve

sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans.

- 5 Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, *Science* 257:789-792, 1992 and Turka *et al.*, *Proc. Natl. Acad. Sci USA*, 89:11102-11105, 1992. In addition, murine models of GVHD (see Paul ed., *Fundamental*  
10 *Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

- Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and  
15 autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor/ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which potentially involved in the disease process.  
20 Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/pr/pr mice or  
25 NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in OD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., *supra*, pp. 840-856).

- Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may involve either enhancing an existing immune response or eliciting an  
30 initial immune response as shown by the following examples. For instance, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases

of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory form of B lymphocyte antigens systemically.

Alternatively, antiviral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen-pulsed APCs either expressing a peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention or together with a stimulatory form of a soluble peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention and reintroducing the *in vitro* primed T cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to T cells *in vivo*, thereby activating the T cells.

In another application, upregulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide encoded by extended cDNAs derived from the 5' ESTs of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules can be transfected with nucleic acids encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I  $\alpha$  chain and  $\beta_2$  microglobulin or an MHC class II  $\alpha$  chain and an MHC class II  $\beta$  chain to thereby express MHC class I or MHC class II proteins on the cell surface, respectively. Expression of the appropriate MHC class I or class II

molecules in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA  
5 encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject. Alternatively, as described in more detail below, genes encoding these immune system regulator proteins or nucleic acids regulating the expression of  
10 such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

#### EXAMPLE 34

##### Assaying the Proteins Expressed from Extended cDNAs

##### or Portions Thereof for Hematopoiesis Regulating Activity

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their hematopoiesis regulating activity. For example, the effect of the proteins on embryonic stem cell differentiation may be evaluated. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following  
20 references, which are incorporated herein by reference: Johansson *et al.* *Cell. Biol.* 15:141-151, 1995; Keller *et al.*, *Mol. Cell. Biol.* 13:473-486, 1993; McClanahan *et al.*, *Blood* 81:2903-2915, 1993.

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their influence on the lifetime of stem cells and stem cell differentiation.  
25 Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Freshney, *Methylcellulose Colony Forming Assays*, in *Culture of Hematopoietic Cells*, Freshney, *et al.* Eds. pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama *et al.*, *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; McNiece and Briddell, in *Culture of Hematopoietic Cells*,  
30 *supra*; Neben *et al.*, *Exp. Hematol.* 22:353-359, 1994; Ploemacher and Cobblestone In

*Culture of Hematopoietic Cells, supra* 1-21, Spooncer *et al*, in *Culture of Hematopoietic Cells, supra* 163-179 and Sutherland in *Culture of Hematopoietic Cells, supra*. 139-162.

Those proteins which exhibit hematopoiesis regulatory activity may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of hematopoiesis is beneficial, such as in the treatment of myeloid or lymphoid cell deficiencies. Involvement in regulating hematopoiesis is indicated even by marginal biological activity in support of colony forming cells or of factor-dependent cell lines. For example, proteins supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, indicates utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells. Proteins supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (*i.e.*, traditional CSF activity) may be useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression. Proteins supporting the growth and proliferation of megakaryocytes and consequently of platelets allows prevention or treatment of various platelet disorders such as thrombocytopenia, and generally may be used in place of or complementary to platelet transfusions. Proteins supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells may therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in vivo* or *ex vivo* (*i.e.*, in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy. Alternatively, as described in more detail below, genes encoding hematopoiesis regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

#### EXAMPLE 35

Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof  
for Regulation of Tissue Growth



The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their effect on tissue growth. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in International Patent Publication No. WO95/16035, International Patent Publication No. WO95/05846 and International Patent  
5 Publication No. WO91/07491, which are incorporated herein by reference.

Assays for wound healing activity include, without limitation, those described in: Winter, *Epidermal Wound Healing*, pps. 71-112, Maibach and Rovee, eds., Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, *J. Invest. Dermatol.* 71:382-84, 1978, which are incorporated herein by reference.

10 Those proteins which are involved in the regulation of tissue growth may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of tissue growth is beneficial. For example, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound  
15 healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention  
20 may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone synthesis induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease,  
25 and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of bone-forming cell progenitors. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast  
30 activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein encoded by extended cDNAs derived from the 5' ESTs of the present invention is tendon/ligament formation. A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition encoded by extended cDNAs derived from the 5' ESTs of the present invention contributes to the repair of tendon or ligaments defects of congenital, traumatic or other origin and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions encoded by extended cDNAs derived from the 5' ESTs of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.*, for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders

head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium) muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to generate. A protein of the invention may also exhibit angiogenic activity.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

Alternatively, as described in more detail below, genes encoding tissue growth regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

### EXAMPLE 36

#### Assaying the Proteins Expressed from Extended cDNAs or Portions

##### Thereof for Regulation of Reproductive Hormones

The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their ability to regulate reproductive hormones, such as follicle stimulating hormone. Numerous assays for such activity are familiar to those skilled in the art, including

the assays disclosed in the following references, which are incorporated herein by reference: Vale *et al.*, *Endocrinol.* 91:562-572, 1972; Ling *et al.*, *Nature* 321:779-782, 1986; Vale *et al.*, *Nature* 321:776-779, 1986; Mason *et al.*, *Nature* 318:659-663, 1985; Forage *et al.*, *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986, Chapter 6.12 in *Current Protocols in Immunology*, Coligan *et al.* Eds. Greene Publishing Associates and Wiley-Interscience; Taub *et al.*, *J. Clin. Invest.* 95:1370-1376, 1995; Lind *et al.*, *APMIS* 103:140-146, 1995; Muller *et al.*, *Eur. J. Immunol.* 25:1744-1748; Gruber *et al.*, *J. Immunol.* 152:5860-5867, 1994; Johnston *et al.*, *J. Immunol.* 153:1762-1768, 1994.

Those proteins which exhibit activity as reproductive hormones or regulators of cell movement may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of reproductive hormones are beneficial. For example, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of FSH. Thus, a protein encoded by extended cDNAs derived from the 5' ESTs of the present invention, alone or in heterodimers with a member of the inhibin  $\alpha$  family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin-B group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885, the disclosure of which is incorporated herein by reference. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

Alternatively, as described in more detail below, genes encoding reproductive hormone regulating activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

**EXAMPLE 37****Assaying the Proteins Expressed from Extended cDNAs or  
Portions Thereof for Chemotactic/Chemokinetic Activity**

The proteins encoded by the extended cDNAs or portions thereof may also be  
5 evaluated for chemotactic/chemokinetic activity. For example, a protein encoded by  
extended cDNAs derived from the 5' ESTs of the present invention may have chemotactic or  
chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example,  
monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or  
endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a  
10 desired cell population to a desired site of action. Chemotactic or chemokinetic proteins  
provide particular advantages in treatment of wounds and other trauma to tissues, as well as  
in treatment of localized infections. For example, attraction of lymphocytes, monocytes or  
neutrophils to tumors or sites of infection may result in improved immune responses against  
the tumor or infecting agent.

15 A protein or peptide has chemotactic activity for a particular cell population if it can  
stimulate, directly or indirectly, the directed orientation or movement of such cell population.  
Preferably, the protein or peptide has the ability to directly stimulate directed movement of  
cells. Whether a particular protein has chemotactic activity for a population of cells can be  
readily determined by employing such protein or peptide in any known assay for cell  
20 chemotaxis.

The activity of a protein of the invention may, among other means, be measured by  
the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent  
chemotaxis) consist of assays that measure the ability of a protein to induce the migration of  
25 cells across a membrane as well as the ability of a protein to induce the adhesion of one cell  
population to another cell population. Suitable assays for movement and adhesion include,  
without limitation, those described in: *Current Protocols in Immunology*, Ed by Coligan,  
Kruisbeek, Margulies, Shevach and Strober, Pub. Greene Publishing Associates and Wiley-  
Interscience, Chapter 6.12: 6.12.1-6.12.28; Taub *et al.*, *J. Clin. Invest.* 95:1370-1376, 1995;  
30 Lind *et al.*, *APMIS* 103:140-146, 1995; Mueller *et al.*, *Eur. J. Immunol.* 25:1744-1748;

Gruber *et al.*, *J. Immunol.* 152:5860-5867, 1994; Johnston *et al.* *J. Immunol.*, 153:1762-1768, 1994.

### EXAMPLE 38

#### Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Regulation of Blood Clotting

5 The proteins encoded by the extended cDNAs or portions thereof may also be evaluated for their effects on blood clotting. Numerous assays for such activity are familiar to those skilled in the art, including the assays disclosed in the following references, which are  
10 incorporated herein by reference: Linet *et al.*, *J. Clin. Pharmacol.* 26:131-140, 1986; Burdick *et al.*, *Thrombosis Res.* 45:413-419, 1987; Humphrey *et al.*, *Fibrinolysis* 5:71-79, 1991; Schaub, *Prostaglandins* 35:467-474, 1988.

Those proteins which are involved in the regulation of blood clotting may then be formulated as pharmaceuticals and used to treat clinical conditions in which regulation of  
15 blood clotting is beneficial. For example, a protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulations disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful  
20 for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as infarction of cardiac and central nervous system vessels (e.g., stroke)). Alternatively, as described in more detail below, genes encoding blood clotting activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as  
25 desired.

### EXAMPLE 39

#### Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Involvement in Receptor/Ligand Interactions

30 The proteins encoded by the extended cDNAs or a portion thereof may also be evaluated for their involvement in receptor/ligand interactions. Numerous assays for such

involvement are familiar to those skilled in the art, including the assays disclosed in the following references, which are incorporated herein by reference: Chapter 7. 7.28.1-7.28.22 in *Current Protocols in Immunology*, Coligan *et al.* Eds. Greene Publishing Associates and Wiley-Interscience; Takai *et al.*, *Proc. Natl. Acad. Sci. USA* 84:6864-6868, 1987; Bierer *et al.*, *J. Exp. Med.* 168:1145-1156, 1988; Rosenstein *et al.*, *J. Exp. Med.* 169:149-160, 1989; 5 Stoltenborg *et al.*, *J. Immunol. Methods* 175:59-68, 1994; Stitt *et al.*, *Cell* 80:661-670, 1995; Gyuris *et al.*, *Cell* 75:791-803, 1993.

For example, the proteins encoded by extended cDNAs derived from the 5' ESTs of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors 10 or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen 15 recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein encoded by extended cDNAs derived from the 5' ESTs of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions. Alternatively, 20 as described in more detail below, genes encoding proteins involved in receptor/ligand interactions or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

#### EXAMPLE 40

##### 25 Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Anti-Inflammatory Activity

The proteins encoded by the extended cDNAs or a portion thereof may also be evaluated for anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or 30 promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting

cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions, including without limitation inflammation associated with infection (such as septic shock, sepsis or  
5 systemic inflammatory response syndrome), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine- or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material. Alternatively, as  
10 described in more detail below, genes encoding anti-inflammatory activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

#### EXAMPLE 41

##### Assaying the Proteins Expressed from Extended cDNAs or Portions Thereof for Tumor Inhibition Activity

The proteins encoded by the extended cDNAs or a portion thereof may also be evaluated for tumor inhibition activity. In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other  
20 anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or  
25 inhibiting factors, agents or cell types which promote tumor growth. Alternatively, as described in more detail below, genes tumor inhibition activity proteins or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

30 A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents.



including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein. Alternatively, as described in more detail below, genes encoding proteins involved in any of the above mentioned activities or nucleic acids regulating the expression of such proteins may be introduced into appropriate host cells to increase or decrease the expression of the proteins as desired.

#### EXAMPLE 42

##### Identification of Proteins which Interact with Polypeptides Encoded by Extended cDNAs

Proteins which interact with the polypeptides encoded by cDNAs derived from the 5' ESTs or fragments thereof, such as receptor proteins, may be identified using two hybrid systems such as the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech). As described in the manual accompanying the kit which is incorporated herein by reference, the the cDNAs derived from 5' ESTs, or fragments thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA binding domain of the yeast

transcriptional activator GAL4. cDNAs in a cDNA library which encode proteins which might interact with the polypeptides encoded by the extended cDNAs or portions thereof are inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into yeast and the yeast are plated on selection medium which selects for expression of selectable markers on each of the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay contain plasmids encoding proteins which interact with the polypeptide encoded by the extended cDNAs or portions thereof.

Alternatively, the system described in Lustig *et al.*, *Methods in Enzymology* 283: 83-99, 1997, and in U.S. Patent No. 5,654,150, the disclosure of which is incorporated herein by reference, may be used for identifying molecules which interact with the polypeptides encoded by extended cDNAs. In such systems, *in vitro* transcription reactions are performed on a pool of vectors containing extended cDNA inserts cloned downstream of a promoter which drives *in vitro* transcription. The resulting pools of mRNAs are introduced into *Xenopus laevis* oocytes. The oocytes are then assayed for a desired activity.

Alternatively, the pooled *in vitro* transcription products produced as described above may be translated *in vitro*. The pooled *in vitro* translation products can be assayed for a desired activity or for interaction with a known polypeptide.

Proteins or other molecules interacting with polypeptides encoded by extended cDNAs can be found by a variety of additional techniques. In one method, affinity columns containing the polypeptide encoded by the extended cDNA or a portion thereof can be constructed. In some versions, of this method the affinity column contains chimeric proteins in which the protein encoded by the extended cDNA or a portion thereof is fused to glutathione S-transferase. A mixture of cellular proteins or pool of expressed proteins as described above and is applied to the affinity column. Proteins interacting with the polypeptide attached to the column can then be isolated and analyzed on 2-D electrophoresis gel as described in Ramunsen *et al.*, *Electrophoresis* 18:588-598, 1997, the disclosure of which is incorporated herein by reference. Alternatively, the proteins retained on the affinity column can be purified by electrophoresis based methods

and sequenced. The same method can be used to isolate antibodies, to screen phage display products, or to screen phage display human antibodies.

Proteins interacting with polypeptides encoded by extended cDNAs or portions thereof can also be screened by using an Optical Biosensor as described in Edwards and  
5 Leatherbarrow, *Analytical Biochemistry* 246:1-6, 1997, the disclosure of which is incorporated herein by reference. The main advantage of the method is that it allows the determination of the association rate between the protein and other interacting molecules. Thus, it is possible to specifically select interacting molecules with a high or low association rate. Typically a target molecule is linked to the sensor surface (through  
10 a carboxymethyl dextran matrix) and a sample of test molecules is placed in contact with the target molecules. The binding of a test molecule to the target molecule causes a change in the refractive index and/ or thickness. This change is detected by the Biosensor provided it occurs in the evanescent field (which extend a few hundred nanometers from the sensor surface). In these screening assays, the target molecule can  
15 be one of the polypeptides encoded by extended cDNAs or a portion thereof and the test sample can be a collection of proteins extracted from tissues or cells, a pool of expressed proteins, combinatorial peptide and/ or chemical libraries, or phage displayed peptides. The tissues or cells from which the test proteins are extracted can originate from any species.

20 In other methods, a target protein is immobilized and the test population is a collection of unique polypeptides encoded by the extended cDNAs or portions thereof.

To study the interaction of the proteins encoded by the extended cDNAs or portions thereof with drugs, the microdialysis coupled to HPLC method described by Wang *et al.*, *Chromatographia* 44:205-208, 1997 or the affinity capillary electrophoresis  
25 method described by Busch *et al.*, *J. Chromatogr.* 777:311-328, 1997, the disclosures of which are incorporated herein by reference can be used.

It will be appreciated by those skilled in the art that the proteins expressed from the extended cDNAs or portions may be assayed for numerous activities in addition to those  
30 specifically enumerated above. For example, the expressed proteins may be evaluated for applications involving control and regulation of inflammation, tumor proliferation or

metastasis, infection, or other clinical conditions. In addition, the proteins expressed from the extended cDNAs or portions thereof may be useful as nutritional agents or cosmetic agents.

The proteins expressed from the cDNAs or portions thereof may be used to generate antibodies capable of specifically binding to the expressed protein or fragments thereof as described in Example 40 below. The antibodies may be capable of binding a full length protein encoded by a cDNA derived from a 5' EST, a mature protein (*i.e.* the protein generated by cleavage of the signal peptide) encoded by a cDNA derived from a 5' EST, or a signal peptide encoded by a cDNA derived from a 5' EST. Alternatively, the antibodies may be capable of binding fragments of at least 10 amino acids of the proteins encoded by the above cDNAs. In some embodiments, the antibodies may be capable of binding fragments of at least 15 amino acids of the proteins encoded by the above cDNAs. In other embodiments, the antibodies may be capable of binding fragments of at least 25 amino acids of the proteins expressed from the extended cDNAs which comprise at least 25 amino acids of the proteins encoded by the above cDNAs. In further embodiments, the antibodies may be capable of binding fragments of at least 40 amino acids of the proteins encoded by the above cDNAs.

### EXAMPLE 43

#### Production of an Antibody to a Human Protein

Substantially pure protein or polypeptide is isolated from the transfected or transformed cells as described in Example 30. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter device, to the level of a few  $\mu\text{g/ml}$ . Monoclonal or polyclonal antibody to the protein can then be prepared as follows:

#### 1. Monoclonal Antibody Production by Hybridoma Fusion

Monoclonal antibody to epitopes of any of the peptides identified and isolated as described can be prepared from murine hybridomas according to the classical method of Kohler, and Milstein, *Nature* 256:495, 1975 or derivative methods thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the selected protein or peptides derived therefrom over a period of a few weeks. The mouse is then sacrificed, and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells, and the excess unfused cells

destroyed by growth of the system on selective media comprising aminopterin (HAT media). The successfully fused cells are diluted and aliquots of the dilution placed in wells of a microtiter plate where growth of the culture is continued. Antibody-producing clones are identified by detection of antibody in the supernatant fluid of the wells by immunoassay procedures, such as ELISA, as originally described by Engvall, *Meth. Enzymol.* 70:419, 1980, the disclosure of which is incorporated herein by reference and derivative methods thereof. Selected positive clones can be expanded and their monoclonal antibody product harvested for use. Detailed procedures for monoclonal antibody production are described in Davis *et al.* in *Basic Methods in Molecular Biology* Elsevier, New York. Section 21-2, the disclosure of which is incorporated herein by reference.

## 2. Polyclonal Antibody Production by Immunization

Polyclonal antiserum containing antibodies to heterogenous epitopes of a single protein can be prepared by immunizing suitable animals with the expressed protein or peptides derived therefrom, which can be unmodified or modified to enhance immunogenicity. Effective polyclonal antibody production is affected by many factors related both to the antigen and the host species. For example, small molecules tend to be less immunogenic than others and may require the use of carriers and adjuvant. Also, host animals response vary depending on site of inoculations and doses, with both inadequate or excessive doses of antigen resulting in low titer antisera. Small doses (ng level) of antigen administered at multiple intradermal sites appears to be most reliable. An effective immunization protocol for rabbits can be found in Vaitukaitis, *et al.*, *J. Clin. Endocrinol. Metab.* 33:988-991 (1971), the disclosure of which is incorporated herein by reference.

Booster injections can be given at regular intervals, and antiserum harvested when antibody titer thereof, as determined semi-quantitatively, for example, by double immunodiffusion in agar against known concentrations of the antigen, begins to fall. See, for example, Ouchterlony, *et al.*, Chap. 19 in: *Handbook of Experimental Immunology* D. Wier (ed) Blackwell (1973), the disclosure of which is incorporated herein by reference. Plateau concentration of antibody is usually in the range of 0.1 to 0.2 mg/ml of serum (about 12  $\mu$ M). Affinity of the antisera for the antigen is determined by preparing competitive binding curves,

as described, for example, by Fisher, D., Chap. 42 in: *Manual of Clinical Immunology*, 2d Ed. (Rose and Friedman, Eds.) Amer. Soc. For Microbiol., Washington, D.C. (1980), the disclosure of which is incorporated herein by reference..

Antibody preparations prepared according to either protocol are useful in  
5 quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

10

#### V. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof as Reagents

The 5' ESTs of the present invention (or cDNAs or genomic DNAs obtainable therefrom) may be used as reagents in isolation procedures, diagnostic assays, and  
15 forensic procedures. For example, sequences from the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be detectably labeled and used as probes to isolate other sequences capable of hybridizing to them. In addition, sequences from 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be used to design PCR primers to be used in isolation, diagnostic, or forensic procedures.

20

#### 1. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof in Isolation, Diagnostic and Forensic Procedures

#### EXAMPLE 44

25

#### Preparation of PCR Primers and Amplification of DNA

The 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) may be used to prepare PCR primers for a variety of applications, including isolation procedures for cloning nucleic acids capable of hybridizing to such sequences, diagnostic techniques and forensic techniques. The PCR primers are at least 10 bases, and preferably at least 12, 15, or  
30 17 bases in length. More preferably, the PCR primers are at least 20-30 bases in length. In some embodiments, the PCR primers may be more than 30 bases in length. It is preferred

that the primer pairs have approximately the same G/C ratio, so that melting temperatures are approximately the same. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see *Molecular Cloning to Genetic Engineering*, White Ed. in *Methods in Molecular Biology* 67: Humana Press, Totowa 1997, the disclosure of which is incorporated herein by reference. In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites.

#### EXAMPLE 45

##### 15      Use of 5'ESTs as Probes

Probes derived from 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom), including full length cDNAs or genomic sequences, may be labeled with detectable labels familiar to those skilled in the art, including radioisotopes and non-radioactive labels, to provide a detectable probe. The detectable probe may be single stranded or double stranded and may be made using techniques known in the art, including *in vitro* transcription, nick translation, or kinase reactions. A nucleic acid sample containing a sequence capable of hybridizing to the labeled probe is contacted with the labeled probe. If the nucleic acid in the sample is double stranded, it may be denatured prior to contacting the probe. In some applications, the nucleic acid sample may be immobilized on a surface such as a nitrocellulose or nylon membrane. The nucleic acid sample may comprise nucleic acids obtained from a variety of sources, including genomic DNA, cDNA libraries, RNA, or tissue samples.

Procedures used to detect the presence of nucleic acids capable of hybridizing to the detectable probe include well known techniques such as Southern blotting, Northern blotting, dot blotting, colony hybridization, and plaque hybridization. In some applications, the nucleic acid capable of hybridizing to the labeled probe may be cloned into vectors such as expression vectors, sequencing vectors, or *in vitro* transcription vectors to facilitate the characterization

and expression of the hybridizing nucleic acids in the sample. For example, such techniques may be used to isolate and clone sequences in a genomic library or cDNA library which are capable of hybridizing to the detectable probe as described in Example 30 above.

5 PCR primers made as described in Example 44 above may be used in forensic analyses, such as the DNA fingerprinting techniques described in Examples 46-50 below. Such analyses may utilize detectable probes or primers based on the sequences of the the 5' ESTs or of cDNAs or genomic DNAs isolated using the 5' ESTs.

#### EXAMPLE 46

##### 10 Forensic Matching by DNA Sequencing

In one exemplary method, DNA samples are isolated from forensic specimens of, for example, hair, semen, blood or skin cells by conventional methods. A panel of PCR primers based on a number of the 5' ESTs of Example 25, or cDNAs or genomic DNAs isolated therefrom as described above, is then utilized in accordance with Example 44 to amplify DNA  
15 of approximately 100-200 bases in length from the forensic specimen. Corresponding sequences are obtained from a test subject. Each of these identification DNAs is then sequenced using standard techniques, and a simple database comparison determines the differences, if any, between the sequences from the subject and those from the sample. Statistically significant differences between the suspect's DNA sequences and those from the  
20 sample conclusively prove a lack of identity. This lack of identity can be proven, for example, with only one sequence. Identity, on the other hand, should be demonstrated with a large number of sequences, all matching. Preferably, a minimum of 50 statistically identical sequences of 100 bases in length are used to prove identity between the suspect and the sample.

25

#### EXAMPLE 47

##### Positive Identification by DNA Sequencing

The technique outlined in the previous example may also be used on a larger scale to provide a unique fingerprint-type identification of any individual. In this technique, primers  
30 are prepared from a large number of 5'EST sequences from Example 25, or cDNA or genomic DNA sequences obtainable therefrom. Preferably, 20 to 50 different primers are



used. These primers are used to obtain a corresponding number of PCR-generated DNA segments from the individual in question in accordance with Example 44. Each of these DNA segments is sequenced, using the methods set forth in Example 46. The database of sequences generated through this procedure uniquely identifies the individual from whom the sequences were obtained. The same panel of primers may then be used at any later time to absolutely correlate tissue or other biological specimen with that individual.

#### EXAMPLE 48

##### Southern Blot Forensic Identification

The procedure of Example 47 is repeated to obtain a panel of at least 10 amplified sequences from an individual and a specimen. Preferably, the panel contains at least 50 amplified sequences. More preferably, the panel contains 100 amplified sequences. In some embodiments, the panel contains 200 amplified sequences. This PCR-generated DNA is then digested with one or a combination of, preferably, four base specific restriction enzymes. Such enzymes are commercially available and known to those of skill in the art. After digestion, the resultant gene fragments are size separated in multiple duplicate wells on an agarose gel and transferred to nitrocellulose using Southern blotting techniques well known to those with skill in the art. For a review of Southern blotting see Davis *et al.* (Basic Methods in Molecular Biology, 1986, Elsevier Press. pp 62-65), the disclosure of which is incorporated herein by reference.

A panel of probes based on the sequences of 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom), or fragments thereof of at least 10 bases, are radioactively or colorimetrically labeled using methods known in the art, such as nick translation or end labeling, and hybridized to the Southern blot using techniques known in the art (Davis *et al.*, supra). Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom). More preferably, the probe comprises at least 20-30 consecutive nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom). In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST (or cDNAs or genomic DNAs obtainable therefrom).

Preferably, at least 5 to 10 of these labeled probes are used, and more preferably at least about 20 or 30 are used to provide a unique pattern. The resultant bands appearing

from the hybridization of a large sample of 5' EST (or cDNAs or genomic DNAs obtainable therefrom) will be a unique identifier. Since the restriction enzyme cleavage will be different for every individual, the band pattern on the Southern blot will also be unique. Increasing the number of 5' EST (or cDNAs or genomic DNAs obtainable therefrom) probes will provide a statistically higher level of confidence in the identification since there will be an increased number of sets of bands used for identification.

#### EXAMPLE 49

##### Dot Blot Identification Procedure

Another technique for identifying individuals using the 5' EST sequences disclosed herein utilizes a dot blot hybridization technique.

Genomic DNA is isolated from nuclei of subject to be identified. Oligonucleotide probes of approximately 30 bp in length are synthesized that correspond to at least 10, preferably 50 sequences from the 5' ESTs or cDNAs or genomic DNAs obtainable therefrom. The probes are used to hybridize to the genomic DNA through conditions known to those in the art. The oligonucleotides are end labeled with  $P^{32}$  using polynucleotide kinase (Pharmacia). Dot Blots are created by spotting the genomic DNA onto nitrocellulose or the like using a vacuum dot blot manifold (BioRad, Richmond California). The nitrocellulose filter containing the genomic sequences is baked or UV linked to the filter, prehybridized and hybridized with labeled probe using techniques known in the art (Davis *et al.*, *supra*). The  $^{32}P$  labeled DNA fragments are sequentially hybridized with successively stringent conditions to detect minimal differences between the 30 bp sequence and the DNA. Tetramethylammonium chloride is useful for identifying clones containing small numbers of nucleotide mismatches (Wood *et al.*, *Proc. Natl. Acad. Sci. USA* **82**(6):1585-1588, 1985) which is hereby incorporated by reference. A unique pattern of dots distinguishes one individual from another individual.

5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) or oligonucleotides containing at least 10 consecutive bases from these sequences can be used as probes in the following alternative fingerprinting technique. Preferably, the probe comprises at least 12, 15, or 17 consecutive nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom). More preferably, the probe comprises at least 20-30

consecutive nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom). In some embodiments, the probe comprises more than 30 nucleotides from the 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom).

Preferably, a plurality of probes having sequences from different genes are used in the alternative fingerprinting technique. Example 50 below provides a representative alternative fingerprinting procedure in which the probes are derived from 5'EST.

#### EXAMPLE 50

##### Alternative "Fingerprint" Identification Technique

20-mer oligonucleotides are prepared from a large number, e.g. 50, 100, or 200, of 5'EST using commercially available oligonucleotide services such as Genset, Paris, France. Cell samples from the test subject are processed for DNA using techniques well known to those with skill in the art. The nucleic acid is digested with restriction enzymes such as EcoRI and XbaI. Following digestion, samples are applied to wells for electrophoresis. The procedure, as known in the art, may be modified to accommodate polyacrylamide electrophoresis, however in this example, samples containing 5 ug of DNA are loaded into wells and separated on 0.8% agarose gels. The gels are transferred onto nitrocellulose using standard Southern blotting techniques.

10 ng of each of the oligonucleotides are pooled and end-labeled with  $^{32}\text{P}$ . The nitrocellulose is prehybridized with blocking solution and hybridized with the labeled probes. Following hybridization and washing, the nitrocellulose filter is exposed to X-Omat AR X-ray film. The resulting hybridization pattern will be unique for each individual.

It is additionally contemplated within this example that the number of probe sequences used can be varied for additional accuracy or clarity.

The proteins encoded by the extended cDNAs may also be used to generate antibodies as explained in Examples 30 and 43 in order to identify the tissue type or cell species from which a sample is derived as described in example 51.

**EXAMPLE 51****Identification of Tissue Types or Cell Species by Means of  
Labeled Tissue Specific Antibodies**

Identification of specific tissues is accomplished by the visualization of tissue specific  
5 antigens by means of antibody preparations according to Examples 30 and 43 which are  
conjugated, directly or indirectly to a detectable marker. Selected labeled antibody species  
bind to their specific antigen binding partner in tissue sections, cell suspensions, or in extracts  
of soluble proteins from a tissue sample to provide a pattern for qualitative or semi-qualitative  
interpretation.

10 Antisera for these procedures must have a potency exceeding that of the native  
preparation, and for that reason, antibodies are concentrated to a mg/ml level by isolation  
of the gamma globulin fraction, for example, by ion-exchange chromatography or by  
ammonium sulfate fractionation. Also, to provide the most specific antisera, unwanted  
antibodies, for example to common proteins, must be removed from the gamma globulin  
15 fraction, for example by means of insoluble immunoabsorbents, before the antibodies are  
labeled with the marker. Either monoclonal or heterologous antisera is suitable for either  
procedure.

***A. Immunohistochemical techniques***

Purified, high-titer antibodies, prepared as described above, are conjugated to a  
20 detectable marker, as described, for example, by Fudenberg, Chap. 26 in: *Basic and Clinical  
Immunology*, 3rd Ed. Lange, Los Altos, California, 1980, or Rose, *et al.*, Chap. 12 in:  
*Methods in Immunodiagnosis*, 2d Ed. John Wiley and Sons, New York (1980), the  
disclosures of which are incorporated herein by reference.

A fluorescent marker, either fluorescein or rhodamine, is preferred, but antibodies can  
25 also be labeled with an enzyme that supports a color producing reaction with a substrate, such  
as horseradish peroxidase. Markers can be added to tissue-bound antibody in a second step,  
as described below. Alternatively, the specific antitissue antibodies can be labeled with ferritin  
or other electron dense particles, and localization of the ferritin coupled antigen-antibody  
complexes achieved by means of an electron microscope. In yet another approach, the  
30 antibodies are radiolabeled, with, for example  $^{125}\text{I}$ , and detected by overlaying the antibody  
treated preparation with photographic emulsion.

Preparations to carry out the procedures can comprise monoclonal or polyclonal antibodies to a single protein or peptide identified as specific to a tissue type, for example, brain tissue, or antibody preparations to several antigenically distinct tissue specific antigens can be used in panels, independently or in mixtures, as required.

5 Tissue sections and cell suspensions are prepared for immunohistochemical examination according to common histological techniques. Multiple cryostat sections (about 4  $\mu\text{m}$ , unfixed) of the unknown tissue and known control, are mounted and each slide covered with different dilutions of the antibody preparation. Sections of known and unknown tissues should also be treated with preparations to provide a positive control, a negative  
10 control, for example, pre-immune sera, and a control for non-specific staining, for example, buffer.

Treated sections are incubated in a humid chamber for 30 min at room temperature, rinsed, then washed in buffer for 30-45 min. Excess fluid is blotted away, and the marker developed.

15 If the tissue specific antibody was not labeled in the first incubation, it can be labeled at this time in a second antibody-antibody reaction, for example, by adding fluorescein- or enzyme-conjugated antibody against the immunoglobulin class of the antiserum-producing species, for example, fluorescein labeled antibody to mouse IgG. Such labeled sera are commercially available.

20 The antigen found in the tissues by the above procedure can be quantified by measuring the intensity of color or fluorescence on the tissue section, and calibrating that signal using appropriate standards.

#### *B. Identification of tissue specific soluble proteins*

The visualization of tissue specific proteins and identification of unknown tissues from  
25 that procedure is carried out using the labeled antibody reagents and detection strategy as described for immunohistochemistry; however the sample is prepared according to an electrophoretic technique to distribute the proteins extracted from the tissue in an orderly array on the basis of molecular weight for detection.

A tissue sample is homogenized using a Virtis apparatus; cell suspensions are  
30 disrupted by Dounce homogenization or osmotic lysis, using detergents in either case as required to disrupt cell membranes, as is the practice in the art. Insoluble cell components

such as nuclei, microsomes, and membrane fragments are removed by ultracentrifugation, and the soluble protein-containing fraction concentrated if necessary and reserved for analysis.

A sample of the soluble protein solution is resolved into individual protein species by conventional SDS polyacrylamide electrophoresis as described, for example, by Davis, *et al.*,  
5 Section 19-2 in: *Basic Methods in Molecular Biology*, Leder ed., Elsevier, New York, 1986, the disclosure of which is incorporated herein by reference, using a range of amounts of polyacrylamide in a set of gels to resolve the entire molecular weight range of proteins to be detected in the sample. A size marker is run in parallel for purposes of estimating molecular weights of the constituent proteins. Sample size for analysis is a convenient volume of from 5  
10 to 55  $\mu$ l, and containing from about 1 to 100  $\mu$ g protein. An aliquot of each of the resolved proteins is transferred by blotting to a nitrocellulose filter paper, a process that maintains the pattern of resolution. Multiple copies are prepared. The procedure, known as Western Blot Analysis, is well described in Davis, L. *et al.*, *supra* Section 19-3. One set of nitrocellulose blots is stained with Coomassie blue dye to visualize the entire set of proteins for comparison  
15 with the antibody bound proteins. The remaining nitrocellulose filters are then incubated with a solution of one or more specific antisera to tissue specific proteins prepared as described in Examples 30 and 43. In this procedure, as in procedure A above, appropriate positive and negative sample and reagent controls are run.

In either procedure A or B, a detectable label can be attached to the primary tissue  
20 antigen-primary antibody complex according to various strategies and permutations thereof. In a straightforward approach, the primary specific antibody can be labeled; alternatively, the unlabeled complex can be bound by a labeled secondary anti-IgG antibody. In other approaches, either the primary or secondary antibody is conjugated to a biotin molecule, which can, in a subsequent step, bind an avidin conjugated marker. According to yet another  
25 strategy, enzyme labeled or radioactive protein A, which has the property of binding to any IgG, is bound in a final step to either the primary or secondary antibody.

The visualization of tissue specific antigen binding at levels above those seen in control tissues to one or more tissue specific antibodies, prepared from the gene sequences identified from extended cDNA sequences, can identify tissues of unknown origin, for  
30 example, forensic samples, or differentiated tumor tissue that has metastasized to foreign bodily sites.

In addition to their applications in forensics and identification, 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be mapped to their chromosomal locations. Example 52 below describes radiation hybrid (RH) mapping of human chromosomal regions using 5'ESTs. Example 53 below describes a representative procedure for mapping an 5' EST to its location on a human chromosome. Example 54 below describes mapping of 5' ESTs on metaphase chromosomes by Fluorescence In Situ Hybridization (FISH). Those skilled in the art will appreciate that the method of Examples 52-54 may also be used to map cDNAs or genomic DNAs obtainable from the 5' ESTs to their chromosomal locations.

10

2. Use of 5' ESTs or Sequences Obtainable Therefrom or Portions Thereof in Chromosome Mapping

**EXAMPLE 52**

Radiation hybrid mapping of 5'ESTs to the human genome

15 Radiation hybrid (RH) mapping is a somatic cell genetic approach that can be used for high resolution mapping of the human genome. In this approach, cell lines containing one or more human chromosomes are lethally irradiated, breaking each chromosome into fragments whose size depends on the radiation dose. These fragments are rescued by fusion with cultured rodent cells, yielding subclones containing different portions of the human  
20 genome. This technique is described by Benham *et al.*, *Genomics* 4:509-517, 1989; and Cox *et al.*, *Science* 250:245-250, 1990, the entire contents of which are hereby incorporated by reference. The random and independent nature of the subclones permits efficient mapping of any human genome marker. Human DNA isolated from a panel of 80-100 cell lines provides a mapping reagent for ordering 5'EST. In this approach, the frequency of breakage between  
25 markers is used to measure distance, allowing construction of fine resolution maps as has been done using conventional ESTs (Schuler *et al.*, *Science* 274:540-546, 1996, hereby incorporated by reference).

RH mapping has been used to generate a high-resolution whole genome radiation hybrid map of human chromosome 17q22-q25.3 across the genes for growth hormone (GH)  
30 and thymidine kinase (TK) (Foster *et al.*, *Genomics* 33:185-192, 1996), the region surrounding the Gorlin syndrome gene (Obermayr *et al.*, *Eur. J. Hum. Genet.* 4:242-245,

1996), 60 loci covering the entire short arm of chromosome 12 (Raeymaekers *et al.*, *Genomics* 29:170-178, 1995), the region of human chromosome 22 containing the neurofibromatosis type 2 locus (Frazer *et al.*, *Genomics* 14:574-584, 1992) and 13 loci on the long arm of chromosome 5 (Warrington *et al.*, *Genomics* 11:701-708, 1991).

5

### EXAMPLE 53

#### Mapping of 5'ESTs to Human Chromosomes using PCR techniques

5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) may be assigned to human chromosomes using PCR based methodologies. In such approaches, oligonucleotide primer pairs are designed from the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) to minimize the chance of amplifying through an intron. Preferably, the oligonucleotide primers are 18-23 bp in length and are designed for PCR amplification. The creation of PCR primers from known sequences is well known to those with skill in the art. For a review of PCR technology see Erlich in *PCR Technology, Principles and Applications for DNA Amplification*, Freeman and Co., New York, 1992, the disclosure of which is incorporated herein by reference.

The primers are used in polymerase chain reactions (PCR) to amplify templates from total human genomic DNA. PCR conditions are as follows: 60 ng of genomic DNA is used as a template for PCR with 80 ng of each oligonucleotide primer, 0.6 unit of Taq polymerase, and 1  $\mu$ Cu of a  $^{32}$ P-labeled deoxycytidine triphosphate. The PCR is performed in a microplate thermocycler (Techne) under the following conditions: 30 cycles of 94°C, 1.4 min; 55°C, 2 min; and 72°C, 2 min; with a final extension at 72°C for 10 min. The amplified products are analyzed on a 6% polyacrylamide sequencing gel and visualized by autoradiography. If the length of the resulting PCR product is identical to the distance between the ends of the primer sequences in the extended cDNA from which the primers are derived, then the PCR reaction is repeated with DNA templates from two panels of human-rodent somatic cell hybrids, BIOS PCRable DNA (BIOS Corporation) and NIGMS Human-Rodent Somatic Cell Hybrid Mapping Panel Number 1 (NIGMS, Camden, NJ).

PCR is used to screen a series of somatic cell hybrid cell lines containing defined sets of human chromosomes for the presence of a given 5' EST (or cDNA or genomic DNA obtainable therefrom). DNA is isolated from the somatic hybrids and used as starting



templates for PCR reactions using the primer pairs from the 5' EST (or cDNA or genomic DNA obtainable therefrom). Only those somatic cell hybrids with chromosomes containing the human gene corresponding to the 5' EST (or cDNA or genomic DNA obtainable therefrom) will yield an amplified fragment. The 5' EST (or cDNA or genomic DNA obtainable therefrom) are assigned to a chromosome by analysis of the segregation pattern of PCR products from the somatic hybrid DNA templates. The single human chromosome present in all cell hybrids that give rise to an amplified fragment is the chromosome containing that 5'EST (or cDNA or genomic DNA obtainable therefrom). For a review of techniques and analysis of results from somatic cell gene mapping experiments, see Ledbetter *et al.*,  
10 *Genomics* 6:475-481, 1990, the disclosure of which is incorporated herein by reference.

#### EXAMPLE 54

##### Mapping of Extended 5' ESTs to Chromosomes Using Fluorescence *In Situ*

##### Hybridization

15 Fluorescence in situ hybridization allows the 5'EST (or cDNA or genomic DNA obtainable therefrom) to be mapped to a particular location on a given chromosome. The chromosomes to be used for fluorescence in situ hybridization techniques may be obtained from a variety of sources including cell cultures, tissues, or whole blood.

In a preferred embodiment, chromosomal localization of an 5'EST (or cDNA or genomic DNA obtainable therefrom) is obtained by FISH as described by Cherif *et al.* (*Proc. Natl. Acad. Sci. U.S.A.*, 87:6639-6643, 1990), the disclosure of which is incorporated herein by reference.. Metaphase chromosomes are prepared from phytohemagglutinin (PHA)-stimulated blood cell donors. PHA-stimulated lymphocytes from healthy males are cultured for 72 h in RPMI-1640 medium. For synchronization, methotrexate (10  $\mu$ M) is added for 17  
20 h, followed by addition of 5-bromodeoxyuridine (5-BrdU, 0.1 mM) for 6 h. Colcemid (1  $\mu$ g/ml) is added for the last 15 min before harvesting the cells. Cells are collected, washed in RPMI, incubated with a hypotonic solution of KCl (75 mM) at 37°C for 15 min and fixed in three changes of methanol:acetic acid (3:1). The cell suspension is dropped onto a glass slide and air dried. The 5'EST (or cDNA or genomic DNA obtainable therefrom) is labeled with  
25 biotin-16 dUTP by nick translation according to the manufacturer's instructions (Bethesda Research Laboratories, Bethesda, MD), purified using a Sephadex G-50 column (Pharmacia,  
30

Upsala, Sweden) and precipitated. Just prior to hybridization, the DNA pellet is dissolved in hybridization buffer (50% formamide, 2 X SSC, 10% dextran sulfate, 1 mg/ml sonicated salmon sperm DNA, pH 7) and the probe is denatured at 70°C for 5-10 min.

Slides kept at -20°C are treated for 1 h at 37°C with RNase A (100 µg/ml), rinsed  
5 three times in 2 X SSC and dehydrated in an ethanol series. Chromosome preparations are denatured in 70% formamide, 2 X SSC for 2 min at 70°C, then dehydrated at 4°C. The slides are treated with proteinase K (10 µg/100 ml in 20 mM Tris-HCl, 2 mM CaCl<sub>2</sub>) at 37°C for 8 min and dehydrated. The hybridization mixture containing the probe is placed on the slide, covered with a coverslip, sealed with rubber cement and incubated overnight in a humid  
10 chamber at 37°C. After hybridization and post-hybridization washes, the biotinylated probe is detected by avidin-FITC and amplified with additional layers of biotinylated goat anti-avidin and avidin-FITC. For chromosomal localization, fluorescent R-bands are obtained as previously described (Cherif *et al.*, *supra*). The slides are observed under a LEICA fluorescence microscope (DMRXA). Chromosomes are counterstained with propidium  
15 iodide and the fluorescent signal of the probe appears as two symmetrical yellow-green spots on both chromatids of the fluorescent R-band chromosome (red). Thus, a particular 5'EST (or cDNA or genomic DNA obtainable therefrom) may be localized to a particular cytogenetic R-band on a given chromosome.

20 Once the 5'EST (or cDNA or genomic DNA obtainable therefrom) have been assigned to particular chromosomes using the techniques described in Examples 52-54 above, they may be utilized to construct a high resolution map of the chromosomes on which they are located or to identify the chromosomes in a sample.

25

### EXAMPLE 55

#### Use of 5'EST to Construct or Expand Chromosome Maps

Chromosome mapping involves assigning a given unique sequence to a particular chromosome as described above. Once the unique sequence has been mapped to a given chromosome, it is ordered relative to other unique sequences located on the same  
30 chromosome. One approach to chromosome mapping utilizes a series of yeast artificial chromosomes (YACs) bearing several thousand long inserts derived from the chromosomes

of the organism from which the extended cDNAs (or genomic DNAs obtainable therefrom) are obtained. This approach is described in Nagaraja *et al.*, *Genome Research* 7:210-222, 1997, the disclosure of which is incorporated herein by reference. Briefly, in this approach each chromosome is broken into overlapping pieces which are inserted into the YAC vector.

5 The YAC inserts are screened using PCR or other methods to determine whether they include the 5'EST (or cDNA or genomic DNA obtainable therefrom) whose position is to be determined. Once an insert has been found which includes the 5'EST (or cDNA or genomic DNA obtainable therefrom), the insert can be analyzed by PCR or other methods to determine whether the insert also contains other sequences known to be on the chromosome

10 or in the region from which the 5'EST (or cDNA or genomic DNA obtainable therefrom) was derived. This process can be repeated for each insert in the YAC library to determine the location of each of the extended cDNAs (or genomic DNAs obtainable therefrom) relative to one another and to other known chromosomal markers. In this way, a high resolution map of the distribution of numerous unique markers along each of the organisms chromosomes may

15 be obtained.

As described in Example 56 below extended cDNAs (or genomic DNAs obtainable therefrom) may also be used to identify genes associated with a particular phenotype, such as hereditary disease or drug response.

20

### 3. Use of 5'ESTs or Sequences Obtained Therefrom or Fragments Thereof in Gene Identification

#### **EXAMPLE 56**

##### Identification of genes associated with hereditary diseases or drug response

25

This example illustrates an approach useful for the association of 5'ESTs (or cDNA or genomic DNA obtainable therefrom) with particular phenotypic characteristics. In this example, a particular 5'EST (or cDNA or genomic DNA obtainable therefrom) is used as a test probe to associate that 5'EST (or cDNA or genomic DNA obtainable therefrom) with a particular phenotypic characteristic.

30

5'ESTs (or cDNA or genomic DNA obtainable therefrom) are mapped to a particular location on a human chromosome using techniques such as those described in Examples 52 and 53 or other techniques known in the art. A search of Mendelian Inheritance in Man (McKusick in *Mendelian Inheritance in Man* (available on line through Johns Hopkins University Welch Medical Library) reveals the region of the human chromosome which contains the 5'EST (or cDNA or genomic DNA obtainable therefrom) to be a very gene rich region containing several known genes and several diseases or phenotypes for which genes have not been identified. The gene corresponding to this 5'EST (or cDNA or genomic DNA obtainable therefrom) thus becomes an immediate candidate for each of these genetic diseases.

Cells from patients with these diseases or phenotypes are isolated and expanded in culture. PCR primers from the 5'EST (or cDNA or genomic DNA obtainable therefrom) are used to screen genomic DNA, mRNA or cDNA obtained from the patients. 5'ESTs (or cDNA or genomic DNA obtainable therefrom) that are not amplified in the patients can be positively associated with a particular disease by further analysis. Alternatively, the PCR analysis may yield fragments of different lengths when the samples are derived from an individual having the phenotype associated with the disease than when the sample is derived from a healthy individual, indicating that the gene containing the 5'EST may be responsible for the genetic disease.

20

#### **VI. Use of 5'EST (or cDNA or Genomic DNA Obtainable Therefrom) to Construct Vectors**

The present 5'ESTs (or cDNA or genomic DNA obtainable therefrom) may also be used to construct secretion vectors capable of directing the secretion of the proteins encoded by genes therein. Such secretion vectors may facilitate the purification or enrichment of the proteins encoded by genes inserted therein by reducing the number of background proteins from which the desired protein must be purified or enriched. Exemplary secretion vectors are described in Example 57 below.

30

## 1. Construction of Secretion Vectors

### **EXAMPLE 57**

#### Construction of Secretion Vectors

The secretion vectors include a promoter capable of directing gene expression in the  
5 host cell, tissue, or organism of interest. Such promoters include the Rous Sarcoma Virus  
promoter, the SV40 promoter, the human cytomegalovirus promoter, and other promoters  
familiar to those skilled in the art.

A signal sequence from a 5' EST (or cDNAs or genomic DNAs obtainable  
therefrom) is operably linked to the promoter such that the mRNA transcribed from the  
10 promoter will direct the translation of the signal peptide. The host cell, tissue, or organism  
may be any cell, tissue, or organism which recognizes the signal peptide encoded by the signal  
sequence in the 5' EST (or cDNA or genomic DNA obtainable therefrom). Suitable hosts  
include mammalian cells, tissues or organisms, avian cells, tissues, or organisms, insect cells,  
tissues or organisms, or yeast.

15 In addition, the secretion vector contains cloning sites for inserting genes encoding  
the proteins which are to be secreted. The cloning sites facilitate the cloning of the insert  
gene in frame with the signal sequence such that a fusion protein in which the signal peptide is  
fused to the protein encoded by the inserted gene is expressed from the mRNA transcribed  
from the promoter. The signal peptide directs the extracellular secretion of the fusion protein.

20 The secretion vector may be DNA or RNA and may integrate into the chromosome  
of the host, be stably maintained as an extrachromosomal replicon in the host, be an artificial  
chromosome, or be transiently present in the host. Many nucleic acid backbones suitable for  
use as secretion vectors are known to those skilled in the art, including retroviral vectors,  
SV40 vectors, Bovine Papilloma Virus vectors, yeast integrating plasmids, yeast episomal  
25 plasmids, yeast artificial chromosomes, human artificial chromosomes, P element vectors,  
baculovirus vectors, or bacterial plasmids capable of being transiently introduced into the  
host.

The secretion vector may also contain a polyA signal such that the polyA signal is  
located downstream of the gene inserted into the secretion vector.

30 After the gene encoding the protein for which secretion is desired is inserted into the  
secretion vector, the secretion vector is introduced into the host cell, tissue, or organism using

calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection, viral particles or as naked DNA. The protein encoded by the inserted gene is then purified or enriched from the supernatant using conventional techniques such as ammonium sulfate precipitation, immunoprecipitation, immunochromatography, size  
5 exclusion chromatography, ion exchange chromatography, and HPLC. Alternatively, the secreted protein may be in a sufficiently enriched or pure state in the supernatant or growth media of the host to permit it to be used for its intended purpose without further enrichment.

The signal sequences may also be inserted into vectors designed for gene therapy. In such vectors, the signal sequence is operably linked to a promoter such that mRNA  
10 transcribed from the promoter encodes the signal peptide. A cloning site is located downstream of the signal sequence such that a gene encoding a protein whose secretion is desired may readily be inserted into the vector and fused to the signal sequence. The vector is introduced into an appropriate host cell. The protein expressed from the promoter is secreted extracellularly, thereby producing a therapeutic effect.

15

The 5' ESTs may also be used to clone sequences located upstream of the 5' ESTs which are capable of regulating gene expression, including promoter sequences, enhancer sequences, and other upstream sequences which influence transcription or translation levels. Once identified and cloned, these upstream regulatory sequences may  
20 be used in expression vectors designed to direct the expression of an inserted gene in a desired spatial, temporal, developmental, or quantitative fashion. Example 58 describes a method for cloning sequences upstream of the extended cDNAs or 5' ESTs.

## 2. Identification of Upstream Sequences With Promoting or Regulatory Activities

25

### EXAMPLE 58

#### Use of Extended cDNAs or 5' ESTs to Clone Upstream Sequences from Genomic DNA

Sequences derived from extended cDNAs or 5' ESTs may be used to isolate the promoters of the corresponding genes using chromosome walking techniques. In one chromosome walking technique, which utilizes the GenomeWalker™ kit available from  
30 Clontech, five complete genomic DNA samples are each digested with a different restriction

enzyme which has a 6 base recognition site and leaves a blunt end. Following digestion, oligonucleotide adapters are ligated to each end of the resulting genomic DNA fragments.

For each of the five genomic DNA libraries, a first PCR reaction is performed according to the manufacturer's instructions (which are incorporated herein by reference) using an outer adaptor primer provided in the kit and an outer gene specific primer. The gene specific primer should be selected to be specific for the extended cDNA or 5' EST of interest and should have a melting temperature, length, and location in the extended cDNA or 5' EST which is consistent with its use in PCR reactions. Each first PCR reaction contains 5 ng of genomic DNA, 5  $\mu$ l of 10X Tth reaction buffer, 0.2 mM of each dNTP, 0.2  $\mu$ M each of outer adaptor primer and outer gene specific primer, 1.1 mM of  $Mg(OAc)_2$ , and 1  $\mu$ l of the Tth polymerase 50X mix in a total volume of 50  $\mu$ l. The reaction cycle for the first PCR reaction is as follows: 1 min - 94°C / 2 sec - 94°C, 3 min - 72°C (7 cycles) / 2 sec - 94°C, 3 min - 67°C (32 cycles) / 5 min - 67°C.

The product of the first PCR reaction is diluted and used as a template for a second PCR reaction according to the manufacturer's instructions using a pair of nested primers which are located internally on the amplicon resulting from the first PCR reaction. For example, 5  $\mu$ l of the reaction product of the first PCR reaction mixture may be diluted 180 times. Reactions are made in a 50  $\mu$ l volume having a composition identical to that of the first PCR reaction except the nested primers are used. The first nested primer is specific for the adaptor, and is provided with the GenomeWalker™ kit. The second nested primer is specific for the particular extended cDNA or 5' EST for which the promoter is to be cloned and should have a melting temperature, length, and location in the extended cDNA or 5' EST which is consistent with its use in PCR reactions. The reaction parameters of the second PCR reaction are as follows: 1 min - 94°C / 2 sec - 94°C, 3 min - 72°C (6 cycles) / 2 sec - 94°C, 3 min - 67°C (25 cycles) / 5 min - 67°C. The product of the second PCR reaction is purified, cloned, and sequenced using standard techniques.

Alternatively, two or more human genomic DNA libraries can be constructed by using two or more restriction enzymes. The digested genomic DNA is cloned into vectors which can be converted into single stranded, circular, or linear DNA. A biotinylated oligonucleotide comprising at least 15 nucleotides from the extended cDNA or 5' EST

sequence is hybridized to the single stranded DNA. Hybrids between the biotinylated oligonucleotide and the single stranded DNA containing the extended cDNA or EST sequence are isolated as described in Example 29 above. Thereafter, the single stranded DNA containing the extended cDNA or EST sequence is released from the beads and  
5 converted into double stranded DNA using a primer specific for the extended cDNA or 5' EST sequence or a primer corresponding to a sequence included in the cloning vector. The resulting double stranded DNA is transformed into bacteria. DNAs containing the 5' EST or extended cDNA sequences are identified by colony PCR or colony hybridization.

10 Once the upstream genomic sequences have been cloned and sequenced as described above, prospective promoters and transcription start sites within the upstream sequences may be identified by comparing the sequences upstream of the extended cDNAs or 5' ESTs with databases containing known transcription start sites, transcription factor binding sites, or promoter sequences.

15 In addition, promoters in the upstream sequences may be identified using promoter reporter vectors as described in Example.

### EXAMPLE 59

#### Identification of Promoters in Cloned Upstream Sequences

20 The genomic sequences upstream of the extended cDNAs or 5' ESTs are cloned into a suitable promoter reporter vector, such as the pSEAP-Basic, pSEAP-Enhancer, p $\beta$ gal-Basic, p $\beta$ gal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline  
25 phosphatase,  $\beta$  galactosidase, or green fluorescent protein. The sequences upstream of the extended cDNAs or 5' ESTs are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert  
30 with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer for



augmenting transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert indicates that a promoter sequence is present in the inserted upstream sequence.

Appropriate host cells for the promoter reporter vectors may be chosen based on the results of the above described determination of expression patterns of the extended cDNAs and ESTs. For example, if the expression pattern analysis indicates that the mRNA corresponding to a particular extended cDNA or 5' EST is expressed in fibroblasts, the promoter reporter vector may be introduced into a human fibroblast cell line.

Promoter sequences within the upstream genomic DNA may be further defined by constructing nested deletions in the upstream DNA using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory sites within the promoter may be identified using site directed mutagenesis or linker scanning to obliterate potential transcription factor binding sites within the promoter individually or in combination. The effects of these mutations on transcription levels may be determined by inserting the mutations into the cloning sites in the promoter reporter vectors.

## EXAMPLE 60

### Cloning and Identification of Promoters

Using the method described in Example 58 above with 5' ESTs, sequences upstream of several genes were obtained. Using the primer pairs GGG AAG ATG GAG ATA GTA TTG CCT G (SEQ ID NO:29) and CTG CCA TGT ACA TGA TAG AGA GAT TC (SEQ ID NO:30), the promoter having the internal designation P13H2 (SEQ ID NO:31) was obtained.

Using the primer pairs GTA CCA GGGG ACT GTG ACC ATT GC (SEQ ID NO:32) and CTG TGA CCA TTG CTC CCA AGA GAG (SEQ ID NO:33), the promoter having the internal designation P15B4 (SEQ ID NO:34) was obtained.

Using the primer pairs CTG GGA TGG AAG GCA CGG TA (SEQ ID NO:35) and GAG ACC ACA CAG CTA GAC AA (SEQ ID NO:36), the promoter having the internal designation P29B6 (SEQ ID NO:37) was obtained.

Figure 4 provides a schematic description of the promoters isolated and the way they are assembled with the corresponding 5' tags. The upstream sequences were screened for the presence of motifs resembling transcription factor binding sites or known transcription start sites using the computer program MatInspector release 2.0, August 1996.

Table VII describes the transcription factor binding sites present in each of these promoters. The columns labeled matrice provides the name of the MatInspector matrix used. The column labeled position provides the 5' position of the promoter site. Numeration of the sequence starts from the transcription site as determined by matching the genomic sequence with the 5' EST sequence. The column labeled "orientation" indicates the DNA strand on which the site is found, with the + strand being the coding strand as determined by matching the genomic sequence with the sequence of the 5' EST. The column labeled "score" provides the MatInspector score found for this site. The column labeled "length" provides the length of the site in nucleotides. The column labeled "sequence" provides the sequence of the site found.

Bacterial clones containing plasmids containing the promoter sequences described above described above are presently stored in the inventor's laboratories under the internal identification numbers provided above. The inserts may be recovered from the deposited materials by growing an aliquot of the appropriate bacterial clone in the appropriate medium. The plasmid DNA can then be isolated using plasmid isolation procedures familiar to those skilled in the art such as alkaline lysis minipreps or large scale alkaline lysis plasmid isolation procedures. If desired the plasmid DNA may be further enriched by centrifugation on a cesium chloride gradient, size exclusion chromatography, or anion exchange chromatography.

The plasmid DNA obtained using these procedures may then be manipulated using standard cloning techniques familiar to those skilled in the art. Alternatively, a PCR can be done with primers designed at both ends of the EST insertion. The PCR product which corresponds to the 5' EST can then be manipulated using standard cloning techniques familiar to those skilled in the art.

The promoters and other regulatory sequences located upstream of the extended cDNAs or 5' ESTs may be used to design expression vectors capable of directing the expression of an inserted gene in a desired spatial, temporal, developmental, or quantitative manner. A promoter capable of directing the desired spatial, temporal, developmental, and quantitative patterns may be selected using the results of the expression analysis described in Example 26 above. For example, if a promoter which confers a high level of expression in muscle is desired, the promoter sequence upstream of an extended cDNA or 5' EST derived from an mRNA which is expressed at a high level in muscle, as determined by the method of Example 26, may be used in the expression vector.

Preferably, the desired promoter is placed near multiple restriction sites to facilitate the cloning of the desired insert downstream of the promoter, such that the promoter is able to drive expression of the inserted gene. The promoter may be inserted in conventional nucleic acid backbones designed for extrachromosomal replication, integration into the host chromosomes or transient expression. Suitable backbones for the present expression vectors include retroviral backbones, backbones from eukaryotic episomes such as SV40 or Bovine Papilloma Virus, backbones from bacterial episomes, or artificial chromosomes.

Preferably, the expression vectors also include a polyA signal downstream of the multiple restriction sites for directing the polyadenylation of mRNA transcribed from the gene inserted into the expression vector.

Following the identification of promoter sequences using the procedures of Examples 58-60, proteins which interact with the promoter may be identified as described in Example 61 below.

#### EXAMPLE 61

##### Identification of Proteins Which Interact with Promoter Sequences, Upstream Regulatory Sequences, or mRNA

Sequences within the promoter region which are likely to bind transcription factors may be identified by homology to known transcription factor binding sites or through conventional mutagenesis or deletion analyses of reporter plasmids containing the promoter sequence. For example, deletions may be made in a reporter plasmid containing the promoter sequence of interest operably linked to an assayable reporter gene. The reporter plasmids

carrying various deletions within the promoter region are transfected into an appropriate host cell and the effects of the deletions on expression levels is assessed. Transcription factor binding sites within the regions in which deletions reduce expression levels may be further localized using site directed mutagenesis, linker scanning analysis, or other techniques familiar to those skilled in the art.

Nucleic acids encoding proteins which interact with sequences in the promoter may be identified using one-hybrid systems such as those described in the manual accompanying the Matchmaker One-Hybrid System kit available from Clontech (Catalog No. K1603-1), the disclosure of which is incorporated herein by reference. Briefly, the Matchmaker One-hybrid system is used as follows. The target sequence for which it is desired to identify binding proteins is cloned upstream of a selectable reporter gene and integrated into the yeast genome. Preferably, multiple copies of the target sequences are inserted into the reporter plasmid in tandem. A library comprised of fusions between cDNAs to be evaluated for the ability to bind to the promoter and the activation domain of a yeast transcription factor, such as GAL4, is transformed into the yeast strain containing the integrated reporter sequence. The yeast are plated on selective media to select cells expressing the selectable marker linked to the promoter sequence. The colonies which grow on the selective media contain genes encoding proteins which bind the target sequence. The inserts in the genes encoding the fusion proteins are further characterized by sequencing. In addition, the inserts may be inserted into expression vectors or *in vitro* transcription vectors. Binding of the polypeptides encoded by the inserts to the promoter DNA may be confirmed by techniques familiar to those skilled in the art, such as gel shift analysis or DNase protection analysis.

## VII. Use of 5' ESTs (or cDNAs or Genomic DNAs Obtainable Therefrom) in Gene Therapy

The present invention also comprises the use of 5'ESTs (or cDNA or genomic DNA obtainable therefrom) in gene therapy strategies, including antisense and triple helix strategies as described in Examples 62 and 63 below. In antisense approaches, nucleic acid sequences complementary to an mRNA are hybridized to the mRNA intracellularly, thereby blocking the expression of the protein encoded by the mRNA. The antisense sequences may prevent gene

expression through a variety of mechanisms. For example, the antisense sequences may inhibit the ability of ribosomes to translate the mRNA. Alternatively, the antisense sequences may block transport of the mRNA from the nucleus to the cytoplasm, thereby limiting the amount of mRNA available for translation. Another mechanism through which antisense sequences may inhibit gene expression is by interfering with mRNA splicing. In yet another strategy, the antisense nucleic acid may be incorporated in a ribozyme capable of specifically cleaving the target mRNA.

## EXAMPLE 62

### 10 Preparation and Use of Antisense Oligonucleotides

The antisense nucleic acid molecules to be used in gene therapy may be either DNA or RNA sequences. They may comprise a sequence complementary to the sequence of the 5'EST (or cDNA or genomic DNA obtainable therefrom). The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex with sufficient stability to inhibit the expression of the mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green *et al.*, *Ann. Rev. Biochem.* 55:569-597, 1986; and Izant and Weintraub, *Cell* 36:1007-1015, 1984, which are hereby incorporated by reference.

15 In some strategies, antisense molecules are obtained from a nucleotide sequence encoding a protein by reversing the orientation of the coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules may be transcribed using *in vitro* transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of the antisense nucleic acids *in vivo* by operably linking DNA containing the antisense sequence to a promoter in an expression vector.

25 Alternatively, oligonucleotides which are complementary to the strand normally transcribed in the cell may be synthesized *in vitro*. Thus, the antisense nucleic acids are complementary to the corresponding mRNA and are capable of hybridizing to the mRNA to create a duplex. In some embodiments, the antisense sequences may contain modified sugar phosphate backbones to increase stability and make them less sensitive to RNase activity.

Examples of modifications suitable for use in antisense strategies are described by Rossi *et al.*, *Pharmacol. Ther.* 50(2):245-254, 1991, which is hereby incorporated by reference.

Various types of antisense oligonucleotides complementary to the sequence of the 5'EST (or cDNA or genomic DNA obtainable therefrom) may be used. In one preferred  
5 embodiment, stable and semi-stable antisense oligonucleotides described in International Application No. PCT WO94/23026, hereby incorporated by reference, are used. In these molecules, the 3' end or both the 3' and 5' ends are engaged in intramolecular hydrogen bonding between complementary base pairs. These molecules are better able to withstand exonuclease attacks and exhibit increased stability compared to conventional antisense  
10 oligonucleotides.

In another preferred embodiment, the antisense oligodeoxynucleotides against herpes simplex virus types 1 and 2 described in International Application No. WO 95/04141, hereby incorporated by reference, are used.

In yet another preferred embodiment, the covalently cross-linked antisense  
15 oligonucleotides described in International Application No. WO 96/31523, hereby incorporated by reference, are used. These double- or single-stranded oligonucleotides comprise one or more, respectively, inter- or intra-oligonucleotide covalent cross-linkages, wherein the linkage consists of an amide bond between a primary amine group of one strand and a carboxyl group of the other strand or of the same strand, respectively, the primary  
20 amine group being directly substituted in the 2' position of the strand nucleotide monosaccharide ring, and the carboxyl group being carried by an aliphatic spacer group substituted on a nucleotide or nucleotide analog of the other strand or the same strand, respectively.

The antisense oligodeoxynucleotides and oligonucleotides disclosed in International  
25 Application No. WO 92/18522, incorporated by reference, may also be used. These molecules are stable to degradation and contain at least one transcription control recognition sequence which binds to control proteins and are effective as decoys therefore. These molecules may contain "hairpin" structures, "dumbbell" structures, "modified dumbbell" structures, "cross-linked" decoy structures and "loop" structures.

30 In another preferred embodiment, the cyclic double-stranded oligonucleotides described in European Patent Application No. 0 572 287 A2, hereby incorporated by

reference are used. These ligated oligonucleotide "dumbbells" contain the binding site for a transcription factor and inhibit expression of the gene under control of the transcription factor by sequestering the factor.

Use of the closed antisense oligonucleotides disclosed in International Application  
5 No. WO 92/19732, hereby incorporated by reference, is also contemplated. Because these molecules have no free ends, they are more resistant to degradation by exonucleases than are conventional oligonucleotides. These oligonucleotides may be multifunctional, interacting with several regions which are not adjacent to the target mRNA.

The appropriate level of antisense nucleic acids required to inhibit gene expression  
10 may be determined using *in vitro* expression analysis. The antisense molecule may be introduced into the cells by diffusion, injection, infection, transfection or h-region-mediated import using procedures known in the art. For example, the antisense nucleic acids can be introduced into the body as a bare or naked oligonucleotide, oligonucleotide encapsulated in lipid, oligonucleotide sequence encapsulated by viral protein, or as an oligonucleotide  
15 operably linked to a promoter contained in an expression vector. The expression vector may be any of a variety of expression vectors known in the art, including retroviral or viral vectors, vectors capable of extrachromosomal replication, or integrating vectors. The vectors may be DNA or RNA.

The antisense molecules are introduced onto cell samples at a number of different  
20 concentrations preferably between  $1 \times 10^{-10}$  M to  $1 \times 10^{-4}$  M. Once the minimum concentration that can adequately control gene expression is identified, the optimized dose is translated into a dosage suitable for use *in vivo*. For example, an inhibiting concentration in culture of  $1 \times 10^{-7}$  translates into a dose of approximately 0.6 mg/kg bodyweight. Levels of oligonucleotide approaching 100 mg/kg bodyweight or higher may be possible after testing the toxicity of the  
25 oligonucleotide in laboratory animals. It is additionally contemplated that cells from the vertebrate are removed, treated with the antisense oligonucleotide, and reintroduced into the vertebrate.

It is further contemplated that the antisense oligonucleotide sequence is incorporated into a ribozyme sequence to enable the antisense to specifically bind and cleave its target  
30 mRNA. For technical applications of ribozyme and antisense oligonucleotides see Rossi *et al.*, *supra*.

In a preferred application of this invention, the polypeptide encoded by the gene is first identified, so that the effectiveness of antisense inhibition on translation can be monitored using techniques that include but are not limited to antibody-mediated tests such as RIAs and ELISA, functional assays, or radiolabeling.

5       The 5' ESTs of the present invention (or cDNAs or genomic DNAs obtainable therefrom) may also be used in gene therapy approaches based on intracellular triple helix formation. Triple helix oligonucleotides are used to inhibit transcription from a genome. They are particularly useful for studying alterations in cell activity as it is associated with a particular gene. The 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom)  
10       of the present invention or, more preferably, a portion of those sequences, can be used to inhibit gene expression in individuals having diseases associated with expression of a particular gene. Similarly, a portion of 5' EST sequences (or cDNAs or genomic DNAs obtainable therefrom) can be used to study the effect of inhibiting transcription of a particular gene within a cell. Traditionally, homopurine sequences were considered the most useful for  
15       triple helix strategies. However, homopyrimidine sequences can also inhibit gene expression. Such homopyrimidine oligonucleotides bind to the major groove at homopurine:homopyrimidine sequences. Thus, both types of sequences from the 5'EST or from the gene corresponding to the 5'EST are contemplated within the scope of this invention.

20

### EXAMPLE 63

#### Preparation and Use of Triple Helix Probes

The sequences of the 5' ESTs (or cDNAs or genomic DNAs obtainable therefrom) are scanned to identify 10-mer to 20-mer homopyrimidine or homopurine stretches which  
25       could be used in triple-helix based strategies for inhibiting gene expression. Following identification of candidate homopyrimidine or homopurine stretches, their efficiency in inhibiting gene expression is assessed by introducing varying amounts of oligonucleotides containing the candidate sequences into tissue culture cells which normally express the target gene. The oligonucleotides may be prepared on an oligonucleotide synthesizer or they may  
30       be purchased commercially from a company specializing in custom oligonucleotide synthesis, such as GENSET, Paris, France.



The oligonucleotides may be introduced into the cells using a variety of methods known to those skilled in the art, including but not limited to calcium phosphate precipitation, DEAE-Dextran, electroporation, liposome-mediated transfection or native uptake.

Treated cells are monitored for altered cell function or reduced gene expression using techniques such as Northern blotting, RNase protection assays, or PCR based strategies to monitor the transcription levels of the target gene in cells which have been treated with the oligonucleotide. The cell functions to be monitored are predicted based upon the homologies of the target gene corresponding to the extended cDNA from which the oligonucleotide was derived with known gene sequences that have been associated with a particular function. The cell functions can also be predicted based on the presence of abnormal physiologies within cells derived from individuals with a particular inherited disease, particularly when the extended cDNA is associated with the disease using techniques described in Example 56.

The oligonucleotides which are effective in inhibiting gene expression in tissue culture cells may then be introduced *in vivo* using the techniques described above and in Example 62 at a dosage calculated based on the *in vitro* results, as described in Example 62.

In some embodiments, the natural (beta) anomers of the oligonucleotide units can be replaced with alpha anomers to render the oligonucleotide more resistant to nucleases. Further, an intercalating agent such as ethidium bromide, or the like, can be attached to the 3' end of the alpha oligonucleotide to stabilize the triple helix. For information on the generation of oligonucleotides suitable for triple helix formation see Griffin *et al.*, *Science* 245:967-971, 1989, which is hereby incorporated by this reference.

#### EXAMPLE 64

##### Use of cDNAs Obtained Using the 5' ESTs to Express an Encoded Protein in a Host Organism

25

The cDNAs obtained as described above using the 5' ESTs of the present invention may also be used to express an encoded protein in a host organism to produce a beneficial effect. In such procedures, the encoded protein may be transiently expressed in the host organism or stably expressed in the host organism. The encoded protein may have any of the activities described above. The encoded protein may be a protein which the host organism

30

lacks or, alternatively, the encoded protein may augment the existing levels of the protein in the host organism.

A full length extended cDNA encoding the signal peptide and the mature protein, or an extended cDNA encoding only the mature protein is introduced into the host organism.

5 The extended cDNA may be introduced into the host organism using a variety of techniques known to those of skill in the art. For example, the extended cDNA may be injected into the host organism as naked DNA such that the encoded protein is expressed in the host organism, thereby producing a beneficial effect.

Alternatively, the extended cDNA may be cloned into an expression vector  
10 downstream of a promoter which is active in the host organism. The expression vector may be any of the expression vectors designed for use in gene therapy, including viral or retroviral vectors. The expression vector may be directly introduced into the host organism such that the encoded protein is expressed in the host organism to produce a beneficial effect. In  
15 another approach, the expression vector may be introduced into cells *in vitro*. Cells containing the expression vector are thereafter selected and introduced into the host organism, where they express the encoded protein to produce a beneficial effect.

#### EXAMPLE 65

##### Use of Signal Peptides Encoded by 5' ESTs or Sequences obtained Therefrom 20 to Import Proteins Into Cells

The short core hydrophobic region (h) of signal peptides encoded by the 5'ESTS or extended cDNAs derived from SEQ ID NOs: 38-270 may also be used as a carrier to import a peptide or a protein of interest, so-called cargo, into tissue culture cells (Lin *et al.*, *J. Biol Chem.*, 270: 14225-14258, 1995; Du *et al.*, *J. Peptide Res.*, 51: 235-243, 1998; Rojas *et al.*,  
25 *Nature Biotech.*, 16: 370-375, 1998).

When cell permeable peptides of limited size (approximately up to 25 amino acids) are to be translocated across cell membrane, chemical synthesis may be used in order to add the h region to either the C-terminus or the N-terminus to the cargo peptide of interest. Alternatively, when longer peptides or proteins are to be imported into cells, nucleic acids can  
30 be genetically engineered, using techniques familiar to those skilled in the art, in order to link the extended cDNA sequence encoding the h region to the 5' or the 3' end of a DNA

sequence-coding for a cargo polypeptide. Such genetically engineered nucleic acids are then translated either *in vitro* or *in vivo* after transfection into appropriate cells, using conventional techniques to produce the resulting cell permeable polypeptide. Suitable hosts cells are then simply incubated with the cell permeable polypeptide which is then translocated across the  
5 membrane.

This method may be applied to study diverse intracellular functions and cellular processes. For instance, it has been used to probe functionally relevant domains of intracellular proteins and to examine protein-protein interactions involved in signal transduction pathways (Lin *et al.*, *supra*; Lin *et al.*, *J. Biol. Chem.*, 271: 5305-5308, 1996;  
10 Rojas *et al.*, *J. Biol. Chem.*, 271: 27456-27461, 1996; Liu *et al.*, *Proc. Natl. Acad. Sci. USA*, 93: 11819-11824, 1996; Rojas *et al.*, *Bioch. Biophys. Res. Commun.*, 234: 675-680, 1997).

Such techniques may be used in cellular therapy to import proteins producing therapeutic effects. For instance, cells isolated from a patient may be treated with imported therapeutic proteins and then re-introduced into the host organism.

15 Alternatively, the h region of signal peptides of the present invention could be used in combination with a nuclear localization signal to deliver nucleic acids into cell nucleus. Such oligonucleotides may be antisense oligonucleotides or oligonucleotides designed to form triple helixes, as described in examples 62 and 63 respectively, in order to inhibit processing and/or maturation of a target cellular RNA.

20 As discussed above, the cDNAs or portions thereof obtained using the 5' ESTs of the present invention can be used for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or  
25 at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus  
30 discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination for expression patterns; to raise anti-protein

antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris *et al.*, *Cell* 75:791-803, 1993, the disclosure of which is hereby incorporated by reference) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins or polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, Fritsch and Maniatis eds., 1989, and *Methods in Enzymology: Guide to Molecular Cloning Techniques*, Academic Press, Berger and Kimmel eds., 1987.

Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid

preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Although this invention has been described in terms of certain preferred embodiments,  
5 other embodiments which will be apparent to those of ordinary skill in the art in view of the disclosure herein are also within the scope of this invention. Accordingly, the scope of the invention is intended to be defined only by reference to the appended claims. All documents cited herein are incorporated herein by reference in their entirety.

Step	Search characteristic		Selection Characteristics		
	Program	Strand	Parameters	Identity (%)	Length (bp)
miscellaneous	blastn	both	S=61 X=16	90	17
tRNA	fasta	both	-	80	60
rRNA	blastn	both	S=108	80	40
miRNA	blastn	both	S=108	80	40
Prokaryotic	blastn	both	S=144	90	40
Fungal	blastn	both	S=144	90	40
Alu	fasta*	both	-	70	40
L1	blastn	both	S=72	70	40
Repeats	blastn	both	S=72	70	40
Promoters	blastn	top	S=54 X=16	90	15†
Vertebrate	fasta*	both	S=108	90	30
ESTs	blastn	both	S=108 X=16	90	30
Proteins	blastx <sup>‡</sup>	top	E = 0.001	-	-

Table 1: Parameters used for each step of EST analysis

\* use "Quick Fast" Database scanner

† alignment further constrained to begin closer than 10bp to EST's end

‡ using BLOSUM62 substitution matrix

TABLE II

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELINE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID38	new	13.2	Testis	51-39-3-H2-PU
ID39	new	12	Testis	51-34-3-F8-PU
ID40	new	11	Testis	51-43-1-C5-PU
ID41	new	10.6	Testis	51-2-4-C4-PU
ID42	new	10.4	Ovary	26-49-1-A5-PU
ID43	new	10.1	Testis	51-3-3-B10-PU
ID44	new	9.8	Testis	51-15-4-A12-PU
ID45	new	9.8	Testis	51-14-1-G6-PU
ID46	new	9.5	Spleen	53-1-4-A1-PU
ID47	new	9.4	Ovary	26-40-1-A11-PU
ID48	new	9.4	Testis	51-19-4-A10-PU
ID49	new	9.2	Ovary	26-25-2-D2-PU
ID50	new	9.2	Testis	51-17-2-C6-PU
ID51	new	9.2	Ovary	26-40-3-A6-PU
ID52	new	9.1	Ovary	26-49-1-A9-PU
ID53	new	9.1	Spleen	20-7-2-D6-PU
ID54	new	9.1	Testis	51-2-1-A11-PU
ID55	new	9	Testis	51-43-3-G3-PU
ID56	new	8.9	Ovary	26-47-2-B1-PU
ID57	new	8.8	Ovary	26-11-1-G8-PU
ID58	new	8.8	Testis	51-37-4-E11-PU
ID59	new	8.7	Ovary	26-25-2-G1-PU
ID60	new	8.5	Testis	51-13-1-F7-PU
ID61	new	8.4	Spleen	20-2-1-D7-PU
ID62	new	8.1	Ovary	26-12-2-B5-PU
ID63	new	8	Testis	51-1-1-G12-PU
ID64	new	7.6	Spleen	20-8-2-F3-PU
ID65	new	7.5	Spleen	20-10-3-D4-PU
ID66	new	7.5	Spleen	20-3-3-G4-PU
ID67	new	7.5	Testis	51-10-3-B6-PU
ID68	new	7.5	Ovary	26-27-3-E8-PU
ID69	new	7.4	Testis	51-44-4-A6-PU
ID70	new	7.3	Testis	51-7-2-A6-PU
ID71	new	7.3	Ovary	26-31-1-D11-PU
ID72	new	7.1	Testis	51-28-2-G1-PU
ID73	new	6.9	Spleen	20-10-1-B12-PU
ID74	new	6.9	Testis	51-39-1-A5-PU
ID75	new	6.9	Ovary	26-23-2-A11-PU
ID76	new	6.9	Testis	51-1-4-C5-PU
ID77	new	6.8	Spleen	53-2-4-D8-PU
ID78	new	6.8	Spleen	20-3-2-C11-PU
ID79	new	6.8	Testis	51-29-4-B4-PU
ID80	new	6.8	Ovary	26-27-3-E11-PU
ID81	new	6.6	Ovary	26-10-1-H8-PU
ID82	new	6.5	Testis	51-18-2-G10-PU
ID83	new	6.5	Spleen	20-2-1-H12-PU
ID84	new	6.4	Testis	51-10-3-G3-PU
ID85	new	6.4	Uterus	74-9-4-H2-PU
ID86	new	6.4	Ovary	26-23-3-G2-PU
ID87	new	6.4	Testis	51-2-4-F5-PU
ID88	new	6.4	Uterus	74-4-3-C4-PU

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELNE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID89	new	6.3	Testis	51-31-3-D1-PU
ID90	new	6.3	Spleen	20-5-1-H1-PU
ID91	new	6.2	Ovary	26-41-1-G3-PU
ID92	new	6.2	Uterus	74-11-4-G3-PU
ID93	new	6.1	Ovary	26-4-4-E9-PU
ID94	new	6.1	Spleen	20-2-3-C2-PU
ID95	new	6.1	Ovary	26-48-1-A9-PU
ID96	new	6	Spleen	20-1-2-C7-PU
ID97	new	6	Ovary	26-28-4-H1-PU
ID98	new	6	Uterus	74-8-4-C11-PU
ID99	new	6	Ovary	26-6-3-B9-PU
ID100	new	5.9	Testis	51-16-4-B10-PU
ID101	new	5.9	Testis	51-47-3-F9-PU
ID102	new	5.9	Testis	51-4-2-D10-PU
ID103	new	5.9	Ovary	26-10-4-D9-PU
ID104	new	5.8	Testis	51-18-1-C3-PU
ID105	new	5.8	Ovary	26-45-2-C4-PU
ID106	new	5.7	Ovary	26-26-3-D7-PU
ID107	new	5.7	Ovary	26-5-3-A8-PU
ID108	new	5.7	Ovary	26-47-1-C6-PU
ID109	new	5.6	Testis	51-19-1-F10-PU
ID110	new	5.6	Testis	51-11-4-G10-PU
ID111	new	5.5	Testis	51-39-3-F7-PU
ID112	new	5.5	Testis	51-2-1-E10-PU
ID113	new	5.4	Testis	51-26-2-F5-PU
ID114	new	5.4	Ovary	26-2-2-G10-PU
ID115	new	5.4	Testis	51-35-4-G9-PU
ID116	new	5.4	Ovary	26-39-1-A6-PU
ID117	new	5.3	Ovary	26-47-1-E2-PU
ID118	new	5.3	Testis	51-26-2-C7-PU
ID119	new	5.2	Uterus	74-11-3-F8-PU
ID120	new	5.2	Spleen	53-3-1-E2-PU
ID121	new	5.2	Testis	51-31-3-G12-PU
ID122	new	5.1	Spleen	20-6-4-G5-PU
ID123	new	5.1	Uterus	74-6-3-F1-PU
ID124	new	5.1	Uterus	74-11-1-F8-PU
ID125	new	5.1	Ovary	26-7-4-B3-PU
ID126	new	5	Ovary	26-5-3-F10-PU
ID127	new	5	Ovary	26-49-3-C2-PU
ID128	new	5	Testis	51-29-3-E1-PU
ID129	new	5	Ovary	26-26-3-D2-PU
ID130	new	5	Uterus	74-9-4-B4-PU
ID131	new	5	Testis	51-1-3-E9-PU
ID132	new	4.9	Ovary	26-5-1-C6-PU
ID133	new	4.9	Ovary	26-3-1-H5-PU
ID134	new	4.9	Ovary	26-51-4-D9-PU
ID135	new	4.9	Ovary	26-27-3-D7-PU
ID136	new	4.8	Uterus	74-3-4-D8-PU
ID137	new	4.8	Ovary	26-29-1-E1-PU
ID138	new	4.8	Spleen	20-3-1-H3-PU
ID139	new	4.8	Testis	51-3-3-D8-PU
ID140	new	4.8	Spleen	20-5-3-D9-PU
ID141	new	4.7	Testis	51-44-4-H4-PU



<u>SEQ. ID</u> <u>NO.</u>	<u>CATEGORY</u>	<u>VON HELME</u> <u>SCORE</u>	<u>TISSUE</u> <u>SOURCE</u>	<u>INTERNAL</u> <u>DESIGNATION</u>
ID142	new	4.7	Testis	51-5-4-G12-PU
ID143	new	4.7	Spleen	20-9-2-F7-PU
ID144	new	4.7	Spleen	53-3-2-A10-PU
ID145	new	4.6	Ovary	26-30-4-C1-PU
ID146	new	4.6	Testis	51-29-3-H6-PU
ID147	new	4.6	Testis	51-5-3-G2-PU
ID148	new	4.6	Testis	51-11-3-D5-PU
ID149	new	4.6	Testis	51-7-1-E7-PU
ID150	new	4.6	Testis	51-27-1-G12-PU
ID151	new	4.6	Uterus	74-4-1-F5-PU
ID152	new	4.5	Ovary	26-24-1-F8-PU
ID153	new	4.5	Spleen	20-7-3-F6-PU
ID154	new	4.5	Ovary	26-1-2-A8-PU
ID155	new	4.4	Testis	51-1-3-H9-PU
ID156	new	4.4	Testis	51-27-1-E8-PU
ID157	new	4.3	Testis	51-44-4-B2-PU
ID158	new	4.3	Ovary	26-44-1-C3-PU
ID159	new	4.3	Spleen	20-4-2-E2-PU
ID160	new	4.3	Testis	51-19-4-F5-PU
ID161	new	4.3	Spleen	20-8-4-D7-PU
ID162	new	4.3	Testis	51-24-1-B11-PU
ID163	new	4.3	Spleen	20-6-2-G10-PU
ID164	new	4.2	Testis	51-6-4-F8-PU
ID165	new	4.2	Testis	51-36-2-A9-PU
ID166	new	4.2	Ovary	26-7-3-H10-PU
ID167	new	4.2	Testis	51-1-3-D9-PU
ID168	new	4.2	Spleen	20-2-1-B11-PU
ID169	new	4.2	Uterus	74-6-4-A5-PU
ID170	new	4.2	Testis	51-14-3-F3-PU
ID171	new	4.1	Ovary	26-33-3-E2-PU
ID172	new	4	Testis	51-26-4-C7-PU
ID173	new	4	Testis	51-25-3-F3-PU
ID174	new	4	Ovary	26-8-3-D5-PU
ID175	new	4	Testis	51-42-3-F9-PU
ID176	new	4	Ovary	26-27-1-C5-PU
ID177	new	4	Ovary	26-1-1-G2-PU
ID178	new	3.9	Ovary	26-8-3-H3-PU
ID179	new	3.9	Ovary	26-40-2-A9-PU
ID180	new	3.9	Ovary	26-24-4-A5-PU
ID181	new	3.9	Uterus	74-5-3-B12-PU
ID182	new	3.8	Testis	51-37-2-G12-PU
ID183	new	3.8	Spleen	20-8-2-E7-PU
ID184	new	3.8	Testis	51-2-1-H9-PU
ID185	new	3.8	Ovary	26-46-4-D12-PU
ID186	new	3.8	Ovary	26-40-1-A12-PU
ID187	new	3.7	Testis	51-3-4-E2-PU
ID188	new	3.7	Ovary	26-47-3-G12-PU
ID189	new	3.7	Ovary	26-2-4-E12-PU
ID190	new	3.7	Uterus	74-4-4-D6-PU
ID191	new	3.7	Testis	51-36-4-A3-PU
ID192	new	3.7	Uterus	74-11-1-B8-PU
ID193	new	3.7	Spleen	20-10-2-G2-PU
ID194	new	3.7	Testis	51-37-4-D6-PU

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELME SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID195	new	3.6	Ovary	26-27-4-G9-PU
ID196	new	3.6	Testis	51-2-3-A6-PU
ID197	new	3.6	Ovary	26-24-2-A3-PU
ID198	new	3.6	Uterus	74-3-3-F6-PU
ID199	new	3.5	Spleen	20-10-2-B2-PU
ID200	new	3.5	Testis	51-13-2-G2-PU
ID201	new	3.5	Testis	51-17-4-A4-PU
ID202	new	3.5	Spleen	20-10-3-E5-PU
ID203	new	3.5	Testis	51-30-1-B6-PU
ID204	new	3.5	Ovary	26-40-2-G12-PU
ID205	new	3.5	Ovary	26-9-3-G4-PU
ID206	ext-est-not-vrt	12.7	Testis	51-18-4-A4-PU
ID207	ext-est-not-vrt	7.4	Ovary	26-44-1-B5-PU
ID208	ext-est-not-vrt	7.3	Testis	51-20-1-A2-PU
ID209	ext-est-not-vrt	7.1	Ovary	26-2-1-A12-PU
ID210	ext-est-not-vrt	6.7	Testis	51-2-1-A7-PU
ID211	ext-est-not-vrt	5.6	Spleen	53-1-1-C10-PU
ID212	ext-est-not-vrt	5.6	Uterus	74-10-1-B10-PU
ID213	ext-est-not-vrt	5.3	Testis	51-31-4-A1-PU
ID214	ext-est-not-vrt	4.4	Testis	51-25-1-A2-PU
ID215	ext-est-not-vrt	4.1	Testis	51-35-2-F8-PU
ID216	ext-est-not-vrt	3.9	Testis	51-8-3-E7-PU
ID217	ext-est-not-vrt	3.9	Testis	51-34-2-H6-PU
ID218	ext-est-not-vrt	3.5	Uterus	74-7-2-F11-PU
ID219	est-not-ext	10.5	Testis	51-18-1-G7-PU
ID220	est-not-ext	9.5	Testis	51-23-1-G1-PU
ID221	est-not-ext	8.3	Ovary	26-8-1-B12-PU
ID222	est-not-ext	8.3	Testis	51-41-1-F10-PU
ID223	est-not-ext	8.2	Ovary	26-12-1-A2-PU
ID224	est-not-ext	8.1	Spleen	53-3-3-B8-PU
ID225	est-not-ext	8	Testis	51-4-4-A12-PU
ID226	est-not-ext	7.8	Testis	51-18-1-H7-PU
ID227	est-not-ext	7.6	Spleen	20-6-4-G3-PU
ID228	est-not-ext	7.5	Testis	51-2-3-F10-PU
ID229	est-not-ext	7.1	Testis	51-7-2-C2-PU
ID230	est-not-ext	7.1	Testis	51-6-4-F9-PU
ID231	est-not-ext	6.5	Spleen	20-6-1-D11-PU
ID232	est-not-ext	6.4	Ovary	26-26-1-A11-PU
ID233	est-not-ext	6.4	Testis	51-9-3-A12-PU
ID234	est-not-ext	6.2	Ovary	26-8-3-F5-PU
ID235	est-not-ext	6.1	Ovary	26-27-2-A12-PU
ID236	est-not-ext	6	Uterus	74-11-3-H4-PU
ID237	est-not-ext	5.8	Ovary	26-51-2-G10-PU
ID238	est-not-ext	5.8	Testis	51-23-1-G2-PU
ID239	est-not-ext	5.7	Uterus	74-1-2-H1-PU
ID240	est-not-ext	5.7	Testis	51-9-1-E7-PU
ID241	est-not-ext	5.3	Testis	51-1-4-E9-PU
ID242	est-not-ext	4.8	Testis	51-6-4-G2-PU
ID243	est-not-ext	4.8	Spleen	20-2-1-C5-PU
ID244	est-not-ext	4.7	Testis	51-23-1-H2-PU
ID245	est-not-ext	4.6	Testis	51-19-3-H6-PU
ID246	est-not-ext	4.6	Testis	51-10-3-D11-PU
ID247	est-not-ext	4.6	Testis	51-20-2-G7-PU

<u>SEQ. ID NO.</u>	<u>CATEGORY</u>	<u>VON HELNE SCORE</u>	<u>TISSUE SOURCE</u>	<u>INTERNAL DESIGNATION</u>
ID248	est-not-ext	4.6	Ovary	26-38-4-C2-PU
ID249	est-not-ext	4.5	Ovary	26-44-3-C5-PU
ID250	est-not-ext	4.4	Ovary	26-47-4-H1-PU
ID251	est-not-ext	4.4	Spleen	20-5-2-C3-PU
ID252	est-not-ext	4.3	Testis	51-21-3-B10-PU
ID253	est-not-ext	4.3	Spleen	20-4-4-B3-PU
ID254	est-not-ext	4.2	Ovary	26-5-1-F8-PU
ID255	est-not-ext	4.1	Testis	51-22-3-B10-PU
ID256	est-not-ext	4.1	Testis	51-18-1-G1-PU
ID257	est-not-ext	4.1	Testis	51-12-2-H4-PU
ID258	est-not-ext	3.9	Testis	51-25-1-A12-PU
ID259	est-not-ext	3.8	Spleen	20-2-1-B4-PU
ID260	est-not-ext	3.8	Spleen	20-7-2-A6-PU
ID261	est-not-ext	3.8	Ovary	26-27-4-D3-PU
ID262	est-not-ext	3.8	Ovary	26-5-4-F9-PU
ID263	est-not-ext	3.8	Uterus	74-3-1-B9-PU
ID264	est-not-ext	3.7	Spleen	20-8-4-A11-PU
ID265	est-not-ext	3.6	Testis	51-15-4-G10-PU
ID266	est-not-ext	3.6	Testis	51-2-1-A10-PU
ID267	est-not-ext	3.5	Spleen	53-1-1-A10-PU
ID268	est-not-ext	3.5	Testis	51-15-4-H10-PU
ID269	ext-vrt-not-genomic	8.1	Ovary	26-36-1-D11-PU
ID270	ext-vrt-not-genomic	4	Testis	51-39-2-D9-PU

TABLE III

SEQ. ID NO.	SIGNAL PEPTIDE
ID38	MGEASPPAPARRHLLVLLLLSTLVPSAA
ID39	MAPQTLLPVLVLCVLLLQAQG
ID40	MWTLKSSLVLLLCLTCSYA
ID41	MLPLLLPLWGGSLQ
ID42	METGALRRPQLPLLLLLCGPSQDQC
ID43	MERLVLTLCTLPLAVA
ID44	MMLPQWLLLLFLLFFFLTRG
ID45	MKPVLPLQXLVVFCLALQLVPG
ID46	MFRQRQETAQRSTQSCRCPRDGLFFSLFSAPLASA
ID47	MGSSACEIAVGTKRLLLALPLALVLG
ID48	MSNQRLPLIFSLFICFFGESFC
ID49	MLWFLSFLALLSLNC
ID50	MLXISLEIXSFICCVTVLISLSWT
ID51	MVFRNCILFILTFSSHFTFC
ID52	MLAACPLSPGCQS
ID53	MAWSPLFLTILHCTVSWA
ID54	MLKSVLVSLCSWSPPLTS
ID55	MTSKXLVSFILAALSSTIFS
ID56	MKSLSLXLAVXLGLATAVSA
ID57	MWAMESGHLLWALLFMQSLWP
ID58	MAQTWAXLLVMGSLPSASWS
ID59	MKCGFLAYLLITLLYVWPVINA
ID60	MRKPAAGFLPSLLKVLPLAPAAA
ID61	MRQSLLFLTSVVPFVLA
ID62	MELSOMSELMGLSVLLGLLALMATA
ID63	MQDAPLSCLSPKWSVSSADSTEKSASAAGTRNLPFQFCLRQALRMKAAGILTLIGCLV TGVES
ID64	MALAFCLMAEAILLFSPEHSLFFFCSRKARIRLHWAGQTLAILCAALGLGFISSRTRS ELPHLVSWHSWVGALTLLATAVQALCGLCLLCPRAA
ID65	MLRFPTCFPSXRVXGKQLPQEI XLVWSPXRD XIXLANTAGEVLLHRLASFHRVWS
ID66	MFMVLEVVSRTSSLAMLSDSFHMLSDVLALVVALVAERFA
ID67	MENQLWHNTVRCCNQYQESPHDAEDILLLLGLIVLVNI
ID68	MLSXKITLLTSPNSVCC
ID69	MEGPRGWL VLCVLAISLA
ID70	MKSLLFTLAVFMLLAQLVSG
ID71	MLKLILFSLLISTVC
ID72	MTPWCLACLGRPLASLQWSLTLAWC
ID73	MTMRHNWTPDLSPLWVLLCAHVVTL
ID74	MTGNNRDLFCATLSCMPATS
ID75	MTMRHNWTPDLSPLWVLLCAHVVTL
ID76	MKPLLETLYLLGMLVPGGLG
ID77	MNQADPRLRAVCLWLTSAAMSRGDNCTDLLALGIPSITQAWGLWVLLGAVTLLFLISLA AHL SQ
ID78	MHRQISFLLLRKPRKNWFCQNHVNLKRYLLSLSLTMVIC
ID79	MKQWLCWVLRLEGRQGLGVGEPRGLRLCLGALSAXTFVSFLHA
ID80	MRLGLCFWVPHRGEMSFSSHYSRGTWYQWDSLLMLTLISWFRWCLPAVSTVELLFLFP ILFIRS
ID81	MDFWEEYRRGDVPFWSWCPIRS YLMSVCPVTGKVNHLVKVASARFLHQVTIFPFLYSVK ANYCFLNFDVPQYAWEHSFAAPSILVILVITTSACSA
ID82	MSTSSSSWDNLESLSLSTVWNWQA
ID83	MVFATIGFSLKSGLALGSAGLLWCLA

SEQ. ID NO.	SIGNAL PEPTIDE
ID84	MVLLLSGSVSVGVC
ID85	MCSQKRAVSNQGLMDLGLCXLCXVXNVFA
ID86	MLIVLTLHSPSCDT
ID87	MTRLCLPRPEAREDPIPVPPRGLGAGEGSGSPVRPPVSTWGPSWAQLLDSVLWLGALGLTIQ
ID88	MVLTCLFLSLISTYP
ID89	MLIPVFSFSLQLLSSST
ID90	MAAAXLSGPSAGSAAGVPGGTGGLSAVSSGPRLRLLLESVSGLLQP
ID91	MHNWLFVFTFCNC
ID92	MHVECFYFLSTALGSQA
ID93	MSPGSALALLWSPASDLG
ID94	MALALGPSISSIA
ID95	MLAFLFCTLFSLVVHP
ID96	MAQMPLTGSYQDLEYFLECMFLHLLYTLQTISSLSG
ID97	MALLMGLWVRTVLQG
ID98	MINHLYLAILDXSLKLITG
ID99	MGRQGTLEIEGILCVITWLEANLGKQKDENVHYYKKLSLLYLCSPPLPGTS
ID100	MELTNKQTGTDREHQLRRVKQDKRISAWWCVLLEWSQG
ID101	MAKRQNPTSVLGLFSISDTWA
ID102	MNVLPFSYYYILFCLSLQIFRVSLA
ID103	MKCLKVNPFLFLVFNFFSYISXFLSPVCG
ID104	MSWTVPVVRASQVRSSVGANXCLGMALCPRQA
ID105	MGFLXLMTLTTHVHS
ID106	MLFRVLLLAQLFLGSG
ID107	MRVPEDLASKILLPGCAPGSLPLSTSAPPLRG
ID108	MFPHXETQVKCFWQGLRRSDLCLCQCILARA
ID109	MKSLLFTLAVFMDLAQLVSG
ID110	MHLYSCSCMRLLNVACCIPFSSS
ID111	MRAPLVLSPLSYQCSS
ID112	MQVPHLRVWTQVXDTFIGYRNLGFTSMCILFHCLLS
ID113	MQKLMAVPMITRAQGGDTCTROJLWLMHQSFQKSNS
ID114	MCXAGFXDHPRAARHARTSRHPLPWVCVSQXPAHRSLCLWPACLC
ID115	MTSKFILVSFILAALSLS
ID116	MHLIFILTVHHTPS
ID117	MLSSSLMVQLISQVYS
ID118	MFSYILCMLFCLFS
ID119	MLFLYYVTLAFSLLVLSES
ID120	MLLSGLWLSSVKEC
ID121	MVAFSVFCFSWLMSSSSP
ID122	MVPLALGIGPPGCLQG
ID123	MNLCMGVLLKVGTSRRCLCLLWFCTAMRPGGA
ID124	MSLAKSLFLRVARG
ID125	MRLPPFLPSATLLLSAES
ID126	MSDRKRTKFSYVQLPCPISLLPRSFKRGOIPGPSAPPLLLLREELVTG
ID127	MTPLGSGPPREASIAQVRGFSRTFFRVAFCFFPAFLVXVS
ID128	MRCALFPLLSLLSC
ID129	MLYDQYYLIISLLKLCSCFI
ID130	MANCFLSHKSQTILISKPALTSQSHFTSPAGFLTVEKSHLLTRLFFHWLSVLCSFLSLRFCTLS
ID131	MHGAGLTYLLFLPDWAAV
ID132	MCCLSATLAFSGSFL
ID133	MAELDLMAPGPLPRATAQPPAPLSPDSGLRGLLLQEALG
ID134	MTLTHGNNILHLANFFLVACPLFGVCLX
ID135	MVLRWLPWPRGSHS
ID136	MKARLSGNLICFSFLGTLFHKSNS

<u>SEQ. ID</u> <u>NO.</u>	<u>SIGNAL PEPTIDE</u>
ID137	MSHVCLVPQTPSLCLG
ID138	MYPASFVKIPSTAYVVLTSVNLFIGING
ID139	MSSSRKDHLGAXAQSPSRSSLWVTAPLVSA
ID140	MASPAAATYLVQSSACCPA
ID141	MNAAJNTGPAPAVTKTETEVQNPDLWDLDIPEARSHADQDSNPXAEALLPCNLHXSWLHS
ID142	MINLLVGNCTYLLGAIRASCMCRXMSFAKFGIFLVFCSEFS
ID143	MLCCGPLRFLLRDPGCLLA
ID144	MRKTSFILLRMTVLPWLWT
ID145	MWWKPAPEEGVRVGLVLVXRALC
ID146	MFNLLGNSSCVYQ
ID147	MKRGAFSNLNDSQLSASFLQPSLQANCPALDPAVSLSAFA
ID148	MKSAGLGFLLRFFIFCSLNTLLG
ID149	MDILFPLHSVIGSHP
ID150	MLKVFRAXHPKICHFGILLLSQRQWS
ID151	MLVRNARRGSRGRSPWWRAGCLXWRKLAASWTLS
ID152	MTKGHHHQHPLHPHPLFTLGLGYPIPTL
ID153	MTYHXIQFSERLHILFIVCLARG
ID154	MSQFPLCSPWKPLVKVSRNLKIRMSIPWPLSVLIYCGLSQPLTLG
ID155	MFRSLTTAFRDAMGFLMFDLTSQ
ID156	MVLTTLPLPSANSPVNMPTTGPNSLSYASSALSPCLX
ID157	MQRNATFIHLQLAIRPSLLPTLPWLPSTRL
ID158	MNLCFHSFHPFLQ
ID159	MLTNRNYFNFLVLVQLCILA
ID160	MKLNPGQVPTWWEALCRFVGMQPCTA
ID161	MLAGFRRSAPASQSLCLNLCPCSSSL
ID162	MKEGASFYLLFFLNDVPP
ID163	MGLECCCPHNLRVYIETLLKLSSQSRT
ID164	MQLCPFTSVLSIAASLLQCRL
ID165	MDVTCFDDAVEGSDFRVCCHGCVSWLCLQMLQLLFLKNSTWCRA
ID166	MRQGPAPLHCFCTFLFSYSSS
ID167	MHTLLGIWLTXLQ
ID168	MLYGSWVCLLSAGTAFE
ID169	MLFFPLLSFRFLPSESLLKXCCFLGRRVVG
ID170	MPVWAILGCWGTLSRG
ID171	MGMSGKKHFLPSWDHIQGSTAETSQGLCGSLPGPSLC
ID172	MASKILLNVQEEVTCPICLELLTEPLSLDCGHSLCRA
ID173	MYYMVCLFFRLIFS
ID174	MGAGGXREIRAAAASWLRAAEHSLAGLWSPGLVPA
ID175	MGSKCKGGPDEDAVERQRRQKLLLAQLHHRKRVKAAGQIQAWWRGVLVVRTLLVAA LRA
ID176	MQQGHPLSAGTLSIHSWQLLTSAPQ
ID177	MSRYEXGSSLLPFPDHSVYSFKXXSFEEAYSISDYATCCLSLFQWCAV
ID178	MIYFIKINNKLHLLHHYLLFTT
ID179	MELLYLKVKRGQKDLISWALCLSQSGYY
ID180	MTLAVTSLALGATG
ID181	MLGPPLQPGSHGKVLAPQGSSGLTPFPCCRCLITLPRSCR
ID182	MGNVCSCLLRARYQQLXLILVHFPAYS
ID183	MLYGLGSGPRCVISCHGVWC
ID184	MHRIMTLHLKALQQLQNKIHVPRMLPGPVTPLDSCPPSAHS
ID185	MLFLVLFYSAIFL
ID186	MVSLCVAALFPLQA
ID187	MSSNLFYTPSILTLLA
ID188	MGLLRKCFVMLGGNTHIQITCIKQFILCLGTCRG

<u>SEQ. ID</u> <u>NO.</u>	<u>SIGNAL PEPTIDE</u>
ID189	MMLPLFCSPWESGG
ID190	MAKLLSDLSVDSARC
ID191	MCGYWVCWGHLLPARVST
ID192	MKLSCAGCADTAILGLSTFLNLLS
ID193	MIPFSGTVFSLGSCPAGPLSA
ID194	MIPSSQPRFXNPACKQTVLLXDPAVSLSAFASA
ID195	MAPTFLLISDSFLT
ID196	MISLIVLSLLGIQWCLS
ID197	MACDSFLKDALPQELSQLXFLFPLVDMREDLLYFNTFLPRKVA
ID198	MLLLNENLKAEIQKNEAQGSCILFLFCFESQNMRSKSIFPFLILHFFPQQIRK
ID199	MISKYVHYSLTDLLLPTFLSLKAFL
ID200	MARTMGVPRACKAFCSLLSFCALHFG
ID201	MILCFLLPHRLQEA
ID202	MQDYVSHA VRRHCQCFVCFSPKIYG
ID203	MEFAHAAECVSFALNETHVLLNLALSHFNNC
ID204	MGNQGFPLYSPSLSVQDLLAASWLPRDAPC
ID205	MKYQMVSGSAQLASPLPGATP
ID206	MGPSTPLLILFLLSWGPLQG
ID207	MASLGHILVFCVGLLTMAKA
ID208	MSGSSLPSALALSLLVSGSLLP
ID209	MMEVVVGNGVVALRGIPRTSRKSSRKTRFCGERGSKQSGKCSVGLAVVSLGGSRG
ID210	MARCFSLVLLLTSTWT
ID211	MGSRKCGGCLSCLLIPLALWS
ID212	MGSRKCGGCLSCLLIPLALWS
ID213	MMVMILFGVSFVFLTHC
ID214	MSNTHTVLVSLPHHPALT
ID215	MXVYRLQTOEKPNTTVQVPAFLQELVDRDNSKFEEWCIEMAEMRXKVVWKEKQNTKRLRS CTKGYLELSPMSLSLWNGCKSGWMNQXPNLLIITLACVPMTSFT
ID216	MFPVLGWILIAVVIILLIFT
ID217	MFSCCISVCLCPCLNKGQS
ID218	MRLCLIMYCSFGTLSHLTLYLLLSPIKYP
ID219	MGKGMVAMLILGLLLALLLPVQVSS
ID220	MGSSGLLSLLVLFVLLANVQG
ID221	MVLGGCPVSYLLLCGQAALLGNLLLLHCVSRSHS
ID222	METGRLLSLSSLPLVLLG
ID223	MAASLGQVLALVLVAALWG
ID224	MHIKSHLEGFKSYAQRTEVNGFDPLFNATGLNGSGKSNILDSICFLGISNLSQVRA
ID225	MSPSPRWGFLCVLFTAVHP
ID226	MCSELYPLVTFFLLCLCIAYWAST
ID227	MLPFLFFSTLFSSIFT
ID228	MVALNLILVPCCAA
ID229	MAARGVIAPVGESLRYAEYLQPSAKRPDADVDQQLVRSLIAVGLGVAALAF
ID230	MIKLKLLSLLRPSLC
ID231	MPSVNSAGLCVLQTTAVTS
ID232	MMLGLHFALFLLVSXYMIRS
ID233	MALLSVLRVLLG
ID234	MLKSLWLSLVAWHWGEA
ID235	MGIVTWLLXSFMSA
ID236	MAGIKALISLSFGGAIGLMFLMLGCALP
ID237	MKKQKHQKLWCISVKLVTLSPVPSLA
ID238	MDGIPMSMKNEMPISQLLMIAPSLGFVLFALFVAFLLRG
ID239	MGGFLHLPALSSSCLWTFPPMCVRISYVPLPILTPKTINLIPVLAICSLPGPGPA
ID240	MSPSPRWGFLCVLFTAVHP

<u>SEQ. ID</u> <u>NO.</u>	<u>SIGNAL PEPTIDE</u>
ID241	MTSQPVPNETIIVLPSNVINFSQAEKPEPTNQGQDSLKKHLHAEXKVIGTIQLCGMMVL SLGHILASAFSPNFT
ID242	MRALENDFFNSPPRKTVRFGGTVEVLLKYKKGETNDFELLKNQLLDPDIKDDQIINWLL EFRSSVMYLTKDQFEQLISILRLPWLNRST
ID243	MVFPKRFCLVPSMEGVRWAFSCGTWLPSTRA
ID244	MNCFQGTNASALEKDIGPEQFPINEHYFGLVNFNTCYCNSVLQALYSCRPFRENVLAYK AQKKKENLLTCLADLFHSIAT
ID245	MAAALRVRXXFGTRA
ID246	MKLLTHNLLSSHVRG
ID247	MGXFSRRTFCGRSGRSCRGQLVQVSRPEVSAGSLLLPAQA
ID248	MEGGVRLDLSACGETSGVAVSELPASETAALVPEGHGPGLRACALSLPDAPGASG
ID249	MTLLSFAALTAAFS
ID250	MAAATGDPGLSKLQFAPFSSA
ID251	MFTSTGSSGLYKAPLSKSLLLVPSXLS
ID252	MTSMTQSLREVAMTKARNFERVLGKITLVSAAPGKVIC
ID253	MADFGISAGQFVAVVWDKSSPVEALKGLVDKLQALTGNEGRVSVENIKQLLSAHKESX DIILSGLVPGSTT
ID254	MGILLGLLLLGHLT
ID255	MFLTVKLLLGQRCSLKVSG
ID256	MNVIDHVRDMAAAGLHSNVRLSSLLLTMSNN
ID257	MGTPSLSILLIGAPESPIPYFPHYSGTGRVLCPLLXAAAAP
ID258	MVYHALDSPDDDYHALFVLCCLYAMS
ID259	MFIVLSMWLCCGFE
ID260	MVVVILSSXVPLAAM
ID261	MLAECSSLLHPSVRG
ID262	MQMARLLGLCAWARK
ID263	MTPQYLPHGGKYQVLGDYSLAVVFPLHFSDLVLYLIPKTLT
ID264	MVVLRAKKTFPLPLXRAFACRG
ID265	MKREGGAHLCSDSLPEQQ
ID266	MVTCPGPSSGQPLSSMYTAGDRRGAPSLPYSLAACPCGSQG
ID267	MQRQLALEVIVTLSETAA
ID268	MGDYLLRGYRMLGETCADCGTILLQDKQRKIYCVACQELSDSDVDKDNPALNAQAALSQAR EHQLASASEPLGSRP
ID269	MWLLYLLVPALFCRA
ID270	MKLEFTKNXXSFVLQNLNRQRKRKEYWDMALSVDNHVFFAHRNVLAAVSPLVRSLS



Minimum signal peptide score	false positive rate	false negative rate	proba(0.1)	proba(0.2)
3.5	0.121	0.036	0.467	0.664
4	0.096	0.06	0.519	0.708
4.5	0.078	0.079	0.565	0.745
5	0.062	0.098	0.615	0.782
5.5	0.05	0.127	0.659	0.813
6	0.04	0.163	0.694	0.836
6.5	0.033	0.202	0.725	0.855
7	0.025	0.248	0.763	0.878
7.5	0.021	0.304	0.78	0.889
8	0.015	0.368	0.816	0.909
8.5	0.012	0.418	0.836	0.92
9	0.009	0.512	0.856	0.93
9.5	0.007	0.581	0.863	0.934
10	0.006	0.679	0.835	0.919

TABLE IV

Minimum signal peptide score	All ESTs	New ESTs	ESTs matching public EST closer than 40 bp from beginning	ESTs extending known mRNA more than 40 bp	ESTs extending public EST more than 40 bp
3.5	2674	947	599	23	150
4	2278	784	499	23	126
4.5	1943	647	425	22	112
5	1657	523	353	21	96
5.5	1417	419	307	19	80
6	1190	340	238	18	68
6.5	1035	280	186	18	60
7	893	219	161	15	48
7.5	753	173	132	12	36
8	636	133	101	11	29
8.5	543	104	83	8	26
9	456	81	63	6	24
9.5	364	57	48	6	18
10	303	47	35	6	15

TABLE V

Tissue	All ESTs	New ESTs	ESTs matching public EST closer than 40 bp from beginning	ESTs extending known mRNA more than 40 bp	ESTs extending public EST more than 40 bp
Brain	329	131	75	3	24
Cancerous prostate	134	40	37	1	6
Cerebellum	17	9	1	0	6
Colon	21	11	4	0	0
Dystrophic muscle	41	18	8	0	1
Fetal brain	70	37	16	0	1
Fetal kidney	227	116	46	1	19
Fetal liver	13	7	2	0	0
Heart	30	15	7	0	1
Hypertrophic prostate	86	23	22	2	2
Kidney	10	7	3	0	0
Large intestine	21	8	4	0	1
Liver	23	9	6	0	0
Lung	24	12	4	0	1
Lung (cells)	57	38	6	0	4
Lymph ganglia	163	60	23	2	12
Lymphocytes	23	6	4	0	2
Muscle	33	16	6	0	4
Normal prostate	181	61	45	7	11
Ovary	90	57	12	1	2
Pancreas	48	11	6	0	1
Placenta	24	5	1	0	0
Prostate	34	16	4	0	2
Spleen	58	28	10	0	1
Substantia nigra	108	47	27	1	6
Surrenals	15	3	3	1	0
Testis	131	68	25	1	8
Thyroid	17	8	2	0	2
Umbilical cord	55	17	12	1	3
Uterus	28	15	3	0	2
Non tissue-specific	568	48	177	2	28
Total	2677	947	601	23	150

TABLE VI

128/4

# Description of Transcription Factor Binding Sites present on promoters Isolated from SignalTag sequences

Promoter sequence P13H2 (646 bp):

Matrix	Position	Orientation	Score	Length	Sequence
CMYB_01	-502	+	0.983	9	TGTCAGTTG
MYOD_Q6	-501	.	0.981	10	CCCAACTGAC
S8_01	-444	.	0.960	11	AATAGAATTAG
S8_01	-425	+	0.968	11	AACTAAATTAG
DELTAEF1_01	-390	.	0.960	11	GCACACCTCAG
GATA_C	-364	.	0.964	11	AGATAAATCCA
CMYB_01	-349	+	0.958	9	CTTCAGTTG
GATA1_02	-343	+	0.959	14	TTGTAGATAGGACA
GATA_C	-339	+	0.953	11	AGATAGGACAT
TAL1ALPHA47_01	-235	+	0.973	16	CATAACAGATGGTAAG
TAL1BETA47_01	-235	+	0.983	16	CATAACAGATGGTAAG
TAL1BETA1F2_01	-235	+	0.978	16	CATAACAGATGGTAAG
MYOD_Q6	-232	.	0.954	10	ACCATCTGTT
GATA1_04	-217	.	0.953	13	TCAAGATAAAGTA
IK1_01	-126	+	0.983	13	AGTTGGGAATTCC
IK2_01	-126	+	0.985	12	AGTTGGGAATTC
CREL_01	-123	+	0.962	10	TGGGAATTCC
GATA1_02	-96	+	0.950	14	TCAGTGATATGGCA
SRY_02	-41	.	0.951	12	TAAACAAAACA
E2F_02	-33	+	0.957	8	TTTAGCGC
MZF1_01	-5	.	0.975	8	TGAGGGGA

Promoter sequence P16B4 (861bp):

Matrix	Position	Orientation	Score	Length	Sequence
NFY_Q6	-748	.	0.956	11	GGACCAATCAT
MZF1_01	-738	+	0.962	8	CCTGGGGA
CMYB_01	-684	+	0.994	9	TGACCGTTG
VMYB_02	-682	.	0.985	9	TCCAACGGT
STAT_01	-673	+	0.968	9	TTCTGGAA
STAT_01	-673	.	0.951	9	TTCCAGGAA
MZF1_01	-556	.	0.956	8	TTGGGGGA
IK2_01	-451	+	0.965	12	GAATGGGATTC
MZF1_01	-424	+	0.986	8	AGAGGGGA
SRY_02	-398	.	0.955	12	GAAAACAAAACA
MZF1_01	-216	+	0.960	8	GAAGGGGA
MYOD_Q6	-190	+	0.981	10	AGCATCTGCC
DELTAEF1_01	-176	+	0.958	11	TCCCACCTTCC
S8_01	5	.	0.992	11	GAGGCAATTAT
MZF1_01	16	.	0.986	8	AGAGGGGA

Promoter sequence P29B6 (666 bp):

Matrix	Position	Orientation	Score	Length	Sequence
ARNT_01	-311	+	0.984	18	GGACTCAGTGCTGCT
NMYC_01	-309	+	0.965	12	ACTCAGTGCTG
USF_01	-309	+	0.985	12	ACTCAGTGCTG
USF_01	-309	.	0.985	12	CAGCACGTGAGT
NMYC_01	-309	.	0.956	12	CAGCACGTGAGT
MYCMAx_02	-309	.	0.972	12	CAGCACGTGAGT
USF_C	-307	+	0.997	8	TCACGTGC
USF_C	-307	.	0.991	8	GCACGTGA
MZF1_01	-292	.	0.968	8	CATGGGGA
ELK1_02	-105	+	0.963	14	CTCTCCGGAAGCCT
CETS1P54_01	-102	+	0.974	10	TCCGGAAGCC
AP1_Q4	-42	.	0.963	11	AGTGACTGAAC
AP1FJ_Q2	-42	.	0.981	11	AGTGACTGAAC
PAOS_C	45	+	1.000	9	TGTGGTCTC

TABLE VII

CLAIMS

1. A purified or isolated nucleic acid comprising the sequence of one of SEQ ID NOs: 38-270 or comprising a sequence complementary thereto.
- 5 2. The nucleic acid of Claim 1, wherein said nucleic acid is recombinant.
3. A purified or isolated nucleic acid comprising at least 10 consecutive bases of the sequence of one of SEQ ID NOs: 38-270 or one of the sequences complementary thereto.
4. A purified or isolated nucleic acid comprising at least 15 consecutive bases of  
10 one of the sequences of SEQ ID NOs: 38-270 or one of the sequences complementary thereto.
5. The nucleic acid of Claim 4, wherein said nucleic acid is recombinant.
6. A purified or isolated nucleic acid of at least 15 bases capable of hybridizing under stringent conditions to the sequence of one of SEQ ID NOs: 38-270 or one of the  
15 sequences complementary to the sequences of SEQ ID NOs: 38-270.
7. The nucleic acid of Claim 6, wherein said nucleic acid is recombinant.
8. A purified or isolated nucleic acid encoding a human gene product, said human gene product having a sequence partially encoded by one of the sequences of SEQ ID NO: 38-270.
- 20 9. A purified or isolated nucleic acid having the sequence of one of SEQ ID NOs: 38-270 or having a sequence complementary thereto.
10. A purified or isolated nucleic acid comprising the nucleotides of one of SEQ ID NOs: 38-270 which encode a signal peptide.
11. A purified or isolated polypeptides comprising a signal peptide encoded by  
25 one of the sequences of SEQ ID NOs: 38-270.
12. A vector encoding a fusion protein comprising a polypeptide and a signal peptide, said vector comprising a first nucleic acid encoding a signal peptide encoded by one of the sequences of SEQ ID NOs: 38-270 operably linked to a second nucleic acid encoding a polypeptide.
- 30 13. A method of directing the extracellular secretion of a polypeptide or the insertion of a polypeptide into the membrane comprising the steps of:

obtaining a vector according to Claim 12; and

introducing said vector into a host cell such that said fusion protein is secreted into the extracellular environment of said host cell or inserted into the membrane of said host cell.

14. A method of importing a polypeptide into a cell comprising contacting said cell with a fusion protein comprising a signal peptide encoded by one of the sequences of SEQ ID NOs: 38-270 operably linked to said polypeptide.

15. A method of making a cDNA encoding a human secretory protein that is partially encoded by one of SEQ ID NOs 38-270, comprising the steps of:

obtaining a cDNA comprising one of the sequences of SEQ ID NOs: 38-270;

10 contacting said cDNA with a detectable probe comprising at least 15 consecutive nucleotides of said sequence of SEQ ID NO: 38-270 or a sequence complementary thereto under conditions which permit said probe to hybridize to said cDNA;

identifying a cDNA which hybridizes to said detectable probe; and

isolating said cDNA which hybridizes to said probe.

15 16. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 15.

17. The cDNA of Claim 16 wherein said cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

18. A method of making a cDNA comprising one of the sequences of SEQ ID NOs: 38-270, comprising the steps of:

contacting a collection of mRNA molecules from human cells with a first primer capable of hybridizing to the polyA tail of said mRNA;

25 hybridizing said first primer to said polyA tail;

reverse transcribing said mRNA to make a first cDNA strand;

making a second cDNA strand complementary to said first cDNA strand using at least one primer comprising at least 15 nucleotides of one of the sequences of SEQ ID NOs 38-270; and

30 isolating the resulting cDNA comprising said first cDNA strand and said second cDNA strand.

19. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270 or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 18.

5 20. The cDNA of Claim 19 wherein said cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

21. The method of Claim 18, wherein the second cDNA strand is made by:

contacting said first cDNA strand with a first pair of primers, said first pair of primers comprising a second primer comprising at least 15 consecutive nucleotides of one of the sequences of SEQ ID NOs 38-270 and a third primer having a sequence therein which is  
10 included within the sequence of said first primer;

performing a first polymerase chain reaction with said first pair of nested primers to generate a first PCR product;

contacting said first PCR product with a second pair of primers, said second pair of primers comprising a fourth primer, said fourth primer comprising at least 15 consecutive  
15 nucleotides of said sequence of one of SEQ ID NOs 38-270, and a fifth primer, said fourth and fifth primers being capable of hybridizing to sequences within said first PCR product; and

performing a second polymerase chain reaction, thereby generating a second PCR product.

20 22. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein encoded by one of SEQ ID NOs 38-270, or a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 21.

23. The cDNA of Claim 22 wherein said cDNA comprises the full protein coding  
25 sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

24. The method of Claim 18 wherein the second cDNA strand is made by:

contacting said first cDNA strand with a second primer comprising at least 15 consecutive nucleotides of the sequences of SEQ ID NOs: 38-270;

hybridizing said second primer to said first strand cDNA; and

30 extending said hybridized second primer to generate said second cDNA strand.

25. An isolated or purified cDNA encoding a human secretory protein, said human secretory protein comprising the protein partially encoded by one of SEQ ID NOs 38-270 or comprising a fragment thereof of at least 10 amino acids, said cDNA being obtainable by the method of Claim 24.

5        26. The cDNA of Claim 25, wherein said cDNA comprises the full protein coding sequence partially included in one of the sequences of SEQ ID NOs: 38-270.

27. A method of making a protein comprising one of the sequences of SEQ ID NO: 271-503, comprising the steps of:

obtaining a cDNA encoding the full protein sequence partially included in one of the  
10 sequences of sequence of SEQ ID NO: 38-270;

inserting said cDNA in an expression vector such that said cDNA is operably linked to a promoter;

introducing said expression vector into a host cell whereby said host cell produces the protein encoded by said cDNA; and

15 isolating said protein.

28. An isolated protein obtainable by the method of Claim 27.

29. A method of obtaining a promoter DNA comprising the steps of:

obtaining DNAs located upstream of the nucleic acids of SEQ ID NO: 38-270 or the sequences complementary thereto;

20 screening said upstream DNAs to identify a promoter capable of directing transcription initiation; and

isolating said DNA comprising said identified promoter.

30. The method of Claim 29, wherein said obtaining step comprises chromosome walking from said nucleic acids of SEQ ID NO: 38-270 or sequences complementary thereto.

25 31. The method of Claim 30, wherein said screening step comprises inserting said upstream sequences into a promoter reporter vector.

32. The method of Claim 30, wherein said screening step comprises identifying motifs in said upstream DNAs which are transcription factor binding sites or transcription start sites.

30 33. An isolated promoter obtainable by the method of Claim 32.



34. An isolated or purified protein comprising one of the sequences of SEQ ID NO: 271-503.

35. In an array of discrete ESTs or fragments thereof of at least 15 nucleotides in length, the improvement comprising inclusion in said array of at least one of the sequences of  
5 SEQ ID NOs: 38-270, or one of the sequences complementary to the sequences of SEQ ID NOs: 38-270, or a fragment thereof of at least 15 consecutive nucleotides.

36. The array of Claim 35 including therein at least two of the sequences of SEQ ID NOs: 38-270, the sequences complementary to the sequences of SEQ ID NOs: 38-270, or fragments thereof of at least 15 consecutive nucleotides.

10 37. The array of Claim 35 including therein at least five of the sequences of SEQ ID NOs: 38-270, the sequences complementary to the sequences of SEQ ID NOs: 38-270, or fragments thereof of at least 15 consecutive nucleotides.

1/4




Figure 1

2/4




Figure 2

3/4







Figure 3

**Promoter P13H2****Promoter P15B4****Promoter P29B6****Figure 4**

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME : GENSET SA
- (B) STREET : 24, RUE ROYALE
- (C) CITY: PARIS
- (E) COUNTRY : FRANCE
- (F) POSTAL CODE (ZIP) : 75008

(ii) TITLE OF INVENTION: 5' ESTs FOR SECRETED PROTEINS  
EXPRESSED IN TESTIS AND OTHER TISSUES

(iii) NUMBER OF SEQUENCES: 503

## (v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy Disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: Win95
- (D) SOFTWARE: Word

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

## (ix) FEATURE:

- (A) NAME/KEY: Cap
- (B) LOCATION: 1
- (D) OTHER INFORMATION: m7Gppp added to 1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGCAUCCUAC UCCCAUCCAA UCCACCCUA ACUCCUCCCA UCUCCAC

47

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GGCAUCCUACU GCAUCCAAU UCCACCCUA CUCUCCCAU CUCCAC

46

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

ATCAAGAATT CGCAGGAGAC CATTG

25

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

TAATGGTCTC GTGCGAATTC TTGAT

25

## (2) INFORMATION FOR SEQ ID NO: 5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CCGACAAGAC CAACGTCAAG GCCGC

25

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

TCACCAGCAG GCAGTGGCTT AGGAG

25

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

AGTGATTCTT GCTACTTTGG ATGGC

25

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCTTGGTCTT GTTCTGGAGT TTAGA

25

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TCCAGAATGG GAGACAAGCC AATTT

25

(2) INFORMATION FOR SEQ ID NO: 10:



- (1) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 25 base pairs  
    (B) TYPE: NUCLEIC ACID  
    (C) STRANDEDNESS: SINGLE  
    (D) TOPOLOGY: LINEAR

(11) MOLECULE TYPE: Other nucleic acid

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

AGGGAGGAGG AACAGCGTG AGTCC

25

(2) INFORMATION FOR SEQ ID NO: 11:

- (1) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 25 base pairs  
    (B) TYPE: NUCLEIC ACID  
    (C) STRANDEDNESS: SINGLE  
    (D) TOPOLOGY: LINEAR

(11) MOLECULE TYPE: Other nucleic acid

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

ATGGGAAAGG AAAAGACTCA TATCA

25

(2) INFORMATION FOR SEQ ID NO: 12:

- (1) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 25 base pairs  
    (B) TYPE: NUCLEIC ACID  
    (C) STRANDEDNESS: SINGLE  
    (D) TOPOLOGY: LINEAR

(11) MOLECULE TYPE: Other nucleic acid

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

AGCAGCAACA ATCAGGACAG CACAG

25

(2) INFORMATION FOR SEQ ID NO: 13:

- (1) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 25 base pairs  
    (B) TYPE: NUCLEIC ACID  
    (C) STRANDEDNESS: SINGLE  
    (D) TOPOLOGY: LINEAR

(11) MOLECULE TYPE: Other nucleic acid

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

ATCAAGAATT CGCAGGAGAC CATT

25

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 67 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

ATCGTTGAGA CTCGTACCAG CAGAGTCACG AGAGAGACTA CACGCTACTG GTTTTTTTTTT 60

TTTTTVN 67

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CCAGCAGAST CACGAGAGAG ACTACACGG

29

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

CACGAGAGAG ACTACACGGT ACTGG

25

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 526 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Lymph ganglia

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(261..376)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 96  
 region 166..281  
 id N70479  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(380..486)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
 region 54..160  
 id N70479  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(110..145)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 94  
 region 403..438  
 id N70479  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(195..229)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 94  
 region 315..348  
 id N70479  
 est

(ix) FEATURE:  
 (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 90..140  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8.2  
 seq LLLITAILAVAVG/FP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

```
AAATATPARAG AGCTACAATA TTCCAGGGCC ARTCACTTGC CATTCTCAT AACAGCGTCA      60
GAGATATAATA ACTGACTGRR ACGTTTGAG ATG AAG AAA GTT CTC CTC CTC ATC      113
```

Met Lys Lys Val Leu Leu Leu Ile  
-15 -10

ACA GCC ATC TTG GCA GTG GCT GTW GGT TTC CCA GTC TCT CAA GAC CAG 161  
Thr Ala Ile Leu Ala Val Ala Val Gly Phe Pro Val Ser Gln Asp Gln  
-5 1 5

GAA CGA GAA AAA AGA AGT ATC AGT GAC AGC GAT GAA TTA GCT TCA GGR 209  
Glu Arg Glu Lys Arg Ser Ile Ser Asp Ser Asp Glu Leu Ala Ser Gly  
10 15 20

WTT TTT GTG TTC CCT TAC CCA TAT CCA TTT CGC CCA CTT CCA CCA ATT 257  
Xaa Phe Val Phe Pro Tyr Pro Tyr Pro Phe Arg Pro Leu Pro Pro Ile  
25 30 35

CCA TTT CCA AGA TTT CCA TGG TTT AGA CGT AAN TTT CCT ATT CCA ATA 305  
Pro Phe Pro Arg Phe Pro Trp Phe Arg Arg Xaa Phe Pro Ile Pro Ile  
40 45 50 55

CCT GAA TCT GCC CCT ACA ACT CCC CTT CCT AGC GAA AAG TAAACAARAA 354  
Pro Glu Ser Ala Pro Thr Thr Pro Leu Pro Ser Glu Lys  
60 65

GGAAAAGTCA CRATAAACCT GGTCACCTGA AATTGAAATT GAGCCACTTC CTTGAARAAT 414

CAAAATTCCT GTTAATAAAA RAAAAACAAA TGTAATTGAA ATAGCACACA GCATTCTCTA 474

GTCAATATCT TTAGTGATCT TCTTTAATAA ACATGAAAGC AAAAAAAAAA AA 526

## (2) INFORMATION FOR SEQ ID NO: 18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..17
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.2  
seq LLLITAILAVAVG/FP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Met Lys Lys Val Leu Leu Leu Ile Thr Ala Ile Leu Ala Val Ala Val  
1 5 10 15  
Gly

## (2) INFORMATION FOR SEQ ID NO: 19:

## (ii) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 822 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (iii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (F) TISSUE TYPE: kidney

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 260..464
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 153..357  
id H57434  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 118..184
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 98..164  
id H57434  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 56..113
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 35..92  
id H57434  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 454..485
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 348..379  
id H57434  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 118..545
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..428  
id N27248  
est

## (ix) FEATURE:

- (A) NAME/KEY: other

(B) LOCATION: 65..369  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 41..345  
id H94779  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 61..399  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 99  
region 6..344  
id H09880  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 408..459  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 92  
region 355..405  
id H09880  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 60..399  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 56..395  
id H29351  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 393..432  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 391..430  
id H29351  
est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 346..408  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.5  
seq SFLPSALVIWTS/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

ACTCCTTTTA GCATAGGGGC TTCGGCGCCA GCGGCCAGCG CTAGTCGGTC TGGTAAGTGC	60
GTGATGCCGA GTTCCGTCTC TCGCGTCTTT TCCTGGTCCC AGGCAAAGCG GASGNAGATC	120
GTGAAACGGC CTAGTCGCTC GCGCTTCCGG AGAAAATCAG CCGTCTAATT AATTCCTCTG	180
GTGTGTGAA GCAGTTACCA AGAATCTTCA ACCCTTCCC ACAAAGCTA ATTGAGTACA	240

```

CGTTCCTGTT_GAGTACACGT TCCTGTTGAT TTACAAAAGG TGCAGGTATG AGCAGGTCTG      300
AAGACTAACA TTTTGTGAAG TTGTAAAACA GAAAACCTST TAGAA ATG TGG TGG TTT      357
                                     Met Trp Trp Phe
                                     -20
CAG CAA GGC CTC AGT TTC CTT CCT TCA GCC CTT GTA ATT TGG ACA TCT      405
Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu Val Ile Trp Thr Ser
      -15                      -10                      -5
GCT GCT TTC ATA TTT TCA TAC ATT ACT GCA GTA ACA CTC CAC CAT ATA      453
Ala Ala Phe Ile Phe Ser Tyr Ile Thr Ala Val Thr Leu His His Ile
      1                      5                      10                      15
GAC CCG GCT TTA CCT TAT ATC AGT GAC ACT GGT ACA GTA GCT CCA RAA      501
Asp Pro Ala Leu Pro Tyr Ile Ser Asp Thr Gly Thr Val Ala Pro Xaa
      20                      25                      30
AAA TGC TTA TTT GGG GCA ATG CTA AAT ATT GCG GCA GTT TTA TGT CAA      549
Lys Cys Leu Phe Gly Ala Met Leu Asn Ile Ala Ala Val Leu Cys Gln
      35                      40                      45
AAA TAGAAATCAG GAARATAATT CAACTTAAAG AAKTTCATTT CATGACCAAA      602
Lys
CTCTTCARAA ACATGTCTTT ACAAGCATAT CTCTTGTAAT GCTTTCTACA CTGTTGAATT      662
GTCTGGCAAT ATTTCTGCAG TGGAAAATTT GATTTARMTA GTTCTTGACT GATAAATATG      722
GTAAGGTGGG CTTTCCCCC TGTGTAATTG GCTACTATGT CTTACTGAGC CAAGTTGTAW      732
TTTGAAATAA AATGATATGA GAGTGACACA AAAAAAAAAA      822

```

## (2) INFORMATION FOR SEQ ID NO: 20:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..21
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5  
seq SFLPSALVIWTS/AF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

```

Met Trp Trp Phe Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu Val
1          5          10          15

```

Ile Trp Thr Ser Ala  
20

## (2) INFORMATION FOR SEQ ID NO: 21:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 405 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(103..398)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 1..296  
id AA442893  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 185..295
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq LSYASSALSPCLT/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

```

ATCACCTTCT TCTCCATCCT TSTCTGGGCC AGTCCCCARC CCAGTCCCTC TCCTGACCTG      60
CCCAGCCCAA GTCAGCCTTC AGCAGCGGCT TTTCTGCACA CAGATATTCC AGGCCTACCT      120
GGCATTCCAG GACCTCCGMA ATGATGCTCC AGTCCCTTAC AAGCGCTTCC TGGATGAGGG      180
TGGC ATG GTG CTG ACC ACC CTC CCC TTG CCC TCT GCC AAC AGC CCT GTG      229
Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val
      -35                -30                -25

AAC ATG CCC ACC ACT GGC CCC AAC AGC CTG AGT TAT GCT AGC TCT GCC      277
Asn Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala
      -20                -15                -10

CTG TCC CCC TGT CTG ACC GCT CCA AAK TCC CCC CGG CTT GCT ATG ATG      325
Leu Ser Pro Cys Leu Thr Ala Pro Xaa Ser Pro Arg Leu Ala Met Met
      -5                1                5                10

CCT GAC AAC TAAATATCCT TATCCAAATC AATAAARWRA RAATCCTCCC TCARAAGGS      384
Pro Asp Asn

TTTCTAAAAA CAAAAAAA A      405

```



## (2) INFORMATION FOR SEQ ID NO: 22:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..37
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq LSYASSALSPCLT/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val Asn  
1 5 10 15  
Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala Leu  
20 25 30  
Ser Pro Cys Leu Thr  
35

## (2) INFORMATION FOR SEQ ID NO: 23:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Cancerous prostate

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..331
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..123  
id AA397994  
est

## (ix) FEATURE:

- (A) NAME/KEY: other

..(B) LOCATION: 328..465  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 96  
 region 179..336  
 id AA397994  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: complement(182..496)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
 region 14..328  
 id AA399680  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 196..240  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.5  
 seq ILSTVTALTFAXA/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

AAAAAATTGG TCCCAGTTTT CACCCTGCCG CAGGGCTGGC TGGGGAGGGC AGCGGTTTAG	60
ATTAGCCGTG GCCTAGGCCG TTAAACGGGG TGACACGAGC NTGCAGGGCC GAGTCCAAGG	120
CCCGGAGATA GGACCAACCG TCAGGAATGC GAGGAATGTT TTTCTTCGGA CTCTATCGAG	180
GCACACAGAC AGACC ATG GGG ATT CTG TCT ACA GTG ACA GCC TTA ACA TTT	231
Met Gly Ile Leu Ser Thr Val Thr Ala Leu Thr Phe	
-15 -10 -5	
GCC ARA GCC CTG GAC GGC TGC AGA AAT GGC ATT GCC CAC CCT GCA AGT	279
Ala Xaa Ala Leu Asp Gly Cys Arg Asn Gly Ile Ala His Pro Ala Ser	
1 5 10	
GAG AAG CAC AGA CTC GAG AAA TGT AGG GAA CTC GAG ASC ASC CAC TCG	327
Glu Lys His Arg Leu Glu Lys Cys Arg Glu Leu Glu Xaa Xaa His Ser	
15 20 25	
GCC CCA GGA TCA ACC CAS CAC CGA AGA AAA ACA ACC AGA AGA AAT TAT	375
Ala Pro Gly Ser Thr Xaa His Arg Arg Lys Thr Thr Arg Arg Asn Tyr	
30 35 40 45	
TCT TCA GCC TGAATGAAK CCGCGATCAA ATGGTTGCTG ATCARAGCCC ATATTTAAAT	434
Ser Ser Ala	
TGGAAAAGTC AAATTGASCA TTATTAAATA AAGCTTGTTT AATATGTCTC AAACAAAAAA	494
AA	496

## (xii) INFORMATION FOR SEQ ID NO: 24:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..15
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5  
seq ILSTVTALTFFAXA/LD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

```
Met Gly Ile Leu Ser Thr Val Thr Ala Leu Thr Phe Ala Xaa Ala
 1           5           10          15
```

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 623 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 49..96
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.1  
seq LVLTLCCTLPLAVA/SA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

```
AAAGATCCCT GCAGCCCGGC AGGAGAGAAG GCTGAGCCTT CTGGCGTC ATG GAG AGG      57
                                     Met Glu Arg
                                     -15

CTC GTC CTA ACC CTG TGC ACC CTC CCG CTG GCT GTG GCG TCT GCT GGC      105
Leu Val Leu Thr Leu Cys Thr Leu Pro Leu Ala Val Ala Ser Ala Gly
-10           -5           1

TGC GCC ACG ACG CCA GCT CGC AAC CTG AGC TGC TAC CAG TGC TTC AAG      153
Cys Ala Thr Thr Pro Ala Arg Asn Leu Ser Cys Tyr Gln Cys Phe Lys
 5           10           15

GTC AGC AGC TGG ACG GAG TGC CCG CCC ACC TGG TGC AGC CCG CTG GAC      201
```

Val	Ser	Ser	Trp	Thr	Glu	Cys	Pro	Pro	Thr	Trp	Cys	Ser	Pro	Leu	Asp	
20					25					30					35	
CAA	GTC	TGC	ATC	TCC	AAC	GAG	GTG	GTC	GTC	TCT	TTT	AAA	TGG	AGT	GTA	249
Ser	Val	Cys	Ile	Ser	Asn	Glu	Val	Val	Val	Ser	Phe	Lys	Trp	Ser	Val	
			40						45					50		
CGC	GTC	CTG	CTC	AGC	AAA	CGC	TGT	GCT	CCC	AGA	TGT	CCC	AAC	GAC	AAC	297
Arg	Val	Leu	Leu	Ser	Lys	Arg	Cys	Ala	Pro	Arg	Cys	Pro	Asn	Asp	Asn	
		55						60					65			
ATG	AAK	TTC	GAA	TGG	TCG	CCG	GCC	CCC	ATG	GTG	CAA	GGC	GTG	ATC	ACC	345
Met	Xaa	Phe	Glu	Trp	Ser	Pro	Ala	Pro	Met	Val	Gln	Gly	Val	Ile	Thr	
		70				75						80				
AGG	CGC	TGC	TGT	TCC	TGG	GCT	CTC	TGC	AAC	AGG	GCA	CTG	ACC	CCA	CAG	393
Arg	Arg	Cys	Cys	Ser	Trp	Ala	Leu	Cys	Asn	Arg	Ala	Leu	Thr	Pro	Gln	
	85					90				95						
GAG	GGG	CGC	TGG	GCC	CTG	CRA	GGG	GGG	CTC	CTG	CTC	CAG	GAC	CCT	TCG	441
Glu	Gly	Arg	Trp	Ala	Leu	Xaa	Gly	Gly	Leu	Leu	Leu	Gln	Asp	Pro	Ser	
100				105					110					115		
AGG	GGC	ARA	AAA	ACC	TGG	GTG	CGG	CCA	CAG	CTG	GGG	CTC	CCA	CTC	TGC	489
Arg	Gly	Xaa	Lys	Thr	Trp	Val	Arg	Pro	Gln	Leu	Gly	Leu	Pro	Leu	Cys	
			120					125					130			
CTT	CCC	AWT	TCC	AAC	CCC	CTC	TGC	CCA	RGG	GAA	ACC	CAG	GAA	GGA		534
Leu	Pro	Xaa	Ser	Asn	Pro	Leu	Cys	Pro	Xaa	Glu	Thr	Gln	Glu	Gly		
		135				140						145				
TAACACTGTG	GGTGCCCCCA	CCTGTGCATT	GGGACCACRA	CTTCACCCTC	TTGGARACAA											594
TAAACTCTCA	TGCCCCCAAA	AAAAAAAAAA														623

## (2) INFORMATION FOR SEQ ID NO: 26:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..16
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.1  
seq LVLTLCTLPLAVA/SA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met	Glu	Arg	Leu	Val	Leu	Thr	Leu	Cys	Thr	Leu	Pro	Leu	Ala	Val	Ala
1				5				10					15		

## (2) INFORMATION FOR SEQ ID NO: 27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 848 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (D) DEVELOPMENTAL STAGE: Fetal
- (E) TISSUE TYPE: kidney

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 32..73
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.7  
seq LWLLFFLVTAIHA/EL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

```

AACTTTGCCT TGTGTTTCC ACCCTGAAAG A ATG TTG TGG CTG CTC TTT TTT CTG      55
                               Met Leu Trp Leu Leu Phe Phe Leu
                               -10

GTG ACT GCC ATT CAT GCT GAA CTC TGT CAA CCA GGT GCA GAA AAT GCT      103
Val Thr Ala Ile His Ala Glu Leu Cys Gln Pro Gly Ala Glu Asn Ala
-5                               1                               5                               10

TTT AAA GTG AGA CTT AGT ATC AGA ACA GCT CTG GGA GAT AAA GCA TAT      151
Phe Lys Val Arg Leu Ser Ile Arg Thr Ala Leu Gly Asp Lys Ala Tyr
15                               20                               25

GCC TGG GAT ACC AAT GAA GAA TAC CTC TTC AAA GCG ATG GTA GCT TTC      199
Ala Trp Asp Thr Asn Glu Glu Tyr Leu Phe Lys Ala Met Val Ala Phe
30                               35                               40

TCC ATG AGA AAA GTT CCC AAC AGA GAA GCA ACA GAA ATT TCC CAT GTC      247
Ser Met Arg Lys Val Pro Asn Arg Glu Ala Thr Glu Ile Ser His Val
45                               50                               55

CTA CTT TGC AAT GTA ACC CAG AGG GTA TCA TTC TGG TTT GTG GTT ACA      295
Leu Leu Cys Asn Val Thr Gln Arg Val Ser Phe Trp Phe Val Val Thr
60                               65                               70

GAC CCT TCA AAA AAT CAC ACC CTT CCT GCT GTT GAG GTG CAA TCA GCC      343
Asp Pro Ser Lys Asn His Thr Leu Pro Ala Val Glu Val Gln Ser Ala
75                               80                               85                               90

ATA AGA ATG AAC AAG AAC CGG ATC AAC AAT GCC TTC TTT CTA AAT GAC      391
Ile Arg Met Asn Lys Asn Arg Ile Asn Asn Ala Phe Phe Leu Asn Asp
95                               100                               105

CAA ACT CTG GAA TTT TTA AAA ATC CCT TCC ACA CTT GCA CCA CCC ATG      439

```

Gln Thr Leu Glu Phe Leu Lys Ile Pro Ser Thr Leu Ala Pro Pro Met  
 110 115 120

GAC CCA TCT GTG CCC ATC TGG ATT ATT ATA TTT GGT GTG ATA TTT TGC 437  
 Asp Pro Ser Val Pro Ile Trp Ile Ile Phe Gly Val Ile Phe Cys  
 125 130 135

ATC ATC ATA GTT GCA ATT GCA CTA CTG ATT TTA TCA GGG ATC TGG CAA 535  
 Ile Ile Ile Val Ala Ile Ala Leu Leu Ile Leu Ser Gly Ile Trp Gln  
 140 145 150

CGT ADA ARA AAG AAC AAA GAA CCA TCT GAA GTG GAT GAC GCT GAA RAT 583  
 Arg Xaa Xaa Lys Asn Lys Glu Pro Ser Glu Val Asp Asp Ala Glu Xaa  
 155 160 165 170

AAK TGT GAA AAC ATG ATC ACA ATT GAA AAT GGC ATC CCC TCT GAT CCC 631  
 Xaa Cys Glu Asn Met Ile Thr Ile Glu Asn Gly Ile Pro Ser Asp Pro  
 175 180 185

CTG GAC ATG AAG GGA GGG CAT ATT AAT GAT GCC TTC ATG ACA GAG GAT 679  
 Leu Asp Met Lys Gly Gly His Ile Asn Asp Ala Phe Met Thr Glu Asp  
 190 195 200

GAG AGG CTC ACC CCT CTC TGAAGGGCTG TTGTCTGCT TCCTCAARAA 727  
 Glu Arg Leu Thr Pro Leu  
 205

ATTAAACATT TGTTTCTGTG TGACTGCTGA GCATCCTGAA ATACCAAGAG CAGATCATAT 787

WTTTGTGTTT ACCATTCTTC TTTTGTAAATA AATTTTGAAT GTGCTTGAAA AAAAAAAAAA 847

C 848

## (2) INFORMATION FOR SEQ ID NO: 28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..14
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.7  
 seq LWLLFFLVTAIHA/EL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

Met Leu Trp Leu Leu Phe Phe Leu Val Thr Ala Ile His Ala  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO: 29:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

GGGAAGATGG AGATAGTATT GCCTG

25

## (2) INFORMATION FOR SEQ ID NO: 30:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CTGCCATGTA CATGATAGAG AGATTG

26

## (2) INFORMATION FOR SEQ ID NO: 31:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 546 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Genomic DNA

## (ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..517

## (ix) FEATURE:

- (A) NAME/KEY: transcription start site
- (B) LOCATION: 518

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 17..25
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name CMY3\_01  
score 0.983  
sequence TGTCAGTTG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(18..27)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MYOD\_Q6  
score 0.961  
sequence CCCAACTGAC

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(75..85)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name S8\_01  
score 0.960  
sequence AATAGAATTAG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 94..104
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name S8\_01  
score 0.966  
sequence AACTAAATTAG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(129..139)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name DELTAEF1\_01  
score 0.960  
sequence GCACACCTCAG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(155..165)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name GATA\_C  
score 0.964  
sequence AGATAAATCCA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 170..178
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name CMYB\_01  
score 0.958  
sequence CTCAGTTG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 176..189
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name GATA1\_02  
score 0.959  
sequence TTGTAGATAGGACA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 180..190
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name GATA\_C



score 0.953  
sequence AGATAGGACAT

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 284..299  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name TAL1ALPHA47\_01  
score 0.973  
sequence CATAACAGATGGTAAG

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 284..299  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name TAL1BETA47\_01  
score 0.983  
sequence CATAACAGATGGTAAG

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 284..299  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name TAL1BETA1TF2\_01  
score 0.978  
sequence CATAACAGATGGTAAG

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(287..296)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name MYOD\_Q6  
score 0.954  
sequence ACCATCTGTT

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(302..314)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name GATA1\_04  
score 0.953  
sequence TCAAGATAAAGTA

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 393..405  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name IK1\_01  
score 0.963  
sequence AGTTGGGAATTCC

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 393..404  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name IK2\_01  
score 0.985  
sequence AGTTGGGAATTCC

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site

(B) LOCATION: 396..405  
 (C) IDENTIFICATION METHOD: matinspector prediction  
 (D) OTHER INFORMATION: name CREL\_01  
 score 0.962  
 sequence TGGGAATTCC

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
 (B) LOCATION: 423..436  
 (C) IDENTIFICATION METHOD: matinspector prediction  
 (D) OTHER INFORMATION: name GATA1\_02  
 score 0.950  
 sequence TCAGTGATATGGCA

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
 (B) LOCATION: complement(478..489)  
 (C) IDENTIFICATION METHOD: matinspector prediction  
 (D) OTHER INFORMATION: name SRY\_02  
 score 0.951  
 sequence TAAACAAAACA

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
 (B) LOCATION: 486..493  
 (C) IDENTIFICATION METHOD: matinspector prediction  
 (D) OTHER INFORMATION: name E2F\_02  
 score 0.957  
 sequence TTTAGCGC

## (ix) FEATURE:

(A) NAME/KEY: TF binding-site  
 (B) LOCATION: complement(514..521)  
 (C) IDENTIFICATION METHOD: matinspector prediction  
 (D) OTHER INFORMATION: name MZF1\_01  
 score 0.975  
 sequence TGAGGGGA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

```

TGAGTGCAST GTTACATGTC AGTTGGGTTA AGTTTGTTAA TGTCATTCAA ATCTTCTATG   60
TCTTGATTGG CCTGCTAATT CTATTATTTC TGGAACATAA TTAGTTTGAT GGTTCATTATTA 120
GTTATTGACT GAGGTGTCCT AATCTCCCAT TATGTGGATT TATCTATTTT TTCAGTTGTA   180
GATAGGACAT TGATAGATAC ATAAGTACCA GGACAAAAGC AGGGAGATCT TTTTCCCAA   240
ATCAGGAGAA AAAAATGACA TCTGGAACAC CTATAGGGAA AGGCATAACA GATGGTAAGG   300
ATACTTTATC TTGAGTAGGA GAGCCTTCCT GTGGCAACST GGAGAGGCA AGAGGTCGTA   360
GAATTGAGGA GTCAGCTCAG TTAGAAGCAG GGAGTTGGGA ATTCCGTTCA TGTGATTAG   420
CATCASTGAT ATGGCAATG TGGGACTAAG GGTAGTGATC AGAGGGTTAA AATTGTGTGT   480
TTTGTTTATG CGCTGCTGGG GCATGCGCTT GGGTCCCTTC AACAGATTTC CCATGAATCT   540
CTTCAT                                     546

```

## (2) INFORMATION FOR SEQ ID NO: 32:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GTACCAGGGA CTGTGACCAT TGC

23

## (2) INFORMATION FOR SEQ ID NO: 33:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

CTGTGACCAT TGCTCCCAAG AGAG

24

## (2) INFORMATION FOR SEQ ID NO: 34:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 861 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: Genomic DNA

## (ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION: 1..806

## (ix) FEATURE:

- (A) NAME/KEY: transcription start site
- (B) LOCATION: 807

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(60..79)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name NFY\_Q6  
score 0.956  
sequence GGACCAATCAT

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 70..77
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MZF1\_01  
score 0.962  
sequence CCTGGGGA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 124..132
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name CMYB\_01  
score 0.994  
sequence TGACCGTTG

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(126..134)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name VMYB\_02  
score 0.985  
sequence TCCAACGGT

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 135..143
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name STAT\_01  
score 0.968  
sequence TTCCTGGAA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(135..143)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name STAT\_01  
score 0.951  
sequence TTCCAGGAA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: complement(252..259)
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name MZF1\_01  
score 0.956  
sequence TTGGGGGA

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 357..368
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name IK2\_01  
score 0.965  
sequence GAATGGGATTTC

## (ix) FEATURE:

- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 384..391
- (C) IDENTIFICATION METHOD: matinspector prediction

- (D) OTHER INFORMATION: name MZF1\_01  
score 0.986  
sequence AGAGGGGA
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(410..421)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name SRY\_02  
score 0.955  
sequence GAAACAAAACA
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: 592..599  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name MZF1\_01  
score 0.960  
sequence GAAGGGGA
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: 618..627  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name MYOD\_Q6  
score 0.981  
sequence AGCATCTGCC
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: 632..642  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name DELTAEF1\_01  
score 0.958  
sequence TCCCACCTTC
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(813..823)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name S8\_01  
score 0.992  
sequence GAGGCAATTAT
- (ix) FEATURE:  
(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(824..831)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name MZF1\_01  
score 0.986  
sequence AGAGGGGA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

TACTATAGGG CACGCGTGGT CGACGGCCGG GGTGTTCTGG AGCAGAGGGC ATGTCAGTAA 60  
TGATTGGTCC CTGGGGAAGG TCTGGCTGGC TCCAGTACAG TGAGGCATTT AGGTATCTCT 120  
CGGTGACCGT TGGATTCTCT GAAGCAGTAG CTGTTCTGTT TGGATCTGGT AGGGACAGGG 180

```

CTCAGAGGGG TAGGCACGAG GGAAGGTCAG AGGAGAAGGS AGGSARGGCC CAGTGAGARG 240
GGAGCATGCC TTCCCCAAC CCTGGCTTSC YCTTGGYMAN AGGGCGKTTY TGGMACTTA 300
AAYTCAGGGC CCAASCAGAA SCACAGGCCC AKTCNTGGCT SMAAGCACAA TAGCCTGAAT 360
GGGATTTTAC GTTAGNCAGG GTGAGAGGGG AGGCTCTCTG GCTTAGTTTT GTTTTGTTTT 420
CCAAATCAAG GTAACCTTGT CCCTTCTGCT ACGGGCCTTG GTCTTGGCTT GTCCTCACCC 480
AGTCGGAACCT CCCTACCACT TTCAGGAGAG TGGTTTTAGG CCCGTGGGGC TGTCTGTTC 540
CAAGCAGTGT GAGAACAIGG CTGGTAGAGG CTCTAGCTGT GTGCGGGGCC TGAAGGGGAG 600
TGGGTCTCTG CCCAAAGAGC ATCTGCCCCT TTCCACCTT CCCTTCTCCC ACCAGAAGCT 660
TGCCTGAGCT GTTTGGACAA AAATCCAAAC CCCACTTGGC TACTCTG3CC TGGCTTCAGC 720
TTGGAACCCA ATACCTAGGC TTACAGGCCA TCCTGAGCCA GGGGCCTCTG GAAATTCTCT 780
TCCTGATGGT CCTTTAGGTT TGGGCACAAA ATATAATTGC CTCTCCCCTC TCCCATTTTC 840
TCTCTTGGGA GCAATGGTCA C 861

```

## (2) INFORMATION FOR SEQ ID NO: 35:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: Other nucleic acid

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

```
CTGGGATGGA AGGCACGGTA 20
```

## (2) INFORMATION FOR SEQ ID NO: 36:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: SINGLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: Other nucleic acid

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

```
GAGACCACAC AGCTAGCAA 20
```

## (2) INFORMATION FOR SEQ ID NO: 37:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 555 base pairs  
    (B) TYPE: NUCLEIC ACID  
    (C) STRANDEDNESS: DOUBLE  
    (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: Genomic DNA
- (ix) FEATURE:  
    (A) NAME/KEY: promoter  
    (B) LOCATION: 1..500
- (ix) FEATURE:  
    (A) NAME/KEY: transcription start site  
    (B) LOCATION: 501
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: 191..206  
    (C) IDENTIFICATION METHOD: matinspector prediction  
    (D) OTHER INFORMATION: name ARNT\_01  
                            score 0.964  
                            sequence GGACTCACGTGCTGCT
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: 193..204  
    (C) IDENTIFICATION METHOD: matinspector prediction  
    (D) OTHER INFORMATION: name NMYC\_01  
                            score 0.965  
                            sequence ACTCACGTGCTG
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: 193..204  
    (C) IDENTIFICATION METHOD: matinspector prediction  
    (D) OTHER INFORMATION: name USF\_01  
                            score 0.985  
                            sequence ACTCACGTGCTG
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: complement(193..204)  
    (C) IDENTIFICATION METHOD: matinspector prediction  
    (D) OTHER INFORMATION: name USF\_01  
                            score 0.985  
                            sequence CAGCACGTGAGT
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: complement(193..204)  
    (C) IDENTIFICATION METHOD: matinspector prediction  
    (D) OTHER INFORMATION: name NMYC\_01  
                            score 0.956  
                            sequence CAGCACGTGAGT
- (ix) FEATURE:  
    (A) NAME/KEY: TF binding-site  
    (B) LOCATION: complement(193..204)  
    (C) IDENTIFICATION METHOD: matinspector prediction

.. (D) OTHER INFORMATION: name MYCMAX\_02  
score 0.972  
sequence CAGCACGTGAGT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 195..202  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name USF\_C  
score 0.997  
sequence TCACGTGC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(195..202)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name USF\_C  
score 0.991  
sequence GCACGTGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(210..217)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name MZF1\_01  
score 0.968  
sequence CATGGGGA

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 397..410  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name ELK1\_02  
score 0.963  
sequence CTCTCCGGAAGCCT

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: 400..409  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name CETS1P54\_01  
score 0.974  
sequence TCCGGAAGCC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(460..470)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name AP1\_Q4  
score 0.963  
sequence AGTGACTGAAC

(ix) FEATURE:

(A) NAME/KEY: TF binding-site  
(B) LOCATION: complement(460..470)  
(C) IDENTIFICATION METHOD: matinspector prediction  
(D) OTHER INFORMATION: name AP1FJ\_Q2  
score 0.961  
sequence AGTGACTGAAC

(ix) FEATURE:



- (A) NAME/KEY: TF binding-site
- (B) LOCATION: 547..555
- (C) IDENTIFICATION METHOD: matinspector prediction
- (D) OTHER INFORMATION: name PADS\_C  
score 1.000  
sequence TGTGGTCTC

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

```

CTATAGGGCA CGCKTGGTCG ACGGCCCGGG CTGGTCTGGT CTGKGTGGA CTCGGGTTGA    60
AGGACAGCAT TTGKACATC TGGTCTACTG CACCTTCCCT CTGCCGIGCA CTGGCCCTTT   120
KAWAAGCTCA GCACCGGTGC CCATCACAGG GCCGGCAGCA CACACATCCC ATTACTCAGA   180
AGGAAGTGAC GGAAGTACGT GCTGCTCCGT CCCCATGAGC TCAGTGGACC TGTCTATSTA   240
GAGCAGTCAG ACAGTGCCTG GGATAGAGTG AGAGTTCAGC CAGTAAATCC AAGTGATTGT   300
CATTCTGTGC TGCATTAGTA ACTCCCAACC TAGATGTGAA AACTTAGTTC TTTCTCATAG   360
GTTGCTCTGC CCATGGTCCC ACTGCAGACC CAGGCACTCT CCGGAAGCCT GGAAATCACC   420
CGTGTCTTCT GCCTGCTCCC GCTCACATCC CACACTTGTG TTCAGTCACT GAGTTACAGA   480
TTTTGCCTCC TCAATTTCTC TTGTCTTAGT CCCATCCTCT GTTCCCTGG CCAGTTTGTC   540
TAGCTGTGTG GTCTC                                         555

```

(2) INFORMATION FOR SEQ ID NO: 38:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 464 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 90..179
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 13.2  
seq LLLLSTLVIPSAA/AP

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

```

AAAACAGTAC GTGGGCGGCC GGAATCCGGG AGTCCGCTGA CCCGGGCTGT GGTCTAGCAT    60
AAAGGCGGAG CCAGAAGAAG GGGCGGGGT ATG GGA GAA GCC TCC CCA CCT GCC   120
Met Gly Glu Ala Ser Pro Pro Ala
-30                               -25

```

```

CCC GCA AGG CGG CAT CTG CTG GTC CTG CTG CTC CTC TCT ACC CTG      161
Pro Ala Arg Arg His Leu Leu Val Leu Leu Leu Leu Leu Ser Thr Leu
      -20                      -15                      -10

GTG ATC CCG TCC GCT GCA GGT CCT ATC CAT GAT GGT GAC GCC CAA GAG      209
Val Ile Pro Ser Ala Ala Ala Pro Ile His Asp Ala Asp Ala Gln Glu
      -5                      1                      5                      10

ASC TCC TTG GGT CTC ACA GGC CTC CAG AGC CTA CTC CAA GGC TTC AGC      257
Ser Ser Leu Gly Leu Thr Gly Leu Gln Ser Leu Leu Gln Gly Phe Ser
      15                      20                      25

CGA CTT TTC CTG AAA GGT AAC CTG CTT CGG GGC ATA GAC AGC TTA TTC      305
Arg Leu Phe Leu Lys Gly Asn Leu Leu Arg Gly Ile Asp Ser Leu Phe
      30                      35                      40

TCT GCC CCC ATG GAC TTC CGG GGC CTC CCT GGG AAC TAC CAC AAA GAG      353
Ser Ala Pro Met Asp Phe Arg Gly Leu Pro Gly Asn Tyr His Lys Glu
      45                      50                      55

GAG AAC CAG GAG CAC CAG CTG GGG AAC AAC ACC CTC TCC AGC MAC CTC      401
Glu Asn Gln Glu His Gln Leu Gly Asn Asn Thr Leu Ser Ser Xaa Leu
      60                      65                      70

CAG ATC GAC NNG ATG ACC GAC AAC AAG ACA GGA GAG GTG CTG ATC TCC      449
Gln Ile Asp Xaa Met Thr Asp Asn Lys Thr Gly Glu Val Leu Ile Ser
      75                      80                      85                      90

GAG AAT GTG GTG GCA
Glu Asn Val Val Ala
      95

```

## (2) INFORMATION FOR SEQ ID NO: 39:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 199 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 56..118
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 12  
seq VLVLCVLLLQAQG/GY

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

```

AAGGGAAGTC TGTGACTGCC TGGCCAGACT TAGGGGTCAC GGTCTGGTCA GAGTT ATG      58
Met

```

```

SCA CCC CAG ACT CTG CTG CCT GTC CTG GTT CTC TGT STG CTG CTG CTG      106
Ala Pro Gln Thr Leu Leu Pro Val Leu Val Leu Cys Val Leu Leu Leu
-20                      -15                      -10                      -5

CAG GCC CAG GGA GGA TAC CGT GAC AAG ATG AGG ATG CAG AGA ATC AAG      154
Gln Ala Gln Gly Gly Tyr Arg Asp Lys Met Arg Met Gln Arg Ile Lys
                      1                      5                      10

GTC TGT GAG AAG CGA CCC AGC ATA GAT CTA TGC ATC CAC CAC AGG      199
Val Cys Glu Lys Arg Pro Ser Ile Asp Leu Cys Ile His His Arg
                      15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 40:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 349 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 47..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 11  
seq SLVLLCLTCSYA/FM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

```

AAAGCAAACC CGTCATGAGC AACTCCCTTC CCCATCTCTG TTCACC ATG TGG ACG      55
                                   Met Trp Thr

CTG AAA TCG TCC CTG GTC CTG CTT CTG TGC CTC ACC TGC AGC TAT GCC      103
Leu Lys Ser Ser Leu Val Leu Leu Leu Cys Leu Thr Cys Ser Tyr Ala
-15                      -10                      -5

TTT ATG TTC TCT TCT CTG AGA CAG AAA ACT AGC GAA CCC CAG GGG AAG      151
Phe Met Phe Ser Ser Leu Arg Gln Lys Thr Ser Glu Pro Gln Gly Lys
 1                      5                      10                      15

GTG CAA TAC GGA GAG CAC TTT CGG ATT CGG CAG AAT CTA CCA GAG CAC      199
Val Gln Tyr Gly Glu His Phe Arg Ile Arg Gln Asn Leu Pro Glu His
                20                      25                      30

ACC CAA CCC TGG CTT GGG AGC AAA TGG CTC TGG CTT CTT KTT GTT GTT      247
Thr Gln Gly Trp Leu Gly Ser Lys Trp Leu Trp Leu Xaa Val Val
                35                      40                      45

GTG CCG TTT GTG ATA CTG CAG TGT CAA AGA GAC AGT GAG AAG AAT AAG      295
Val Pro Phe Val Ile Leu Gln Cys Gln Arg Asp Ser Glu Lys Asn Lys
                50                      55                      60

```

GAG CAG AAT CCT CCT GGC CTT CGA GGC GGC CAA CTT CAC TCT CCA TTA 343  
 Glu Gln Ser Pro Pro Gly Leu Arg Gly Gly Gln Leu His Ser Pro Leu  
 65 70 75 80

AAG AAA 349  
 Lys Lys

(2) INFORMATION FOR SEQ ID NO: 41:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 414 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 70..117
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.6  
seq LLLLPLLWGGSLQ/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

AAATTTTGGG GCATTTCCTT CCCTGACAGC CGGACCTGGG ACTGGGCTGG GGCCCTGGCG 50  
 GATGGAGAC ATG CTG CCC CTG CTG CTG CTG CCC CTG CTG TGG GGG GGG TCC 111  
 Met Leu Pro Leu Leu Leu Leu Pro Leu Leu Trp Gly Gly Ser  
 -15 -10 -5  
 CTG CAG GAG AAG CCA GTG TAC GAG CTG CAA GTG CAG AAG TCG GTG ACG 159  
 Leu Gln Glu Lys Pro Val Tyr Glu Leu Gln Val Gln Lys Ser Val Thr  
 1 5 10  
 GTG CAG GAG GGC CTG TGC GTC CTT GTG CCC TGC TCC TTC TCT TAC CCC 207  
 Val Gln Glu Gly Leu Cys Val Leu Val Pro Cys Ser Phe Ser Tyr Pro  
 15 20 25 30  
 TGG AGA TCC TGG TAT TCC TCT CCC CCA CTC TAC GTC TAC TGG TTC CGG 255  
 Trp Arg Ser Trp Tyr Ser Ser Pro Pro Leu Tyr Val Tyr Trp Phe Arg  
 35 40 45  
 GAC GGG GAG ATC CCA TAC TAC GCT GAG GTT GTG GCC ACA AAC AAC CCA 303  
 Asp Gly Glu Ile Pro Tyr Tyr Ala Glu Val Val Ala Thr Asn Asn Pro  
 50 55 60  
 GAC AGA AGA GTG AAG CCA GAG ACC CAG GGC CGA TTC CGC CTC CTT GGG 351  
 Asp Arg Arg Val Lys Pro Glu Thr Gln Gly Arg Phe Arg Leu Leu Gly  
 65 70 75  
 GAT GTC CAG AAG AAG AAC TGC TCC CTG AGC ATC GGA GAT SCC AGA ATG 399

Asp Val Gln Lys Lys Asn Cys Ser Leu Ser Ile Gly Asp Xaa Arg Met  
 80 85 90

GAG GAC ACG GGC GGG  
 Glu Asp Thr Gly Gly  
 95

414

## (2) INFORMATION FOR SEQ ID NO: 42:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 215 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 24..101
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.4  
 seq LLLLLCGPSQDQC/RP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

AANCCACCTG CSGCCGGCCA CCC ATG GAG ACT GGA GCG CTG CGG CGC CCG CAA 53  
 Met Glu Thr Gly Ala Leu Arg Arg Pro Gln  
 -25 -20

CTT CTC CCG TTG CTG CTG CTG CTC TGC GGC CCT TCC CAG GAT CAA TGC 101  
 Leu Leu Pro Leu Leu Leu Leu Cys Gly Pro Ser Gln Asp Gln Cys  
 -15 -10 -5

CGA CCT GTA CTC CAG AAT CTG TTG CAG AGC CCA GGC TTG ACA TGG AGC 149  
 Arg Pro Val Leu Gln Asn Leu Leu Gln Ser Pro Gly Leu Thr Trp Ser  
 1 5 10 15

TTG GAA GTG CCC ACT GGG AGA GAA GGA AAG GAA GGT ACT ATG AGA GTT 197  
 Leu Glu Val Pro Thr Gly Arg Glu Gly Lys Glu Gly Thr Met Arg Val  
 20 25 30

TCA CCA ACT GCA CCA AGG 215  
 Ser Pro Thr Ala Pro Arg  
 35

## (2) INFORMATION FOR SEQ ID NO: 43:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 297 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

- (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis
- (ix) FEATURE:
- (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 49..96
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 10.1  
seq LVLTLCPLAVA/SA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

```

AAAGATCCCT GCAGCCCGGC AGGAGAGAAG GCTGAGCCTT CTGGCGTC ATG GAG AGG      57
                                         Met Glu Arg
                                         -15

CTC GTC CTA ACC CTG TGC ACC CTC CCG CTG GCT GTG GCG TCT GCT GGC      105
Leu Val Leu Thr Leu Cys Thr Leu Pro Leu Ala Val Ala Ser Ala Gly
          -10                      -5                      1

TGC GCC ACG ACG CCA GCT CGC AAC CTG AGC TGC TAC CAG TGC TTC AAG      153
Cys Ala Thr Thr Pro Ala Arg Asn Leu Ser Cys Tyr Gln Cys Phe Lys
          5                      10                      15

GTC AGC AGC TGG ACG GAG TGC CCG CCC ACC TGG TGC AGC CCG CTG GAC      201
Val Ser Ser Trp Thr Glu Cys Pro Pro Thr Trp Cys Ser Pro Leu Asp
          20                      25                      30                      35

CAA GTC TGC ATC TCC AAC GAG GTG GTC GTC TCT TTT AAA TGG AGT GTA      249
Gln Val Cys Ile Ser Asn Glu Val Val Val Ser Phe Lys Trp Ser Val
          40                      45                      50

CGC GTC CTG CTC AGC AAA CGC TGT GGT CCC AGA TGT CCC AAC TCA GGG      297
Arg Val Leu Leu Ser Lys Arg Cys Ala Pro Arg Cys Pro Asn Ser Gly
          55                      60                      65

```

(2) INFORMATION FOR SEQ ID NO: 44:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 421 base pairs
  - (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: DOUBLE
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis
- (ix) FEATURE:
- (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 62..130

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 9.8

seq FLLFFFLFLLTRG/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

```

ACATGGTCGG YGTGCAGGAT ATTCGCTGG ACCCTAGAAA AGCCACCACG ACCTGTGGGC   60
C ATG ATG CTA CCC CAA TGG CTG CTG CTG CTG TTC CTT CTC TTC TTC TTT   109
Met Met Leu Pro Gln Trp Leu Leu Leu Leu Phe Leu Leu Phe Phe Phe
      -20                      -15                      -10

CTC TTC CTC CTC ACC AGG GGC TCA CTT TCT CCA ACA AAA TAC AAC CTT   157
Leu Phe Leu Leu Thr Arg Gly Ser Leu Ser Pro Thr Lys Tyr Asn Leu
      -5                      1                      5

TTG GAG CTC AAG GAG KSK KGC ATS GGG AAC CAG GAC TGC GAG ACT GGC   205
Leu Glu Leu Lys Glu Xaa Xaa Xaa Gly Asn Gln Asp Cys Glu Thr Gly
      10                      15                      20                      25

TGC TGC CAA CGT GCT CCA GAC AAT TGC GAG TCG CAC TGC GCG GAG AAG   253
Cys Cys Gln Arg Ala Pro Asp Asn Cys Glu Ser His Cys Ala Glu Lys
      30                      35                      40

GGG TCC GAG GGC AGT CTG TGT CAA ACG CAG GTG TTC TTT GGC CAA TAT   301
Gly Ser Glu Gly Ser Leu Cys Gln Thr Gln Val Phe Phe Gly Gln Tyr
      45                      50                      55

AGA GCG TGT CCC TGC CTG CGG AAC CTG ACT TGT ATA TAT TCA AAG AAT   349
Arg Ala Cys Pro Cys Leu Arg Asn Leu Thr Cys Ile Tyr Ser Lys Asn
      60                      65                      70

GAG AAA TGG CTT AGC ATC GCC TAT GCC CGT TGT CAG AAA ATT GGA AGG   397
Glu Lys Trp Leu Ser Ile Ala Tyr Gly Arg Cys Gln Lys Ile Gly Arg
      75                      80                      85

CAG AAG TTG GCT AGR AAA TGT TCT   421
Gln Lys Leu Ala Arg Lys Cys Ser
      90                      95

```

(2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 63..133

(C) IDENTIFICATION METHOD: Von Heijne matrix

-- (D) OTHER INFORMATION: score 9.8  
seq LVVFCLALQLVPS/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

```

AAACAGCAGT GCCTGGTCAA ACCCAGCAAC COTTGGCCAG AACTTACTCA CCCATCCCAC   60
TGACACC ATG AAG CCT GTG CTG CCT CTC CAG TWC CTG GTG GTG TTC TGC   109
    Met Lys Pro Val Leu Pro Leu Gln Xaa Leu Val Val Phe Cys
          -20                      -15                      -10

CTA GCA CTG CAG CTG GTG CCT GGG AGT CCC AAG CAG CTA GGG   151
Leu Ala Leu Gln Leu Val Pro Gly Ser Pro Lys Gln Leu Gly
          -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 253 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 134..238
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.5  
seq LFFSLFSAPLASA/VR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

```

AAACAGATAC TCCCAGCACA TGTTCCAWAG CAGCCCCCTG ATCCAATTTT CCTTAGCAGC   60
TAGGCTCAAG ACAATGCCCC ACTTCCCAAA GGCCTTGTGG CAATGTCCTC TTTTCTTTTC   120
ACATATATGA TTT ATG TTC CGT CAA CGA CAG GAA ACT GCT CAA AGA TCC   169
    Met Phe Arg Gln Arg Gln Glu Thr Ala Gln Arg Ser
          -35                      -30                      -25

ACC CAG TCC TGC CGC TGC CCC CGT GAT GGT TTG TTT TTC TCA TTG TTT   217
Thr Gln Ser Cys Arg Cys Pro Arg Asp Gly Leu Phe Phe Ser Leu Phe
          -20                      -15                      -10

AGC GCT CCA TTA GCT TCC GCA GTG AGA GCC GCC ASG   253
Ser Ala Pro Leu Ala Ser Ala Val Arg Ala Ala Xaa
          -5                      1                      5

```

2. INFORMATION FOR SEQ ID NO: 47:



## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 14..91
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.4  
seq RLLALPLALVLG/FE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

```
AGTACCACAG GCA ATG GGG TCA AGT GCC TGT GAA ATA GCT GTC GGG ACT    43
      Met Gly Ser Ser Ala Cys Glu Ile Ala Val Gly Thr
      -25                      -20                      -15

AAA AGG TTA TTA TTA GCT CTG CCT CTC GCT CTT GTT CTG GGC TTT GAA    97
Lys Arg Leu Leu Leu Ala Leu Pro Leu Ala Leu Val Leu Gly Phe Glu
      -10                      -5                      1

GGC TCA TCA GTT CCC CCA AGA AAT TTT    124
Gly Ser Ser Val Pro Pro Arg Asn Phe
      5                      10
```

## (2) INFORMATION FOR SEQ ID NO: 48:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 353 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 186..254
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.4  
seq SLLFICFFGESFC/IC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

```

AATATTTTGC TGACTGGCAA GGTTATATGA AGTGCCTTTA TTGAAGCACC ATTTCAACTA   60
ATAGCTCCTG GTATTTTCTG CTTCCCTTCG TAGGGAATTT AGTTATTITA TTTTATTATT  120
TAGCTAAATTT AGCTATTTTA AAATAGCTAA ATTTTAGCTA CTTTTTTTTC AATTGACAAA  180
GAAGG ATG TCT AAT CAA AGA CTA CCS CTG ATT TTT TCT CTG ITG TTT ATC   230
    Met Ser Asn Gln Arg Leu Pro Leu Ile Phe Ser Leu Leu Phe Ile
          -20          -15          -10

TGC TTC TTC GGG GAG AGT TTC TGC ATT TGT GAT GGA ACT GTC TGG ACA   273
Cys Phe Phe Gly Glu Ser Phe Cys Ile Cys Asp Gly Thr Val Trp Thr
          -5          1          5

WWG GTT KRA TGG GAG ATT CTT CCA GAA GAA STA CAT TAT TGG AAA GTT   326
Xaa Val Xaa Trp Glu Ile Leu Pro Glu Glu Val His Tyr Trp Lys Val
      10          15          20

AAG GGT TCT CCA TCT CAC TGC CTG CGG   353
Lys Gly Ser Pro Ser His Cys Leu Arg
    25          30

```

## (2) INFORMATION FOR SEQ ID NO: 49:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 167 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 108..155
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.2  
seq FLSFLLALLSLNC/IP

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

```

ACATCCAGTT TACAAAATTT ATATTATCAT GAAATACTTC AATAGAGGGT TGGGAAATCT   60
AATCTGGAG GAAATGCCAC AAATTTCCAC TGCTGGGGTT TTTGAAG ATG CTC TGG   116
                                Met Leu Trp
                                -15

TTC STA TCT TTT CTT CTA GCT CTC CTT TCC CTC AAT TGT ATC CCC ATC   164
Phe Leu Ser Phe Leu Leu Ala Leu Leu Ser Leu Asn Cys Ile Pro Ile
      -10          -5          1

GGG   167
Gly

```

## (2) INFORMATION FOR SEQ ID NO: 50:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 203 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 84..155
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.2  
seq ICCVIVLISLSWT/S?

## (x) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

```

ATGTCGAAT TTATTGCTGA GACTTTCCAG TGCATTTTGC ATCTCTCTGA GTGTGTCCTT   60
GATTTCCAA AGTTGTTTAA TTT ATG CTG TKT ATT TCA CTC GAG ATT KTT TCC   113
                Met Leu Xaa Ile Ser Leu Glu Ile Xaa Ser
                -20                               -15

TTC ATA TGC TGT GTC ATT GTT TTG ATT TCT TTA AGT TGG ACT TCA CCT   161
Phe Ile Cys Cys Val Ile Val Leu Ile Ser Leu Ser Trp Thr Ser Pro
                -10                               -5                               1

TTC ACT GGT GTG TAC TTG ATT GGT TTA ATA ATC GAG CCA GGG           203
Phe Thr Gly Val Tyr Leu Ile Gly Leu Ile Ile Glu Pro Gly
                5                               10                               15

```

## (2) INFORMATION FOR SEQ ID NO: 51:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 266 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 183..239
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.2

seq ILFILTFSSHTEC/SR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

```

AATTTCACTG ATGTCTAGCT GTGGCTCTCT TTTTATACCT CCTATTTAAT ACCACATGGT   60
CTTTGAAACC TGGAGACTTA CTGATTCTTT GAGCTCTAGT AAATGTTCTT TTCTCATTTA   120
ATTGATCATT TTCTCCCATTT TGTGTCTCTC TTACATCCCC AGGGCATTAC TATTTTGTAG   180
CT ATG GTA TTC AGG AAC TGC ATT TTA TTT ATT TTA ACT TTT TTT TCT   227
  Met Val Phe Arg Asn Cys Ile Leu Phe Ile Leu Thr Phe Phe Ser
                    -15                -10                -5

CAT ACT TTC TGT AGT AGG CAG AAT AAA GCC CAG CCC TGG   266
  His Thr Phe Cys Ser Arg Gln Asn Lys Ala Gln Pro Trp
                    1                5

```

(2) INFORMATION FOR SEQ ID NO: 52:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 7..45
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1  
seq MLAACPLSPGCQS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

```

GACGGC ATG CTG GCT GCG TGT CCC CTC TCA CCA CGT TGC CAA AGC GCT   49
  Met Leu Ala Ala Cys Pro Leu Ser Pro Gly Cys Gln Ser Ala
                    -10                -5                1

CCA TCA ACG TGG AAT CAT TTT CCT CCT GAA AGA ATA ACC ACT GGA GCC   96
  Pro Ser Thr Trp Asn His Phe Pro Pro Glu Arg Ile Thr Thr Gly Ala
                    5                10                15

GCC AGC CTT CTG AAA CCA GGG GGT GGC CTC TGG CCA CGC ACA GTC TCT   144
  Gly Ser Leu Leu Lys Pro Gly Gly Gly Leu Trp Pro Arg Thr Val Ser
                    20                25                30

CTG CCC TCC CCT GCG   159
  Leu Pro Ser Pro Ala
                    35

```

## (2) INFORMATION FOR SEQ ID NO: 53:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 43..99
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1  
seq FLTLITHCTVSWA/QS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

```

AAGCTGCGGG TAGAGAAGAC AGGACTCAGG ACAATCTCCA GC ATG GCC TGG TCC      54
                                   Met Ala Trp Ser

CCT CTC TTC CTC ACC CTC ATC ACT CAC TGT ACA GTG TCC TGG GCC CAG      102
Pro Leu Phe Leu Thr Leu Ile Thr His Cys Thr Val Ser Trp Ala Gln
-15                               -10                               -5                               1

TCT GTT CTG ACT CAG CCA CCC TCG GTG TCT GAA GCC CCC AGA CAG AGG      150
Ser Val Leu Thr Gln Pro Pro Ser Val Ser Glu Ala Pro Arg Gln Arg
                               5                               10                               15

GTC ACC ATC TCC TGT TTT GGA AGC AGC TCC AAT ATC GGA CGA AAT GCT      198
Val Thr Ile Ser Cys Phe Gly Ser Ser Ser Asn Ile Gly Arg Asn Ala
                20                               25                               30

GTA AAC TGG TAT CAG CAA CTC CCA GGA AGG TCT CCC AGA CTT CTC ATT      246
Val Asn Trp Tyr Gln Gln Leu Pro Gly Arg Ser Pro Arg Leu Leu Ile
                35                               40                               45

TTT TAT AAT AAT CTC CCG GCA TCG      270
Phe Tyr Asn Asn Leu Pro Ala Ser
50                               55

```

## (2) INFORMATION FOR SEQ ID NO: 54:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 111 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vii) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 49..102  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 9.1  
seq LVSLCSWSPPLTS/SP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

```

ACAGGCATAG ATCTAGCCCC ACCATCAAGA CAAACAACAT TTTCTATT ATG TTA AAA    57
                                     Met Leu Lys

AGT GTC CTT GTA AGC CTT TGC AGT TSG TCT CCT CCC CTG ACT TCC AGC    105
Ser Val Leu Val Ser Leu Cys Ser Trp Ser Pro Pro Leu Thr Ser Ser
-15          -10          -5          1

CCC AGG                                111
Pro Arg

```

## (2) INFORMATION FOR SEQ ID NO: 55:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 285 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 154..219  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 9  
seq FILAALSLSTTFS/LQ

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

```

AAKACCCTCC CTCCCGTTGC TCCAACTAA TACGSACTGA ACGGATCGCT GCGAGGGTGG    60

GAGAGAAAAT TAGGGGGAGA AAGGACAGAK AGAKCAACTA CCATCCATAG CCAGATAGAT    120

TATCTTACAC TGAACTGATC AAGTACTKTG AAA ATG ACT TCG AAA TTN ATC TTG    174
                                     Met Thr Ser Lys Xaa Ile Leu
                                     -20

ATG TCC TTC ATA CTT GCT GCA CTG AGT CTT TCA ACC ACC TTT TCT CTC    222
Val Ser Phe Ile Leu Ala Ala Leu Ser Leu Ser Thr Thr Phe Ser Leu
-15          -10          -5          1

```

```

CAA CCA TAC CAG CAR AAG GTT CTA CTA GTT TCT TTT GAT GGA TTC CGT   170
Gln Pro Tyr Gln Gln Lys Val Leu Leu Val Ser Phe Asp Gly Phe Arg
          5              10              15

TGG GAT TAC TTA TAT   235
Trp Asp Tyr Leu Tyr
          20

```

## (2) INFORMATION FOR SEQ ID NO: 56:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 123 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..120
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 44..79  
id AA280744  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 52..111
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.9  
seq LAVXLGLATAVSA/GP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

```

GAAAGTGAAA GGAGGAAGAG GAGGCTAAAT GGCTGAGGAG GTCGCAGCGC C ATG AAG   57
                                   Met Lys
                                   -20

TCC CTG TCT CTV MTC CTM GCT GTG GMT TTG GGC CTG GCG ACC GCC GTC   105
Ser Leu Ser Leu Xaa Leu Ala Val Xaa Leu Gly Leu Ala Thr Ala Val
          -15              -10              -5

TCA GCA GCA CCC GCG TGG   123
Ser Ala Gly Pro Ala Trp
          1

```

## (2) INFORMATION FOR SEQ ID NO: 57:

## (1) SEQUENCE CHARACTERISTICS:

- .. (A) LENGTH: 345 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 106..168
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.8  
seq LLWALLFMQSLWP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

```

AAAGATACTG ACTGAACATG GCTGGCGGAC TCAGGCTGGG GTCTGCAGTG CAGCATTAA 60
GGGCGGCTGA CATGAATATG GAGTAGTTTT CTCTAGCAAA GAGTA ATG TGG GCC ATG 117
                                     Met Trp Ala Met
                                     -20

GAG TCA GGC CAC CTC CTC TGG GCT CTG CTG TTC ATG CAG TCC TTG TGG 165
Glu Ser Gly His Leu Leu Trp Ala Leu Leu Phe Met Gln Ser Leu Trp
      -15                -10                -5

CCT CAA CTG ACT GAT GGA GCC ACT CGA GTC TAC TAC CTG GGC ATC CGG 213
Pro Gln Leu Thr Asp Gly Ala Thr Arg Val Tyr Tyr Leu Gly Ile Arg
      1                5                10                15

GAT GTG CAG TGG AAC TAT GCT CCC AAG GGA AGA AAT GTC ATC ACG AAC 261
Asp Val Gln Trp Asn Tyr Ala Pro Lys Gly Arg Asn Val Ile Thr Asn
                20                25                30

CAG CCT CTG GAC AGT GAC ATA GTG GCT TCC AGC TTC TTA AAG TCT GAC 309
Gln Pro Leu Asp Ser Asp Ile Val Ala Ser Ser Phe Leu Lys Ser Asp
      35                40                45

AAG AAC CGG ATA GGG GGA ACT ACA AGA AGA CCA TGG 345
Lys Asn Arg Ile Gly Gly Thr Thr Arg Arg Pro Trp
      50                55

```

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 246 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens



.. (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 130..159  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8.3  
 seq LLVMGSLPSASWS/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

```

ACGGTGCAGG GCAGAGAAGG AGCAGCCTTG GACTGGGGAT CCTGAGTAGT CCTGTCTGGG   60
AATGGAGGGC ACTGAATTGG CACCCTCCTT GGAGGCCAC ATG GCC CAA ACA TGG   114
                               Met Ala Gln Thr Trp
                               -20
GCA TTD CTG CTG GTG ATG GGA TCT CTC CCT TCT GCC AGC TGG TCT CTG   162
Ala Xaa Leu Leu Val Met Gly Ser Leu Pro Ser Ala Ser Trp Ser Leu
-15                -10                -5                1
CCC TGT TTG AGC TGG GAA AGT TTG CTG AAG GCT GCA GCC TGT TCT GAG   210
Pro Cys Leu Ser Trp Glu Ser Leu Leu Lys Ala Ala Ala Cys Ser Glu
                5                10                15
TTG GAT GGT AGA AAT GTA GGA AAT ACA CCA ACT CGG   246
Leu Asp Gly Arg Asn Val Gly Asn Thr Pro Thr Arg
                20                25

```

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 201 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 130..195  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8.7  
 seq LITLLYVWPVINA/CQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

```

ATGAATGAST GTTAATAGGC AATTTTAAAG GACAGAACCT CTGGGGAACC ATCCTGCAGT   60
TCTCCATGTG TACTTAAATT GATTTTGGAA ACCAGAAACA TATACTACTT CCTTAGAAGT  120
TCTACATTS ATG AAA TGT GGG TTT CTG GCT TAC TTG CTA ATC ACA CTC TTG  171

```

Met Lys Cys Gly Phe Leu Ala Tyr Leu Leu Ile Thr Leu Leu  
 -20 -15 -10

TAT GTT TGG CCA GTT ATT AAT GCT TGC CAG 201  
 Tyr Val Trp Pro Val Ile Asn Ala Cys Gln  
 -5 1

## (2) INFORMATION FOR SEQ ID NO: 60:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 128 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 21..95
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.5  
seq LKVLLLPLAPAAA/QD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

AGGGCGGATC TTCTCCGGCC ATG AGG AAG CCA GCC GCT GGC TTC CTT CCC TCA 53  
 Met Arg Lys Pro Ala Ala Gly Phe Leu Pro Ser  
 -25 -20 -15

CTC CTG AAG GTG CTG CTC CTG CCT CTG GCA CCT GCC GCA GCC CAG GAT 101  
 Leu Leu Lys Val Leu Leu Leu Pro Leu Ala Pro Ala Ala Ala Gln Asp  
 -10 -5 1

TCG ACT CAG GCC TCC ACT CCA GGC AGG 128  
 Ser Thr Gln Ala Ser Thr Pro Gly Arg  
 5 10

## (2) INFORMATION FOR SEQ ID NO: 61:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 313 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 152..202
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.4  
seq LLFLTSVVPFVLA/PR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

```

AAGAATCTTC CCAGTAGGCG GCGCGGGAGG GAAAAGAGGA TTGAGGGGCT AGGCGGGGCG    50
GATCCCGTCC TCCCCCGATG TGAGCAGTTT TCCGAAACCC CGTCAGGCGA AGGCTGCCCA    120
GAGAGGTGGA GTCGGTAGCG GGGCCGGGAA C ATG AGG CAG TCT CTC CTA TTC    172
                               Met Arg Gln Ser Leu Leu Phe
                               -15

CTG ACC AGC GTG GTT CCT TTC GTG CTG GCG CCG CGA CCT CCG GAT GAC    220
Leu Thr Ser Val Val Phe Val Leu Ala Pro Arg Pro Pro Asp Asp
-10                -5                1                5

CCG GGC TTC GGC CCC CAC CAG AGA CTC GAG AAG CTT GAT TCT TTG CTC    268
Pro Gly Phe Gly Pro His Gln Arg Leu Glu Lys Leu Asp Ser Leu Leu
          10                15                20

TCA GAC TAC GAT ATT CTC TCT TTA TCT AAT ATC CAG CAG CAG CSG    313
Ser Asp Tyr Asp Ile Leu Ser Leu Ser Asn Ile Gln Gln Gln Xaa
          25                30                35

```

## (2) INFORMATION FOR SEQ ID NO: 62:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 142 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 29..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.1  
seq SVLLGLLALMATA/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

```

AAGCTGAGGT GGCAGTGGTT CCACCAAC ATG GAG CTC TCG CAG ATG TCG GAG    52
                               Met Glu Leu Ser Gln Met Ser Glu
                               -25                -20

CTG ATG GGG CTG TCG GTG TTG CTT GGG CTG CTG GCC CTG ATG GGG ACG    100

```

Leu Met Gly Leu Ser Val Leu Leu Gly Leu Leu Ala Leu Met Ala Thr  
 -15 -10 -5

GCG GCG GTA GCG CGG GGG TGG CTG CGC GCG GGG GAG GTG AGG 142  
 Ala Ala Val Ala Arg Gly Trp Leu Arg Ala Gly Glu Val Arg  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO: 63:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 358 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 50..244
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8  
seq LTLIGCLVTGVES/KI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

AAGGAAGGA TTACTCGAGC CTTGTTAGAA TCAGACATGG CTCAGGGG ATG CAG GAC 58  
 Met Gln Asp  
 -65

GCT CCC CTG AGC TGC CTG TCA CCG ACT AAG TGG AGC AGT GTT TCT TCC 106  
 Ala Pro Leu Ser Cys Leu Ser Pro Thr Lys Trp Ser Ser Val Ser Ser  
 -60 -55 -50

GCA GAC TCA ACT GAG AAG TCA GCC TCT GCG GCA GGC ACC AGG AAT CTG 154  
 Ala Asp Ser Thr Glu Lys Ser Ala Ser Ala Ala Gly Thr Arg Asn Leu  
 -45 -40 -35

CCT TTT CAG TTC TGT CTC CGG CAG GCT TTG AGG ATG AAG GCT GCG GGC 202  
 Pro Phe Gln Phe Cys Leu Arg Gln Ala Leu Arg Met Lys Ala Ala Gly  
 -30 -25 -20 -15

ATT CTG ACC CTC ATT GGC TGC CTG GTC ACA GGC GTC GAG TCC AAA ATC 250  
 Ile Leu Thr Leu Ile Gly Cys Leu Val Thr Gly Val Glu Ser Lys Ile  
 -10 -5 1

TAC ACT CGT TGC AAA CTG GCA AAA ATA TTC TCG AGG GCT GGC CTG GAC 298  
 Tyr Thr Arg Cys Lys Leu Ala Lys Ile Phe Ser Arg Ala Gly Leu Asp  
 5 10 15

AAT CAG AGG GGC TTC AGC CTT GGA AAS TGG ATC TGC ATG GCC TAT TAT 346  
 Asn Xaa Arg Gly Phe Ser Leu Gly Xaa Trp Ile Cys Met Ala Tyr Tyr  
 20 25 30

GAG AGC GGC TGG  
Glu Ser Gly Trp  
35

353

## (2) INFORMATION FOR SEQ ID NO: 64:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 419 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 24..311
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.6  
seq ALCGLCLLCPRAA/RV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

ATTTCCTCTG GCACCCTGTA TTC ATG GCC TTG GCG TTC TGC CTC TGC ATG GCT	53
Met Ala Leu Ala Phe Cys Leu Cys Met Ala	
-95 -90	
GAA GCC ATC CTA CTC TTC TCA CCT GAA CAC TCC CTG TTC TTC TTC TGC	101
Glu Ala Ile Leu Leu Phe Ser Pro Glu His Ser Leu Phe Phe Phe Cys	
-85 -80 -75	
TCC CGA AAA GCA CGG ATC CGG CTC CAC TGG GCA GGG CAG ACC CTA GCC	149
Ser Arg Lys Ala Arg Ile Arg Leu His Trp Ala Gly Gln Thr Leu Ala	
-70 -65 -60 -55	
ATC CTC TGT GCA GCT CTG GGC CTG GGC TTC ATC ATC TCC AGC AGG ACC	197
Ile Leu Cys Ala Ala Leu Gly Leu Gly Phe Ile Ile Ser Ser Arg Thr	
-50 -45 -40	
CGC AGT GAG CTG CCT CAT CTG GTG TCC TGG CAC AGC TGG GTG GGA GCC	245
Arg Ser Glu Leu Pro His Leu Val Ser Trp His Ser Trp Val Gly Ala	
-35 -30 -25	
CTG ACA CTG CTG GCC ACT GCT GTC CAG GCA CTG TGT GGG CTC TGC CTC	293
Leu Thr Leu Leu Ala Thr Ala Val Gln Ala Leu Cys Gly Leu Cys Leu	
-20 -15 -10	
CTT TCT CCC CCC GCA CCC AGG GTC TCA AGG GTG GCT CGC CTC AAG CTC	341
Leu Cys Pro Arg Ala Ala Arg Val Ser Arg Val Ala Arg Leu Lys Leu	
-5 1 5 10	
TAC CAT CTG ACA TGT GGA CTG GTG GTC TAC CTG ATG GCT ACA GTA ACG	389
Tyr His Leu Thr Cys Gly Leu Val Val Tyr Leu Met Ala Thr Val Thr	
15 20 25	

GTG CTT CTG GGC ATG TAC TCA GTA TGG TTC  
 Val Leu Leu Gly Met Tyr Ser Val Trp Phe  
                   30                  35

419

## (2) INFORMATION FOR SEQ ID NO: 65:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 336 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 37..207
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq LLHRLASFHRVWS/FP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

AAGKCTGCCG GTGGGGACTC TTGCAGGGCC GTCCCC ATG TTR CGT TTT CCG ACC	54
Met Leu Arg Phe Pro Thr	
-55	
TGT TTC CCA TCC KTC CGG GTG RTG GGA GAK AAG CAG CTC CCG CAG GAG	102
Cys Phe Pro Ser Xaa Arg Val Xaa Gly Xaa Lys Gln Leu Pro Gln Glu	
-50 -45 -40	
ATT ATT TWC CTG GTC TGG TCG CCC AAK CGG GAT CKC ATT GST TTG GCC	150
Ile Ile Xaa Leu Val Trp Ser Pro Xaa Arg Asp Xaa Ile Xaa Leu Ala	
-35 -30 -25 -20	
AAC ACA GCT GGC GAG GTT TTA CTT CAT CGA CTG GCA AGT TTT CAT CGA	198
Asn Thr Ala Gly Glu Val Leu Leu His Arg Leu Ala Ser Phe His Arg	
-15 -10 -5	
GTT TGG AGT TTT CCA CCA AAT GAA AAT ACA GGA AWK GAG GTG ACG TGT	246
Val Trp Ser Phe Pro Pro Asn Glu Asn Thr Gly Xaa Glu Val Thr Cys	
1 5 10	
CTG GCA TGG AGA CCA GAT CCC AAA CTT TTG GCC TTT GCT CTT GCT GAT	294
Leu Ala Trp Arg Pro Asp Gly Lys Leu Leu Ala Phe Ala Leu Ala Asp	
15 20 25	
ACC AAG AAA ATT GTT TTG TGT GAT GTA GAA AAA CCT GAG AGC	336
Thr Lys Lys Ile Val Leu Cys Asp Val Glu Lys Pro Glu Ser	
30 35 40	

## (2) INFORMATION FOR SEQ ID NO: 66:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 398 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 9..134
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq LALVVALVAERFA/RR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

AGACCTTC ATG TTC ATG GTG CTG GAG GTG GTG GTG AGC CGG GTG ACC TCG	50
Met Phe Met Val Leu Glu Val Val Val Ser Arg Val Thr Ser	
-40 -35 -30	
TCG CTG GCG ATG CTC TCC GAC TCC TTC CAC ATG CTG TCG GAC GTG CTG	98
Ser Leu Ala Met Leu Ser Asp Ser Phe His Met Leu Ser Asp Val Leu	
-25 -20 -15	
GCG CTG GTG GTG GCG CTG GTG GCC GAG CGC TTC GCC CGG CGG ACC CAC	146
Ala Leu Val Val Ala Leu Val Ala Glu Arg Phe Ala Arg Arg Thr His	
-10 -5 1	
GCC ACC CAG AAG AAC ACG TTC GGC TGG ATC CGA GCC GAG GTA ATG GGG	194
Ala Thr Gln Lys Asn Thr Phe Gly Trp Ile Arg Ala Glu Val Met Gly	
5 10 15 20	
GCT CTG GTG AAC GCC ATC TTC CTG ACT GGC CTC TGT TTC GCG ATC CTG	242
Ala Leu Val Asn Ala Ile Phe Leu Thr Gly Leu Cys Phe Ala Ile Leu	
25 30 35	
CTG GAG GCC ATC GAG CGC TTC ATC GAG CCG CAC GAG ATG CAG CAG CCG	290
Leu Glu Ala Ile Glu Arg Phe Ile Glu Pro His Glu Met Gln Gln Pro	
40 45 50	
CTG GTG GTS CNT TGG GGT CGG GCG TGG CSG GCT GCT GGT CAA CGT GCT	338
Leu Val Val Xaa Trp Gly Arg Ala Trp Xaa Ala Ala Gly Gln Arg Ala	
55 60 65	
GGG GCT CTG CCT CTT CCA CCA TCA CAG CGG CTT CAG CCA GGA CTC CGG	386
Gly Ala Leu Pro Leu Pro Pro Ser Gln Arg Leu Gln Pro Gly Leu Arg	
70 75 80	
CCG CGG CCA TGG	393
Pro Arg Pro Trp	
85	

## (2) INFORMATION FOR SEQ ID NO: 67:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 306 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 70..186
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq LLLLLGLIVLVNI/GI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

```

AASGTGSST TGGGGCCGGG GGTGGGGGGC AGAGGGGGGT GGCCAGGTG GCCCTAGGAC   60
CCCCCTCC ATG GAA AAC CAG CTA TGG CAT AAC ACC GTG AGA TGT TGC AAT   111
      Met Glu Asn Gln Leu Trp His Asn Thr Val Arg Cys Cys Asn
                -35                      -30

CAA TAC CAA GAA AGC CCC CAC GAT GCC GAG GAC ATC TTA CTC CTG CTG   159
Gln Tyr Gln Glu Ser Pro His Asp Ala Glu Asp Ile Leu Leu Leu Leu
-25                -20                      -15                      -10

CTG GGC CTC ATC GTT CTT GTC AAC ATT GGC ATC AAC GTG GCA ACT ATG   207
Leu Gly Leu Ile Val Leu Val Asn Ile Gly Ile Asn Val Ala Thr Met
                -5                      1                      5

ATG TGG CAT GGA CTC CAG AAC GCC TTA GAC AAG ATG ATT GAT TGG GCT   255
Met Trp His Gly Leu Gln Asn Ala Leu Asp Lys Met Ile Asp Trp Ala
                10                      15                      20

ACT CAG AAA ATA GCA GTC TTC TTC GCT GTG TTC GTC GCC GCC GCC GCC   303
Thr Gln Lys Ile Ala Val Phe Phe Ala Val Phe Val Ala Ala Ala Ala
                25                      30                      35

CGG
Arg
40

```

## (2) INFORMATION FOR SEQ ID NO: 68:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 178 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR



(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 23..76

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 7.5  
seq ITLLTLSPNSVCC/CP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

```

AATACTGAGG TATATTGCCA AA ATG CTC TCC AKW AAG ATC ACC CTC TTG ACA      52
           Met Leu Ser Xaa Lys Ile Thr Leu Leu Thr
                               -15                -10

CTG TCA CCC AAT AGT GTG TGT TGC TGC CCC TCA GCA ACC CTG GGT GCC      100
Leu Ser Pro Asn Ser Val Cys Cys Cys Pro Ser Ala Thr Leu Gly Ala
           -5                      1                      5

AGC AAT CAT TCT CAT CTT TGG AGA TCT ACT AGC AGA CAT GGC ATC TCC      148
Ser Asn His Ser His Leu Trp Arg Ser Thr Ser Arg His Gly Ile Ser
          10                      15                      20

TTC CCA TGG GCA TTC CTT TTA ATT AAC GGG      178
Phe Pro Trp Ala Phe Leu Leu Ile Asn Gly
          25                      30

```

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 234 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 79..132

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 7.4  
seq GWLVLCVLAISLA/SM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

```

CTGCTGGGCA GCCACAGGG TCCCTGGGCG GAGGSCAGGA GCATCCMSTT GGAGTTGACA      60
ACAGGAGGCA GAGGCATC ATG GAG GGT CCC CGG GGA TGG CTG GTG CTC TGT      111

```

```

--          Met Glu Gly Pro Arg Gly Trp Leu Val Leu Cys
              -15                      -10

GTG CTG GCC ATA TCG CTG GCC TCT ATG GTG ACC GAG GAC TTG TGC CGA   159
Val Leu Ala Ile Ser Leu Ala Ser Met Val Thr Glu Asp Leu Cys Arg
      -5                      1                      5

GCA CCA GAC GGG AAG AAA GGG GAG GCA GGV AVA CCT GGC AGA CGG GGG   207
Ala Pro Asp Gly Lys Lys Gly Glu Ala Gly Xaa Pro Gly Arg Arg Gly
    10                      15                      20                      25

CGG CCA GGC CTC AAG GGG GAG CAA CGG                               254
Arg Pro Gly Leu Lys Gly Glu Gln Arg
              30

```

## (2) INFORMATION FOR SEQ ID NO: 70:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 364 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 41..100
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.3  
seq LAVFMLLAQLVSG/NW

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

```

AATAGAGACT TCTGGACTCT ATAGAACCCA CTGCTCTCTG ATG AAG TCC CTA CTG   55
              Met Lys Ser Leu Leu
              -20

TTC ACC CTT GCA GTT TTT ATG CTC CTG GCC CAA TTG GTC TCA GGT AAT   103
Phe Thr Leu Ala Val Phe Met Leu Leu Ala Gln Leu Val Ser Gly Asn
-15                      -10                      -5                      1

TGG TAT GTG AAA AAG TGT CTA AAC GAC GTT GGA ATT TGC AAG AAG AAG   151
Trp Tyr Val Lys Lys Cys Leu Asn Asp Val Gly Ile Cys Lys Lys Lys
              5                      10                      15

TGC AAA CCT GAA GAG ATG CAT GTA AAG AAT GGT TGG GCA ATG TGC GGC   199
Cys Lys Pro Glu Glu Met His Val Lys Asn Gly Trp Ala Met Cys Gly
    20                      25                      30

AAA CAA AGG GAC TGC TGT GTT CCA GGT GAC ACA CGT SCT AAT TAT CCT   247
Lys Gln Arg Asp Cys Cys Val Pro Ala Asp Arg Arg Ala Asn Tyr Pro
    35                      40                      45

```

```

GTT TTC TGT GTC CAG ACA AAG ACT ACA AGA ATT TCA ACA GTC ACA GCA      295
Val Phe Cys Val Gln Thr Lys Thr Thr Arg Ile Ser Thr Val Thr Ala
 50                      55                      60                      65

ACA ACA GCA ACA ACA ACT TTG ATG ATG ACT ACT GCT TCG ATG TCT TCG      343
Thr Thr Ala Thr Thr Thr Leu Met Met Thr Thr Ala Ser Met Ser Ser
                      70                      75                      80

ATG GCT CCT ACC CGT TTC TCC
Met Ala Pro Thr Arg Phe Ser
                      85

```

## (2) INFORMATION FOR SEQ ID NO: 71:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 62 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 9..56
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.3  
seq LILLFSLISIVC/MI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

```

ATAGTAAA ATG TTA AAG TTG ATC TTA CTT TTT TCG CTC CTC ATC TCT ATT      50
Met Leu Lys Leu Ile Leu Leu Phe Ser Leu Leu Ile Ser Ile
  -15                      -10                      -5

GTT TGT ATG ATT
Val Cys Met Ile
      1

```

## (2) INFORMATION FOR SEQ ID NO: 72:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 296 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 195..272
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1  
seq LASLQWSLTLANC/GS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

```

AAAGTGTAGA ACACGGACCT CTGAGTTATG CTCTTGAGAG GTGCCAAAGC TGGGCTGTTT   60
ACCTACCTTA TCCACAGAGC TCTGAAAGTC AAGCCAGAAA GGAAGGATTC CAAATTCTTG   120
GAATTTTATC TAGAAAAGAA GACTAAGCAG CTTTGTGTCT TCTGTGACCC AGTTGCTGGC   180
CCAAGACATG GACA ATG ACC CCC TGG TGT TTG GCG TGT CTG GGG AGG AGG   230
              Met Thr Pro Trp Cys Leu Ala Cys Leu Gly Arg Arg
              -25                      -20                      -15
CCT CTC GCT TCT TTG CAG TGG AGC CTG ACA CTG GCG TGG TGT GGC TCC   278
Pro Leu Ala Ser Leu Gln Trp Ser Leu Thr Leu Ala Trp Cys Gly Ser
              -10                      -5                      1
GGC AGC CAC TGG ACA GAG                                           296
Gly Ser His Trp Thr Glu
              5

```

## (2) INFORMATION FOR SEQ ID NO: 73:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 315 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 151..228
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9  
seq LWVLLLCARHVVT/LV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

```

AACACGCAGC TAGACACAGC TAMCTTGAGT CTTGGAGCTC CTAGAGGGAM GCTTCTGGAM   60
AGGAAGGCTC TTCAGGACCT CTTAGGAGCC AGAGMMSMGG ACGTKSACAC AGATAAAGAG   120
CCAGGCTCAC CAGCTCCTGA CGCATGCAKS ATG ACC ATG ASA CAC AAC TGG ACA   174
              Met Thr Met Arg His Asn Trp Thr

```

(A) LENGTH: 224 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 114..191  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.9  
seq LWVLLLCARVVTL/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

```

ACTTCCCAGA AGCAGCTCTG GTGCTGAAGA GAGCACTGCC TCCCTGTGTG ACTGGAGAAG   60
AGGACGTTGT CACAGATAAA GAGCCAGGCT CACCAGCTCC TGACGCATGC ATC ATG   116
                                         Met
ACC ATG AGA CAC AAC TGG ACA CCA GAC CTC AGC CCT TTG TGG GTC CTC   164
Thr Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val Leu
-25                      -20                      -15                      -10
CTC CTG TGT GCC CAC GTC GTC ACT CTC CTG GTC AGA GCC ACA CCT GTC   212
Leu Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro Val
                      -5                      1                      5
TCG CAG CCC ACG                                         224
Ser Gln Pro Thr
                      10

```

(2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 333 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 79..138  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.9  
seq LYLLGMLVPGGLG/YD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

```

ACTTGATGCA TGCATTCTCA AGGACACTTG ATCCACTCCC AGAGAGGCCC AGAATTTTCT   50

```

```

AACTTACTGT GTGGCAGA ATG AAG CCT CTG CTT GAA ACC CTT TAT CTT TTG    111
      Met Lys Pro Leu Leu Glu Thr Leu Tyr Leu Leu
      -20                      -15                      -10

GGG ATG CTG GTT CCT GGA GGG CTG GGA TAT GAT AGA TCC TTA GCC CAA    159
Gly Met Leu Val Pro Gly Gly Leu Gly Tyr Asp Arg Ser Leu Ala Gln
      -5                      1                      5

CAC AGA CAA GAG ATT GTG GAC AAG TCA GTG AGT CCA TGG AGC CTG GAG    207
His Arg Gln Glu Ile Val Asp Lys Ser Val Ser Pro Trp Ser Leu Glu
      10                      15                      20

ACG TAT TCC TAT AAC ATA TAC CAC CCC ATG GGA GAG ATC TAT GAG TGG    255
Thr Tyr Ser Tyr Asn Ile Tyr His Pro Met Gly Glu Ile Tyr Glu Trp
      25                      30                      35

ATG AGA GAG ATC AGT GAG AAG TAC AAG GAA GTG GTG ACA CAG CAT TTC    303
Met Arg Glu Ile Ser Glu Lys Tyr Lys Glu Val Val Thr Gln His Phe
      40                      45                      50                      55

CTA GGA GTG ACC TAT GAG ACC CAA CCC GCG    333
Leu Gly Val Thr Tyr Glu Thr Gln Pro Ala
      60                      65

```

## (2) INFORMATION FOR SEQ ID NO: 77:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 295 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 80..274
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq LLFLISLAHLSQ/WT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

```

AAACTATTGG GGATGCTGAG CTGCGGGGTA CGGGCCTGAG GASGGATGGG AGTAAGAAGT    60

GCTGTGGAAA CCGTCAGCC ATG AAC CAG GCT GAC CCT CGG CTC AGA GCA GTG    112
      Met Asn Gln Ala Asp Pro Arg Leu Arg Ala Val
      -65                      -60                      -55

TGC TTG TGG AOT CTC ACA TCT GCA GCC ATG AGC AGA GGC GAC AAC TGC    160
Cys Leu Trp Thr Leu Thr Ser Ala Ala Met Ser Arg Gly Asp Asn Cys
      -50                      -45                      -40

```

ACG	GAT	CTA	CTC	GCA	CTG	GGA	ATC	CCC	TCC	ATA	ACC	CAG	GCC	TGG	GGA	208
Thr	Asp	Leu	Leu	Ala	Leu	Gly	Ile	Pro	Ser	Ile	Thr	Gln	Ala	Trp	Gly	
			-35					-30					-25			
CTG	TGG	GTC	CTC	TTA	GGG	GCT	GTG	ACG	CTG	CTA	TTT	CTC	ATC	TGG	CTG	256
Leu	Trp	Val	Leu	Leu	Gly	Ala	Val	Thr	Leu	Leu	Phe	Leu	Ile	Ser	Leu	
		-20					-15					-10				
GCT	GCA	CAC	TTG	TCC	CAG	TGG	ACC	AGG	GGT	CGG	AGC	GGG				295
Ala	Ala	His	Leu	Ser	Gln	Trp	Thr	Arg	Gly	Arg	Ser	Gly				
	-5					1				5						

(2) INFORMATION FOR SEQ ID NO: 78:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 451 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 317..442  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.8  
seq LLSILSSSLTMVIC/RH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

ACTACACAGA	GAGAAGCCAT	CATTCTAGCT	AGACAAGAAG	CTCGGGAAGA	ATTACTTTTA	60
CATCAGAGTG	AATGGGAGGG	AAGAATATCT	CCCGAGCAGG	TTGACACCTC	TTCCTTACCC	120
CTAGTACCAC	AGCATTCATT	CGCCTCATTA	CCTCTTAATG	AATCTGAAAG	AAACCAAGAA	180
CCATGTTCAA	TTAACAGTGA	TAATATAGTA	TCCTCAGGTC	ACTCAGAGAT	ACCAACATTG	240
CCTGATGGGC	TGTTGGGTTT	ATCACATCTT	GTTTTACCTC	AACAAGATAA	TTTGATTGCA	300
CTTGAAGAAC	ACTTGC	ATG CAC AGA CAG ATT TCC TTC CTT CTA TTG AGA AAA				352
		Met His Arg Gln Ile Ser Phe Leu Leu Leu Arg Lys				
		-40		-35		
CCC AGA AAG AAT TGG TTT TGT CAA AAC CAT GTA AAT TTG AGG AAA AGG						400
Pro Arg Lys Asn Trp Phe Cys Gln Asn His Val Asn Leu Arg Lys Arg						
-30		-25		-20	-15	
TAT CTT CTG AGC ATT TTA TCC AGT CTC ACC ATG GTG ATT TGC AGA CAC						448
Pyr Leu Leu Ser Ile Leu Ser Ser Leu Thr Met Val Ile Cys Arg His						
	-10		-5		1	
GGG						452



Gly

## (2) INFORMATION FOR SEQ ID NO: 79:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 317 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 162..290
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq ALSAXTFVSFLHA/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

```

AGTACGGATC TCTTTAATAT TCTGTGTAAC AAAATAGAAA TGCTCATAAA GTACTTCTGC    60
GGCAAACCAA AGTATAGCAC CTGACTCAAG GAAAAGCAAG GAAAAGCACA TGTGGGATCC    120
CTTGAATGGC AAGTGAAACT AGCCACTAGT TTCATTTTTC C ATG AAA CAA TGG CTG    176
                               Met Lys Gln Trp Leu
                               -40

TGT TGG GTG CTG AGG CTG GAA GGT AGA CAG GGG CTT GGG GTT GGA GAG    224
Cys Trp Val Leu Arg Leu Glu Gly Arg Gln Gly Leu Gly Val Gly Glu
   -35                      -30                      -25

CCT CGT GGG CTG CGT TTG TGC TTG GGG GCC TTG AGC GCA SCC ACC TTT    272
Pro Arg Gly Leu Arg Leu Cys Leu Gly Ala Leu Ser Ala Xaa Thr Phe
   -20                      -15                      -10

GTC AGC TTT CTA CAC GCT GCT CCC CAC TCC CAT CCA GCC CTT GGG    317
Val Ser Phe Leu His Ala Ala Pro His Ser His Pro Ala Leu Gly
   -5                      1                      5

```

## (2) INFORMATION FOR SEQ ID NO: 80:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 235 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 29..226  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.8  
seq LLFFLFPIIFRS/QH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

```

ACAATTGGG TGTGTCTGGT GTYTTGCC ATG AGA CTG GGG TTA TGC TTC TGG      52
                        Met Arg Leu Gly Leu Cys Phe Trp
                        -65                      -60

GTG CCA CAC AGA GGT GAA ATG TCC TTC TCA TCA CAT TAT TCG AGA GGT      100
Val Pro His Arg Gly Glu Met Ser Phe Ser Ser His Tyr Ser Arg Gly
      -55                      -50                      -45

ACA TGG TAC CAA TGG GAC TTA TCG CTG CTG ATG TTA ACC TTG ATC TCT      148
Thr Trp Tyr Gln Trp Asp Leu Ser Leu Leu Met Leu Thr Leu Ile Ser
      -40                      -35                      -30

TGG TTC AGG TGG TGC CTG CCA GCT GTC TCC ACT GTG GAG TTA CTA TTT      196
Trp Phe Arg Trp Cys Leu Pro Ala Val Ser Thr Val Glu Leu Leu Phe
      -25                      -20                      -15

TTC CTT TTC CCC ATT TTA TTC ATC AGA AGC CAG CAC CGG      235
Phe Leu Phe Pro Ile Leu Phe Ile Arg Ser Gln His Arg
      -10                      -5                      1

```

## (2) INFORMATION FOR SEQ ID NO: 81:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 390 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 67..369  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.6  
seq IILVITITSACSA/CI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

```

ACTTCTAAE AATACTTGTT TAGATATTTT GTGTCCCTAG ATTTGTCTGA TGATTAGGCT      60

```

GCASTT ATG GAT TTT TGG GAA GAA TAC CGC AGA GGT GAT GTG CCC TTC 103  
 Met Asp Phe Trp Glu Glu Tyr Arg Arg Gly Asp Val Pro Phe  
 -100 -95 -90

TCA TGG TGT CCT ATC AGG AGC TAC CTG ATG TCA GTA TGT CCT GTT ACT 156  
 Ser Trp Cys Pro Ile Arg Ser Tyr Leu Met Ser Val Cys Pro Val Thr  
 -35 -30 -75

GGC AAA GTT AAC CTT AAT CAT TTG GTT AAG GTA GCC TCT GCC AGG TTT 204  
 Gly Lys Val Asn Leu Asn His Leu Val Lys Val Ala Ser Ala Arg Phe  
 -70 -65 -60

CTC CAC CAA GTT ACT ATT TTT CCT TTT CTG TAC TCT GTT AAG GCA AAT 252  
 Leu His Gln Val Thr Ile Phe Pro Phe Leu Tyr Ser Val Lys Ala Asn  
 -55 -50 -45 -40

TAT TGC TTT TTA AAT TTT GAT GTA CCT CAA TAT GCA TGG GAG ATA CAT 300  
 Tyr Cys Phe Leu Asn Phe Asp Val Pro Gln Tyr Ala Trp Glu Ile His  
 -35 -30 -25

AGC TTT GCA GCT CCC TCA ATC TTA ATT GTA ATA ATA ATA GTA ATA ACA 343  
 Ser Phe Ala Ala Pro Ser Ile Leu Ile Val Ile Ile Ile Val Ile Thr  
 -20 -15 -10

ATA ACT AGC GCT TGC TCC GCC TGC ATA GTT CTA AAC ACA TGT 390  
 Ile Thr Ser Ala Cys Ser Ala Cys Ile Val Leu Asn Thr Cys  
 -5 1 5

## (2) INFORMATION FOR SEQ ID NO: 82:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 349 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 59..139
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5  
seq SLSLSTVWNWIQA/SF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

AATATTTAGA TTCCTGAAGC TTCTGCACAT GTAGTTCCTA GAGCTGCTGC TTATTAAA 35

ATG TCA ACA TCT TCA TCT TCT AGC TGG GAC AAC CTC TTA GAG TCT CTC 105  
 Met Ser Thr Ser Ser Ser Ser Ser Trp Asp Asn Leu Leu Glu Ser Leu  
 -25 -20 -15

TCT CTC AGC ACA GTA TGG AAT TGG ATA CAA GCA ACT TTT TTG GGA GAG 155

```

Ser Leu Ser Thr Val Trp Asn Trp Ile Gln Ala Ser Phe Leu Gly Glu
-10          -5          1          5

ACT AGT GCA CCT CAG CAA ACA AGT TTG GGA CTA TTA GAT AAT CTT GCT   202
Thr Ser Ala Pro Gln Gln Thr Ser Leu Gly Leu Leu Asp Asn Leu Ala
          10          15          20

CCA GCT GTG CAA ATC ATC TTG AGG ATT TCT TTC TTG ATT TTA TTG GGA   250
Pro Ala Val Gln Ile Ile Leu Arg Ile Ser Phe Leu Ile Leu Leu Gly
          25          30          35

ATA GGA ATA TAT GCC TTA TGG AAA CGA AGT ATT CAG TCA ATT CAG AAA   298
Ile Gly Ile Tyr Ala Leu Trp Lys Arg Ser Ile Gln Ser Ile Gln Lys
          40          45          50

ACA TTG TTG TTT GTA ATC ACA CTC TAC AAA CTT TAC AAG AAG GGC TCG   346
Thr Leu Leu Phe Val Ile Thr Leu Tyr Lys Leu Tyr Lys Lys Gly Ser
          55          60          65

GCG
Ala
70
349

```

## (2) INFORMATION FOR SEQ ID NO: 83:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 302 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 27..104
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5  
seq LALGSAGLLWCLA/GF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

```

AGCAGACCGG CCGCCGCTTC ACCGGC ATG GTC TTC GCC ACC ATC GGT TTC TCG   53
          Met Val Phe Ala Thr Ile Gly Phe Ser
          -25          -20

CTG AAG TCG GGC CTG GCC CTT GGC TCG GCG GGC CTG CTG TGG TGC CTG   101
Leu Lys Ser Gly Leu Ala Leu Gly Ser Ala Gly Leu Leu Trp Cys Leu
          -15          -10          -5

GGC GGT TTC TTC GGC TAC GAC ACA CAG CAG CCC ACG GCA CCC AAC GCC   149
Ala Gly Phe Phe Gly Tyr Asp Thr Gln Gln Pro Thr Ala Pro Asn Ala
          1          5          10          15

```

```

ATC GAG GGC TAC CGC GTC ATG TCC AGC TTC GGC GTC GGC GGC CTG TTC   197
Ile Glu Gly Tyr Arg Val Met Ser Ser Phe Gly Val Gly Ala Leu Phe
      20                      25                      30

GCC GCC TGC ACG ATC TGC CTG CTG GCS RAC AAG CTG AAC AAG CAG ACG   245
Ala Ala Cys Thr Ile Cys Leu Leu Ala Xaa Lys Leu Asn Lys Gln Thr
      35                      40                      45

ACG CTG AAG ATG GCC GAC GAC CTC GCC CAA CGG CGC CAG CAG GCC GAC   293
Thr Leu Lys Met Ala Asp Asp Leu Ala Gln Arg Arg Gln Gln Ala Asp
      50                      55                      60

CTT GCG CCG
Leu Ala Pro
      65

```

## (2) INFORMATION FOR SEQ ID NO: 84:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 151 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 35..76
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq VLLSGSVSVGVC/CA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

```

ACAATCTCAC AGTCCGTGGC AGAGCCTTGT CCTG ATG GTT TTA TTG CTT TCT GGC   55
      Met Val Leu Leu Leu Ser Gly
                        -10

AGT GTG AGT GTG GGT GTG TGT TGT GCC TAC TTG TGC ATC TCC ATT TCT   103
Ser Val Ser Val Gly Val Cys Cys Ala Tyr Leu Cys Ile Ser Ile Ser
      -5                      1                      5

AAA ACA CCA ACT GCT TGT GCA TTG TAT GGT CTT TAT TTA CCG TTT TTT   151
Lys Thr Pro Thr Ala Cys Ala Leu Tyr Gly Leu Tyr Leu Pro Phe Phe
      10                      15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 85:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 169 base pairs
- (B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 26..112  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.4  
seq GLCXLCXVXNVFA/GS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

ATAATCTGTA ACTTTAGCCC CAACC ATG TGC TCG CAG AAA CGT GCT GTA TCA	52
Met Cys Ser Gln Lys Arg Ala Val Ser	
-25	
AAT CAA GGT TTA ATG GAT TTA GGG CTG TGC ARG CTG TGC YTT GTT AMC	100
Asn Gln Gly Leu Met Asp Leu Gly Leu Cys Xaa Leu Cys Xaa Val Xaa	
-20 -15 -10 -5	
AAT GTG TTT GCA GGC AGT ATG CCT GGT AAA AGT CAT TGC CAT TCT CCA	148
Asn Val Phe Ala Gly Ser Met Pro Gly Lys Ser His Cys His Ser Pro	
1 5 10	
TTC TCT ATT AAC CAG GGC AGG	169
Phe Ser Ile Asn Gln Gly Arg	
15	

(2) INFORMATION FOR SEQ ID NO: 86:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 157 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 29..70  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.4  
seq LIVLTLHSPSCDT/AQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

ATGGGGGTTT-CTTTGTGCT GCTGGGTG ATG CTA ATA GTC CTG ACT CTC CAC 52  
Met Leu Ile Val Leu Thr Leu His  
-10

TCG CCC TCC TGT GAG ACT GCC CAG GAG GAG ATG GGG AGG GTG CCC ACT 100  
Ser Pro Ser Ser Cys Asp Thr Ala Gln Glu Glu Met Gly Arg Val Pro Thr  
-5 1 5 10

ACT CCC AAG TGC AGG TGG AAG TTA GGG CTC TCC ATG TGT TCT TTG CTG 148  
Thr Pro Lys Cys Arg Trp Lys Leu Gly Leu Ser Met Cys Ser Leu Leu  
15 20 25

ACA CCT GGG 157  
Thr Pro Gly

## (2) INFORMATION FOR SEQ ID NO: 87:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 66..251
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq SVLNLGALGLTIQ/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

AACTCCCA GA ATGCTGACCA AAGTGGGAGG AGCACTAGGT CTTCCCGTCA CCTCCACCTC 60

TCTCC ATG ACC CGG CTC TGC TTA CCC AGA CCC GAA GCA CGT GAG GAT CCG 110  
Met Thr Arg Leu Cys Leu Pro Arg Pro Glu Ala Arg Glu Asp Pro  
-60 -55 -50

ATC CCA GTT CCT CCA AGG GGC CTG GGT GCT GGG GAG GGG TCA GGT AGT 158  
Ile Pro Val Pro Pro Arg Gly Leu Gly Ala Gly Glu Gly Ser Gly Ser  
-45 -40 -35

CCA GTG CGT CCA CCT GTA TCC ACC TGG GGC CCT AGC TGG GCC CAG CTC 206  
Pro Val Arg Pro Pro Val Ser Thr Trp Gly Pro Ser Trp Ala Gln Leu  
-30 -25 -20

CTG GAC AGT GTC CTA TGG CTG GGG GCA CTA GGA CTG ACA ATC CAG GCA 254  
Leu Asp Ser Val Leu Trp Leu Gly Ala Leu Gly Leu Thr Ile Gln Ala  
-15 -10 -5

GTC TTT TCC ACC ACT GGC CCA GCC CTG CTG CTG CTT CTG GTC AGC TCG 302  
Val Phe Ser Thr Thr Gly Pro Ala Leu Leu Leu Leu Val Ser Phe

```

      5          10          15
CTC ACC TTT GAC CTG CTC CAT AGG CCC GCA GTC ACA CTC TGC CAC AGC   350
Leu Thr Phe Asp Leu Leu His Arg Pro Ala Val Thr Leu Cys His Ser
      20          25          30

GCA AAC TTC TCA CCA GGG GCC AGA GTC AGG GGG CCG GTG AAG GTC CTG   398
Ala Asn Phe Ser Pro Gly Ala Arg Val Arg Gly Pro Val Lys Val Leu
      35          40          45

GAC AGC AGG AGG CTC TAC TCC TGC AAA TGG GTA CAG TCT   437
Asp Ser Arg Arg Leu Tyr Ser Cys Lys Trp Val Gln Ser
      50          55          60

```

## (2) INFORMATION FOR SEQ ID NO: 88:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 237 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 133..177
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq LTCLFLSLISTYP/SC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

```

ATTATGGTA GAGAGATATA TTTGTATTGG TTCCAGTTCC ATTGGTTTGT GAAATATTAA   60
TATGCCAACA CAGCCTAGCA TATTGGAGTC ACTGGAAATG CATCAGTGCT AGCCTTACAT   120
GCCTTTTCACT CT ATG GTG TTA ACC TGC CTT TTT CTA AGT CTA ATC TCC ACT   171
      Met Val Leu Thr Cys Leu Phe Leu Ser Leu Ile Ser Thr
      -15          -10          -5

TAC CCC AGC TGT ATC ACA CTT TTT CTT TCC AAA ATT CCT AAT CCT CTG   219
Tyr Pro Ser Cys Ile Thr Leu Phe Leu Ser Lys Ile Pro Asn Pro Leu
      1          5          10

TCT TCA CTC CCC TCA CTG   237
Ser Ser Leu Pro Ser Leu
      15          20

```

## (2) INFORMATION FOR SEQ ID NO: 89:

## (i) SEQUENCE CHARACTERISTICS:



(A) LENGTH: 281 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 171..224  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 6.3  
 seq FSFSLQLLSSSST/NP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

```

ATCTGTCTCT TGTTTATTAA GATATGCACA GTTCTGAAT CAACAAATAT ATCTGTGATT   60
CTTTTATACT ACTACATAAA AGAACAGGGR GTAATTCTTG CCTTATAAAT TAAATGTCAA   120
ACATTTCCCTA TATGTAATCA TTTGTTCCCTA AAATATGATT TAGTCCCAGC ATG CTT     176
                                     Met Leu
ATC CCT GTT TTC TCT TTT TCT CTC CAG CTC CTA TCT AGT TCT TCA ACA     224
Ile Pro Val Phe Ser Phe Ser Leu Gln Leu Leu Ser Ser Ser Ser Thr
  -15                               -10                               -5
AAT CCT GTC AAC TCT ACC TTC CAA ATG CCT TTT GAA TCC AGC CAT STC     272
Asn Pro Val Asn Ser Thr Phe Gln Met Pro Phe Glu Ser Ser His Xaa
   1                               5                               10                               15
ACC ACC AGA
Thr Thr Arg
                                     281
  
```

(2) INFORMATION FOR SEQ ID NO: 90:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 206 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 15..155  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 6.3  
 seq LLLLESVSGLLQP/RT

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

```

AAACCCGGGG GAAG ATG GCG GCA GCG TNT CTG AGT GGG CCC TCT GCG GGC      50
      Met Ala Ala Ala Xaa Leu Ser Gly Pro Ser Ala Gly
              -45                      -40

TCC GCG GCT GGG GTT CCT GGC GGG ACC GGG GGT CTC TCG GCA GTR ACC      90
Ser Ala Ala Gly Val Pro Gly Gly Thr Gly Gly Leu Ser Ala Val Ser
-35              -30              -25              -20

TCG GGC CCG CGG CTC CGC CTG CTG CTG CTG GAG AGT GTT TCT GGT TTG      140
Ser Gly Pro Arg Leu Arg Leu Leu Leu Leu Glu Ser Val Ser Gly Leu
              -15              -10              -5

CTG CAA CCT CGA ACG GGG TCT GCC GTT GCT CCG GTG CAT CCC CCA AAC      194
Leu Gln Pro Arg Thr Gly Ser Ala Val Ala Pro Val His Pro Pro Asn
              1              5              10

CGC TCG GCA AGG      206
Arg Ser Ala Arg
      15

```

## (2) INFORMATION FOR SEQ ID NO: 91:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 78..122
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2  
seq NWLFLEFVETFCNC/FF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

```

CATCTTGTTAC ATCTGKAGC ATGTATCTGT GAACATATCC ATAGGCTGGA TACCTAGCAG      60

GTCAAAATGA CGTGTGC ATG CAT AAT TGG CTT TTT TTG TTT GTW TTT ACT      110
      Met His Asn Trp Leu Phe Leu Phe Val Phe Thr
      -15              -10              -5

TTT TGT AAC TGC TTT TTT AAA AAT AAT GGC      140
Phe Cys Asn Cys Phe Phe Lys Asn Asn Gly
      1              5

```

## (2) INFORMATION FOR SEQ ID NO: 92:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 352 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 245..295
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2  
seq CFYFLSTALGSQA/DS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

```

ACTACCAATG GAAAATGCAG CTCTTGAGGA TGACGATTGC CAAACAAAGG CTCGGAGACG   60
AAGCAATCGG CGTGCGACAC TTTGCAGCCC ATGAGCGTGA AGACTTGGTG CAGCAGCTAG   120
AGCGAGCTAA GGAACAGGTT CTCACTAACA TCTATTCAGA GTGGGGGATG CATTTCACA   180
GCTGGACACA ACACAAACAA GAGTGGACTG TGCCCCCTCGT TTCTCAGAGT ATGGGGTGCC   240
TGGG ATG CAC GTT GAA TGC TTT TAC TTC CTC AGC ACT GCA CTA GGG TCC   289
  Met His Val Glu Cys Phe Tyr Phe Leu Ser Thr Ala Leu Gly Ser
    -15                      -10                      -5

CAA GCT GAC TCT TGG GTT TCT GGC CTC CAG CAG GCA GGT CTG CTC CCT   337
  Gln Ala Asp Ser Trp Val Ser Gly Leu Gln Gln Ala Gly Leu Leu Pro
    1                      5                      10

GCT ATT GGG TAC CGG   352
Ala Ile Gly Tyr Arg
  15

```

## (2) INFORMATION FOR SEQ ID NO: 93:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 353 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 177..233
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.1  
seq LALLWSLPASDLG/RS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

```

ATAAGTGAAC CAGACCACCC TGATGGCATC CACAGTGATG TCAAGGTTGG GGCTGGCCAG   60
GGGTGGGTGG ACTAGAAGCA TTTGGGAGTA GTGGCCAGGG GCCCTGGACG CTAGCCACGG  120
AGCTGCTGCA CAGAGCCTGG TGTCCACAAG CTTCCAGGTT GGGGTTGGAG COTGGG ATG  179
                                         Met
AGC CCC GGC AGC GCC TTG GCC CTT CTG TGG TCC CTG CCA GCC TCT GAC   227
Ser Pro Gly Ser Ala Leu Ala Leu Leu Trp Ser Leu Pro Ala Ser Asp
      -15                      -10                      -5
CTG GGC CGG TCA GTC ATT GCT GGA CTC TGG CCA CAC ACT GGC GTT CTC   275
Leu Gly Arg Ser Val Ile Ala Gly Leu Trp Pro His Thr Gly Val Leu
      1                      5                      10
ATC CAC TTG GAA ACA AGC CAG TCT TTT CTG CAA GGT CAG TTG ACC AAG   323
Ile His Leu Glu Thr Ser Gln Ser Phe Leu Gln Gly Gln Leu Thr Lys
      15                      20                      25                      30
AGC ATA TTT CCC CTC TGT TGT ACA TCG TTG                               353
Ser Ile Phe Pro Leu Cys Cys Thr Ser Leu
      35                      40

```

## (2) INFORMATION FOR SEQ ID NO: 94:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 290 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 180..218
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.1  
seq MALALGSIPSSIA/SS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

```

ACAAGGTCAT GAGGCTGAAT TAGGCTTTGT GTGCTCCTTC TAGGTTCTCA GCGTGTGTCT   60

```

GGTCTTTTAC TTGCTTTAGA TGTCTGCCCC AGCCACTGTA GGCAGGAACA GGTCTCTTCC	120
TTGAGAACTC AAGAGGTTCT CAAGGTAGTA AACTTCAATGG TGCTCTTAGT TTAGTTGTGA	179
ATG GCC TTG GCC TTG GGG TCC ATC CCA AGT TCC ATA GCC AGC AGT TGG	227
Met Ala Leu Ala Leu Gly Ser Ile Pro Ser Ser Ile Ala Ser Ser Trp	
-10 -5 1	
GTC CAT GTC TCA CAT TTT TGT CCC TGT CTC CTC CAC ACA ACA TTG CCA	275
Val His Val Ser His Phe Cys Pro Cys Leu Leu His Thr Thr Leu Pro	
5 10 15	
CAG TCC ACC CCG AAG	290
Gln Ser Thr Pro Lys	
20	

(2) INFORMATION FOR SEQ ID NO: 95:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 31..78  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.1  
seq ELEFTLPSLVVHP/SH

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

AATAGTTCAC ATATTTATGT TTTTCCACAA ATG CTA GCA TTT TTG TTC TGC ACT	54
Mat Leu Ala Phe Leu Phe Cys Thr	
-15 -10	
CTG TTT TCT TTA GTA GTG CAT CCT TCA CAC ATA GAT TTA AAA TGC TCA	102
Leu Phe Ser Leu Val Val His Pro Ser His Ile Asp Leu Lys Cys Ser	
-5 1 5	
TTT TAT	108
Phe Tyr	
10	

(2) INFORMATION FOR SEQ ID NO: 96:

## (2) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 349 base pairs  
(3) TYPE: NUCLEIC ACID

-- (C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 32..139

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 6  
seq LLYTLQTISSLG/CF

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

```

AGGTGCAGGG GAGGTAAGGT GGGAGCAGGT C ATG GCT CAA ATG CCA CTG ACA      52
                               Met Ala Gln Met Pro Leu Thr
                               -35                               -30

GGC TCT TAC CAA GAT TTA GAA TAT TTT CTT GAA TGC ATG TTT CTC CAT      100
Gly Ser Tyr Gln Asp Leu Glu Tyr Phe Leu Glu Cys Met Phe Leu His
                               -25                               -20                               -15

TTA TTA TAT ACT CTT CAA ACA ATT TCC AGT TTA AGT GGT TGT TTT AAA      148
Leu Leu Tyr Thr Leu Gln Thr Ile Ser Ser Leu Ser Gly Cys Phe Lys
                               -10                               -5                               1

CAA TTT TTT TTC CAG TTA AAT TGT TTT TGT TGG GGA GAA ATT CTA TGG      196
Gln Phe Phe Phe Gln Leu Asn Cys Phe Cys Trp Gly Glu Ile Leu Trp
                               5                               10                               15

CAC TCT TCA TTC CTC CAT TCT GGA AGT TGT CTC TTG GTT TTG CTC ATT      244
His Ser Ser Phe Leu His Ser Gly Ser Cys Leu Leu Val Leu Leu Ile
                               20                               25                               30                               35

AAA AAA AAA AAG ATA TAT CTT CAA TCT CAC TAA ATC TAT ACA GGT TAC      292
Lys Lys Lys Lys Ile Tyr Leu Gln Ser Xaa Xaa Xaa Ile Tyr Thr Gly Tyr
                               40                               45                               50

TTW ATA GAT YCT WAA YCT TTA SGT YCC TTC TCC ATC CCT TTA AGT TTC      340
Xaa Ile Asp Xaa Xaa Xaa Leu Xaa Xaa Phe Ser Ile Pro Leu Ser Phe
                               55                               60                               65

ATA CAG TTT
Ile Gln Phe
                               70

```

(2) INFORMATION FOR SEQ ID NO: 97:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 150 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(iii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 91..135

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 6  
seq LLMGLWVRTVLOG/KE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

```

AATAAAGCAT ACAGAAACCC ACCTAAAATA GACTCAGGGA GGTAGGAGGT TTCCTAAGGG    60
CTGAGACTGA AAGATAATAG GGATTGCTTG ATG GCA TTG TTG ATG GGG CTG TGG    114
                               Met Ala Leu Leu Met Gly Leu Trp
                               -15                               -10

GTG AGA ACA GTG CTC CAG GGA AAA GAG GCC AGC GGG                    150
Val Arg Thr Val Leu Gln Gly Lys Glu Ala Ser Gly
      -5              1              5

```

(2) INFORMATION FOR SEQ ID NO: 98:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 180 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 100..156

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 6  
seq LAILIXSLKLITIG/IQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

```

ATTAAAGTTG GAGAGAGATT AGAGGCAGAA TTAACAGAAA GGAGATGTGA GAATCCAGTA    60
GTCATTTAAAT TTTAAAAAAC AGGTATTCAA TAAAATTTT ATG ATT AAC CAT TTA    114
                               Met Ile Asn His Leu
                               -15

TAT TTG GCT ATT CTT ATT KTT TCT TTA AAA TTA ACA ATA GGA ATC CAG    162
Tyr Leu Ala Ile Leu Ile Xaa Ser Leu Lys Leu Thr Ile Gly Ile Gln
      -10              5              1

```

AAA CGT TTC GGA CCA CCG  
 Lys Arg Phe Gly Pro Pro  
 5

130

## (2) INFORMATION FOR SEQ ID NO: 99:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 218 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 12..161
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6  
seq LLYLCSFPLPGTS/LF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

AAAACAAAAT T ATG GGT AGG CAA GGG ACT TTA GAA ATT GAG GGC ATT CTC	50
Met Gly Arg Gln Gly Thr Leu Glu Ile Glu Gly Ile Leu	
-50 -45 -40	
TGT GTC ATC ACT TGG TTA GAG GCA AAT CTA GGG AAA CAA AAA GAT GAG	98
Cys Val Ile Thr Trp Leu Glu Ala Asn Leu Gly Lys Gln Lys Asp Glu	
-35 -30 -25	
AAT CAC TAC TAT AAG AAA TTA TCC CTT TTA TAC CTT TGC TCA TTT CCA	146
Asn His Tyr Tyr Lys Lys Leu Ser Leu Leu Tyr Leu Cys Ser Phe Pro	
-20 -15 -10	
CTG CCT GGA ACG TCC CTT TTT CTT CTC TGC TCT TTC TCA TAT CTT ACT	194
Leu Pro Gly Thr Ser Leu Phe Leu Leu Cys Ser Phe Ser Tyr Leu Thr	
-5 1 5 10	
CAA AGA CTT TCC CAA GGT GGA GGG	218
Gln Arg Leu Ser Gln Gly Gly Gly	
15	

## (2) INFORMATION FOR SEQ ID NO: 100:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 394 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR



(iii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 173..289  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.9  
seq SAWWCVLLEWSQG/AS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

```

ACTGGGGAAA TTGAGCCTAA GAGAACAGAA AGTACTTGAG GTCCACAAAT GAATCTATGG   60
ATGAATGAGT GCTTATTCAT TCACTCATTT TTTAAAAAAA TCCATCCAC AAGTATGTCT  120
TAATCACTGC AGTGTAGGC ACATAGGGAC AAAATAGAAG ATTCCTGTCC TC ATG GAA  178
                                     Met Glu
CTC ACA AAC AAG CAA ACA GGA ACT GAC AGA CAT GAA CAG GTA CTA CGG   226
Leu Thr Asn Lys Gln Thr Gly Thr Asp Arg His Glu Gln Val Leu Arg
      -35                      -30                      -25
AGG GTA AAG CAA GAC AAG AGG ATA AGT GCA TGG TGG TGC GTT TTA CTG   274
Arg Val Lys Gln Asp Lys Arg Ile Ser Ala Trp Trp Cys Val Leu Leu
      -20                      -15                      -10
GAG TGG TCA CAG GGG GCC TCT CTG AGG AGG CAA CAT CGA GGG GAG ACA   322
Glu Trp Ser Gln Gly Ala Ser Leu Arg Arg Gln His Arg Gly Glu Thr
      -5                      1                      5                      10
AGC CCC AAA TCT GGG GAA AGA CTT TCC AGG CAG AGA GAA CAG CAA AAA   370
Ser Pro Lys Ser Gly Glu Arg Leu Ser Arg Gln Arg Glu Gln Gln Lys
          15                      20                      25
CCG CAG ATG AGT GAC AAG AGC CTG   394
Pro Gln Met Ser Asp Lys Ser Leu
          30                      35

```

(2) INFORMATION FOR SEQ ID NO: 101:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 213 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

-- (B) LOCATION: 4..69  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.9  
 seq VLGLLFSDTWA/PA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

ACA ATG GCC AAG CST CAA AAT CCT ACG TCA GTG CTA GGA CTG CTT TTT	48
Met Ala Lys Arg Gln Asn Pro Thr Ser Val Leu Gly Leu Leu Phe	
-20 -15 -10	
TCT ATA TCA GAC ACG TGG GCT CCT GCT GTG TCT TCC TGG AAA GCA GAG	96
Ser Ile Ser Asp Thr Trp Ala Pro Ala Val Ser Ser Trp Lys Ala Glu	
-5 1 5	
GCC AAG GAT GGA GCA GAC CAA GAG GAT GCC AGG WWA WAA TCA CAA AGA	144
Ala Lys Asp Gly Ala Asp Gln Glu Asp Ala Arg Xaa Xaa Ser Gln Arg	
10 15 20 25	
AGC CCA GAW AGC ACA GCT GGA AGC CAA GAA CCC TAT TTT TGG TTT GTG	192
Ser Pro Xaa Ser Thr Ala Gly Ser Gln Glu Pro Tyr Phe Trp Phe Val	
30 35 40	
TGG GTG GAA GGT GAG GGA CGG	213
Trp Val Glu Gly Glu Gly Arg	
45	

(2) INFORMATION FOR SEQ ID NO: 102:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 375 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 250..324  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.9  
 seq FCLSLQIFRVSLA/LA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:

ATAAGGCTAG TTCTATTTTG AAGCCTATGT GTTTTGTGAA ACACAAAAAA AAGTACAGAG	60
AAAGATCGCA TCCTTTTCTG GTAGGGGTTT TCAGGAAAAA GTAGAGTTT TGAATCATGT	120
TGGGATTTCT TGGGCGGTTA TTCTGCAGTG GTCAAAATGG GGGAGCATG TCTGTAAAAG	180
TGTTACTGAT ATGACTAACA CTAATGATC TACTTTCAAA CATTAAGCTTT TTCCTCTCCC	240

```

TCCCTGTTT ATG AAT GTT TTG CCC TTC TCT TAC TAT TAT ATC TTG TTT TGT 291
Met Asn Val Leu Pro Phe Ser Tyr Tyr Tyr Ile Leu Phe Cys
-25 -20 -15

TTG AGT TTA CAA ATT TTC AGA GTT TCC CTA GCT CTG GCA GAS ACT CAT 339
Leu Ser Leu Gln Ile Phe Arg Val Ser Leu Ala Leu Ala Xaa Thr His
-10 -5 1 5

GAG GTT CCT GTC TCT ACT CAT ACT AAC RAA TTG CAT 375
Glu Val Pro Val Ser Thr His Thr Asn Xaa Leu His
10 15

```

## (2) INFORMATION FOR SEQ ID NO: 103:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 190 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 17..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq FSYISXFLSPVCG/CS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

```

ATCAAAATTC TCTTTG ATG AAA TGT TTA AAA GTG AAC CCT TTT TTA TTT CTG 52
Met Lys Cys Leu Lys Val Asn Pro Phe Leu Phe Leu
-25 -20

GTW TTT AAT TTC TTT TCC TAC ATC AGT KGC TTT TTG TCA CCA GTA TGT 100
Val Phe Asn Phe Phe Ser Tyr Ile Ser Xaa Phe Leu Ser Pro Val Cys
-15 -10 -5

GGA TGT TCT GTC TGT AAT TTA AAA CAC TGG GAG AAT GAG CTT CTA TTT 143
Gly Cys Ser Val Cys Asn Leu Lys His Trp Glu Asn Glu Leu Leu Phe
1 5 10 15

CCT TCT CCC CAC TTT TTG CCA TAT AAA TTT TTN TTT CTT TTT 190
Pro Ser Pro His Phe Leu Pro Tyr Lys Phe Xaa Phe Leu Phe
20 25

```

## (2) INFORMATION FOR SEQ ID NO: 104:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 226 base pairs

.. (B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 74..172  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.8  
seq XLCLGMALCPQA/TR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

```

ATCTCTTGGC GTCTCAACGT TCGGATCAGC AGCTTTTTC CATCTCTCT CTCCACTTCT   60
TCAGTGAGCA GCC ATG AGT TGG ACT GTG CCT GTT GTG CGG GCC AGC CAG   109
      Met Ser Trp Thr Val Pro Val Val Arg Ala Ser Gln
      -30                               -25

AGA GTG AGC TCG GTG GGA GCG AAT KTC CTA TGC CTG GGG ATG GCC CTG   157
Arg Val Ser Ser Val Gly Ala Asn Xaa Leu Cys Leu Gly Met Ala Leu
-20                               -15                               -10

TGT CCG CGT CAA GCA ACG CGC ATC CCG CTC AAC GGC ACC TGG CTC TTC   205
Cys Pro Arg Gln Ala Thr Arg Ile Pro Leu Asn Gly Thr Trp Leu Phe
-5                               1                               5                               10

ACC CCC GTG AGC AAG ATG GCG                               226
Thr Pro Val Ser Lys Met Ala
      15

```

(2) INFORMATION FOR SEQ ID NO: 105:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 173 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 111..155  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.8  
seq FLXLNTLTTHVHS/SA

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 105:

ATCCGATACA GAACATGCAG TAATGTGGAC TGCCACCCAG AAGCAGGTGA TTTCCGAGCT 60  
 CAGCAATGCT CAGCTCATAA TGATGTCAAG CACCATGGCC AGTTTATGA ATG GGY 116  
 Met Gly  
 -15  
 TTC CTG WGT CTA ATG ACC CTG ACA ACC CAT GTT CAC TCA AST GCC AAG 164  
 Phe Leu Xaa Leu Met Thr Leu Thr Thr His Val His Ser Ser Ala Lys  
 -10 -5 1  
 CCA AAT GGG 173  
 Pro Asn Gly  
 5

## (2) INFORMATION FOR SEQ ID NO: 106:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 98 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 33..80
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7  
seq RVLLLAQLFLGSG/KT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:

AAATTCTCTG GGCCTGCTTG TCATCACTCC AG ATG TTG TTT AGA GTT CTT CTG 53  
 Met Leu Phe Arg Val Leu Leu  
 -15 -10  
 TTA GCA CAG CTG TTT CTA GGG TCT GGA AAA ACT CTA AGG ACC CCG 98  
 Leu Ala Gln Leu Phe Leu Gly Ser Gly Lys Thr Leu Arg Thr Pro  
 -5 1 5

## (2) INFORMATION FOR SEQ ID NO: 107:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 243 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(i) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 79..174  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.7  
seq SLPLSTSAPPLRG/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

```

AACAGTCCTG CCGGCTGGCT TGGGTGGGTG GTGGGCTGCG GGTAGGGGAG GGGATGGACC   60
GAGTCCCGGC TTGTCGGG ATG AGG GTT CCG GAA GAT CTG GCC AGT AAG ATT   121
      Met Arg Val Pro Glu Asp Leu Ala Ser Lys Ile
      -30                               -25

CTA CTC CCT GGC TGT GCA CCG GGT TCC CTA CCC CTG TCT ACG TCG GCT   159
Leu Leu Pro Gly Cys Ala Pro Gly Ser Leu Pro Leu Ser Thr Ser Ala
  -20                -15                -10

CCG CCA CTT CGC GGC TTG AGA CTA AAA GAG CAT CCC GGC AGG GGG CCT   207
Pro Pro Leu Arg Gly Leu Arg Leu Lys Glu His Pro Gly Arg Gly Pro
  -5                1                5                10

TCC AGC CCC AAA GCA GCC TGT CCA GAG ACC CCC GCG   243
Ser Ser Pro Lys Ala Ala Cys Pro Glu Thr Pro Ala
      15                20

```

(2) INFORMATION FOR SEQ ID NO: 108:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 182 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 63..155  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.7  
seq SDLCLCQCILARA/HD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

```

ACATTTCTAA CGAGTACTGC CCAATCATTT GGTAGAATGC CGCTTAACTT TGCCTTWGTT   60

```



## (2) INFORMATION FOR SEQ ID NO: 110:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 161 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 18..86
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.6  
seq LLNVACCIPFSSS/LF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

```

ATTTTCCAAA CATTGTG ATG CAC CTT TAT AGC TGT TCG TGT ATG CGC CTT      50
      Met His Leu Tyr Ser Cys Ser Cys Met Arg Leu
                        -20                      -15

TTA AAC GTG GCA TGC TGC ATA CCC TTT TCG AGC AGC CTG TTT CCG CAC      98
Leu Asn Val Ala Cys Cys Ile Pro Phe Ser Ser Ser Leu Phe Pro His
      -10                      -5                      1

ATT CTT TTC AAG TCA TTA AAC TAT TCC TTG ACG TCC TTT CTC AAG GCT      146
Ile Leu Phe Lys Ser Leu Asn Tyr Ser Leu Thr Ser Phe Leu Lys Ala
      5                      10                      15                      20

GTG CGT GGC CGG TGG      161
Val Arg Gly Arg Trp
                        25

```

## (2) INFORMATION FOR SEQ ID NO: 111:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 285 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 223..270



(C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.5  
 seq PLVLSPLSYQCSS/QG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

```

AATGTTTAAG ATCTGTTTAA AATTTAAAAC AATGAATTGA ATGCTCTAAG AGGCTCCTAC   60
AGGCGCTCCA GGCCACTCTC AGAGACTCCC AGGAGTTGTT GAACTATATT TGGAGAAAAC   120
AGCCAMTGAA TATTATCATT TCTCCTTTAA AGAGAGTTTG TAAGGGGGGA ACATGCATTT   180
TATCAGACAA TTTATCCAAA GCATTTCAGA ACATGAGTGC TG ATG AGG GCA CCT   234
                                   Met Arg Ala Pro
                                   -15
CTT GTG CTG AGT CCC CTC AGC TAT CAG TGT TCT TCT CAA GGA CAC ATT   282
Leu Val Leu Ser Pro Leu Ser Tyr Gln Cys Ser Ser Gln Gly His Ile
      -10                      -5                      1

TGG                                     285
Trp
5

```

(2) INFORMATION FOR SEQ ID NO: 112:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 262 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 146..253
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.5  
seq FTSMCILFHCLLS/FQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

```

AACTTGGGAC AAGARATCAA ACTTTAAAGA TGGTCTAAG CCCCTCTTAA AGGTCTGACT   60
GTGTCGGACC TCTAGAGCTA ATCTCACTAG ATGTGAGCCA TTGTTTATAT TCTAGCCATC   120
CTTTCATTTT ATTCTAGAAG ACCCC ATG CAA GTT CCC CAC CTA AGG GTC TGG   172
                                   Met Gln Val Pro His Leu Arg Val Trp
                                   -35                      -30
ACA CAG GTG AWA GAT ACC TTC ATT GGT TAT AWA AAT TTG GGA TTT ACA   220
Thr Gln Val Waa Asp Thr Phe Ile Gly Tyr Arg Asn Leu Gly Phe Thr

```

-25                      -20                      -15

AGT	ATG	TGC	ATA	TTG	TTC	CAC	TGT	CTT	GTT	AGC	TTT	CAG	AGG
Ser	Met	Cys	Ile	Leu	Phe	His	Cys	Leu	Leu	Ser	Phe	Gln	Arg
-10						-5					1		

262

(2) INFORMATION FOR SEQ ID NO: 113:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 193 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 46..153  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.4  
seq LWLMHQSFQKSNSS

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

ACTACGAATG CAGATGTGGA AACAACTTCT GTGCATCTCA TCGTT ATG CAG AAA CTC 57  
Met Gln Lys Leu  
-35

ATG GCT GTA CCT ATG ATT ACA AGA GCG CAG GGA GGA GAT ACT TGC ACG 105  
Met Ala Val Pro Met Ile Thr Arg Ala Gln Gly Gly Asp Thr Cys Thr  
-30 -25 -20

AGG CAA ATC CTG TGG TTA ATG CAC CAA AGC TTC CAA AAA TCT AAC TCT 153  
Arg Gln Ile Leu Trp Leu Met His Gln Ser Phe Gln Lys Ser Asn Ser  
-15 -10 -5

TCC TCT ACA TCT TAC TGT TCT GCC CAG GGG 183  
Ser Ser Thr Ser Tyr Cys Ser Ala Gln Gly  
1 5 10

(2) INFORMATION FOR SEQ ID NO: 114:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 162 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(11) MOLECULE TYPE: cDNA

(vii) ORIGINAL SOURCE:

-- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 1..135  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.4  
seq AHRSLCLWPACLC/AR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

```

ATG TGT RIA GCT GGG TTT WAT GAC CAC CCT CGT GCG GCC CGG CAC GCC      48
Met Cys Xaa Ala Gly Phe Xaa Asp His Pro Arg Ala Ala Arg His Ala
-45                      -40                      -35                      -30

CGC ACG TCC CGC CAC CCC CTC CCT TGG GTG TGT GTC TCT CAG CYC CCT      96
Arg Thr Ser Arg His Pro Leu Pro Trp Val Cys Val Ser Gln Xaa Pro
                      -25                      -20                      -15

GCA CAC CGT TCC CTA TGT CTG TGG CCC GCG TGC CTB TGT GCG CGT GTG      144
Ala His Arg Ser Leu Cys Leu Trp Pro Ala Cys Leu Cys Ala Arg Val
                      -10                      -5                      1

CTC CCC CCA GCG CCA GGN                                             162
Leu Pro Pro Ala Pro Gly
5

```

## (2) INFORMATION FOR SEQ ID NO: 115:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 127 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 62..115  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.4  
seq ILVSFILAALSLS/TT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

```

ATCGGACTGA ACGGATCGCT GCGAGGATTA TCTTACACTG AACTGATCAA GTACTTTGAA      60

A ATG ACT TCG AAA TTT ATC TTG GTG TCC TTC ATA CTT GGT GCA CTG AGT      109
Met Thr Ser Lys Phe Ile Leu Val Ser Phe Ile Leu Ala Ala Leu Ser
-15                      -10                      -5

```

CTT TCA ACC ACC ATA GGG  
 Leu Ser Thr Thr Ile Gly  
 1

127

## (2) INFORMATION FOR SEQ ID NO: 116:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 279..323
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4  
seq LLIFILTVHHTPS/LP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

ATCTTTTGTT TGAGTATCTT CAAGAAAAAT CTGTTGTGAG AAAGATCCTA AACATATGTA	60
TGTATAGATG CATATCTTTG AAAGCCTATG TGAATACCAA GGAATCTGA ACTTTTTCTT	120
TGGAGATGTT TACATAATAA ATCTATTTTC ATCAATCTGG CATATTTTTC TCCTAGCACT	180
GACTTACTGA ATGCCGCTGA CCACGTGCTG CCTCTCATGC TAAATGCTTA CTTAATTCAT	240
CACCAATTC TGTAGACTGT ACAGGCTAAA CACCTCTA ATG CAT TTA CTT ATT TTC	296
	Met His Leu Leu Ile Phe
	-15 -10
ATC CTC ACT GTC CAT CAC ACT CCC TCC CTC CCC TCG	332
Ile Leu Thr Val His His Thr Pro Ser Leu Pro Ser	
-5 1	

## (2) INFORMATION FOR SEQ ID NO: 117:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 188 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 129..176
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq SSLMWQLISQVYS/CN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

```

ACAGGAAGTT TGCCTAGAAG GAATAAATTA ACTCTTGTTA CTTGGTGAGA TCATGGAAGG   60
GAATGTAATT TGTTTTAGGT GGTGGTAATT GTGAGTTTGA GGCTGGCCCA GGAAATGAGT  120
TGTCAGAT ATG CTG TCA TCC TCA TTA ATG GTT CAG CTT ATT TCT CAG GTT   170
      Met Leu Ser Ser Ser Leu Met Val Gln Leu Ile Ser Gln Val
      -15                      -10                      -5

TAT AGT TGT ATG AGG AGG                               189
Tyr Ser Cys Met Arg Arg
      1

```

## (2) INFORMATION FOR SEQ ID NO: 118:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 146 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 57..98
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq FSYILCMLFCLFS/QD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

```

ACTAATCCCC TATTTAGGTT GTTTACTTTT AGGTATTCTG CATAGAGCTG TGATGG ATG   59
                                     Met

TTC TCA TAT ATA CTT TGC ATG CTT TTC TGC TTA TTT TCT CAG GAT AAA   107
Phe Ser Tyr Ile Leu Cys Met Leu Phe Cys Leu Phe Ser Gln Asp Lys
      -10                      -5                      1

TTT CTG GAA GTG ACA TTG TTG TGT GAA AGG TAC ATG CTT               146
Phe Leu Glu Val Thr Leu Leu Cys Glu Arg Tyr Met Leu
      5                      10                      15

```

## (2) INFORMATION FOR SEQ ID NO: 119:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 145 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 11..67
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2  
seq VTLAFSLVLSES/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

AYCWTCTTAA ATG TTA TTT TTA TAT TAT GTT ACA CTT GCA TTC TCT TTA	49
Met Leu Phe Leu Tyr Tyr Val Thr Leu Ala Phe Ser Leu	
-15 -10	
TTG GTG TTA TCA GAG TCA GCA GTA CTG AAA AGA AGA GAA ATC TTT TGR	97
Leu Val Leu Ser Glu Ser Ala Val Leu Lys Arg Arg Glu Ile Phe Xaa	
-5 1 5 10	
ACA GGG TTA GGT TGT GTG ACA GGG TTA GGT TGT GTG ACA GGG TTA CGG	145
Thr Gly Leu Gly Cys Val Thr Gly Leu Gly Cys Val Thr Gly Leu Arg	
15 20 25	

## (2) INFORMATION FOR SEQ ID NO: 120:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 235 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 143..184
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2  
seq LLSGLWLSSVKEC/DD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

```

AAGGAGTAGT GGCTTTGTTC CCAGCTCAGT GAAGGCTGGC ATGGTCTCTC CTGTCCACTT   60
CACTCTGGAT TCTTTAAGCC TGTGAATTAC TAGACATGGA TTCCATCTCC AATGTGGATG   120
CCTCTCTTCA CCACAAGART AC ATG CTC CTT TCT GGG CTG TGG CTT AGC TCG   172
                Met Leu Leu Ser Gly Leu Trp Leu Ser Ser
                        -10                        -5

GTC AAG GAG TGT GAT GAC TGG CGA GCA GAT GGC TGC CTT CCA TCC ATC   220
Val Lys Glu Cys Asp Asp Trp Arg Ala Asp Gly Cys Leu Pro Ser Ile
                1                5                10

GTC CAC CCC CTA AGG   235
Val His Pro Leu Arg
                15

```

## (2) INFORMATION FOR SEQ ID NO: 121:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 191 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 59..112
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2  
seq VFCFSWLMSSSSP/SI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

```

ATACAAAGGA AATTAGTATG TTCCTTGAGG TTCAGGGAAT CTATGTATAT TTCAGATC   58
ATG GTT GCA TTT TCA GTC TTC TGT TTT TCA TGG TTG ATG AGT TCA TCA   106
Met Val Ala Phe Ser Val Phe Cys Phe Ser Trp Leu Met Ser Ser Ser
                -15                -10                -5

AGT CCT TCC ATC TTT TGG AGT CAT TTC TAT TCA CCA TTC AAG GAT CTA   154
Ser Pro Ser Ile Phe Trp Ser His Phe Tyr Ser Pro Phe Lys Asp Leu
                1                5                10

TCT AAA ATG TAT AAT TAT GTC TCC CCG   181
Ser Lys Met Tyr Asn Tyr Val Ser Pro
                15                20

```

## (2) INFORMATION FOR SEQ ID NO: 122:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 248 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 123..170
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1  
seq LALGIGPPGCLQG/SP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

```
ATGTTTCTT AAGATCCAGA AGTTTTGCT TTAGCTTAAG GATGTGTGCA ATTTCCATG   60
TGGCTTCATA ATTCATCCAT GACTTTGAAT TTTAAATGG AGAGAAGTTG GCTTCCCAGG  120
AA ATG GTG CCC CTG GCC CTG GGC ATC GGC CCA CCT GGC TGT CTC CAA   167
  Met Val Pro Leu Ala Leu Gly Ile Gly Pro Pro Gly Cys Leu Gln
    -15                -10                -5
GGC TCT CCT TCC CAG TGG CTG GTG CGG GCT CCG GGA GCT CAG CTG AGT   215
Gly Ser Pro Ser Gln Trp Leu Val Arg Ala Pro Gly Ala Gln Leu Ser
   1             5             10             15
CCC ATT GGG GTG GCA ACG GAA AGG GAG CAG AGG   248
Pro Ile Gly Val Ala Thr Glu Arg Glu Gln Arg
      20                25
```

## (2) INFORMATION FOR SEQ ID NO: 123:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 186 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 64..159
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1  
seq LLWFCTAMRPGGA/GL



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 123:

```

AGGATTAAGC AAGCACAGCC CTAGTTGATC ACCCAGCATG AAAAGTCCTG GAATCTCTCA    60
GAG ATG AAC CTG TGT ATG GGA GTT TTG CTT AAA GTK GST ACT TCA AGA    105
  Met Asn Leu Cys Met Gly Val Leu Leu Lys Val Gly Thr Ser Arg
      -30                -25                -20

AGG TGC CTC TGT TTA CTT TGG TTT TGC ACT GCC ATG CGA CCA GGT GGT    156
Arg Cys Leu Cys Leu Leu Trp Phe Cys Thr Ala Met Arg Pro Gly Gly
      -15                -10                -5

GCA GGT CTC CCA AAT GCC ACC CCC GAA TGG    195
Ala Gly Leu Pro Asn Ala Thr Pro Glu Trp
      1                5

```

## (2) INFORMATION FOR SEQ ID NO: 124:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 159 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 112..153
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1  
seq SLAKSLFLRVARG/LG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 124:

```

ATTTTAAAGG GAAAGACCCG GAAACAGCAC ATTCTCTTTT TCCAGTAGCC GGAATTTGCA    60
ACTACATATA GTCGCAAAGA AGACTGGGAG GWWATCTTTA GTTGGGAAGC A ATG AGT    117
                               Met Ser

CTA GCA AAA TCT CTG TTT TTA AGG GTG GCA AGG GGA CTG GGG    159
Leu Ala Lys Ser Leu Phe Leu Arg Val Ala Arg Gly Leu Gly
      -10                -5                1

```

## (2) INFORMATION FOR SEQ ID NO: 125:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 342 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

..(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 61..114

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.1  
seq FLPSATLLLSAES/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 125:

```

AAGGGCTCTG CCTCTTCCT ATACCATGCT GTCTTCATA GCCTTCCTCC TGTCCTACTC   60
ATG AGA CTG CCT CCA TTT CTT CCT TCT GCA ACC CTG CTC CTA TCA GCT   108
Met Arg Leu Pro Pro Phe Leu Pro Ser Ala Thr Leu Leu Leu Ser Ala
      -15                      -10                      -5

GAA TCC TTC TTT CGG AGT GTT AGT GAG TAC CCG TCT CTC CCC AGC CCC   156
Glu Ser Phe Phe Arg Ser Val Ser Glu Tyr Pro Ser Leu Pro Ser Pro
      1                      5                      10

TCA GCT GGT GGG CCT GGG TGT GTC AGC GGC AAA TGG GGC TCT GGT TCC   204
Ser Ala Gly Gly Pro Gly Cys Val Ser Gly Lys Trp Gly Ser Gly Ser
      15                      20                      25                      30

AAT GGG CCA CTC TCA TCT CTC TCT TGT TCC TTG TGC AGA AAA CCT TTG   252
Asn Gly Pro Leu Ser Ser Leu Ser Cys Ser Leu Cys Arg Lys Pro Leu
      35                      40                      45

CTT CAC TCC ACT GCC CTC TCT AGT TCC CGA CCC TTT TTC TCT CCT GGC   300
Leu His Ser Thr Ala Leu Ser Ser Ser Arg Pro Phe Phe Ser Pro Gly
      50                      55                      60

TTT CCC TGC CAA ATT TCT CCA AGG AGT GGT CTA CAY CCT CTG   342
Phe Pro Cys Gln Ile Ser Pro Arg Ser Gly Leu His Pro Leu
      65                      70                      75

```

(2) INFORMATION FOR SEQ ID NO: 126:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 354 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

--(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 202..349  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5  
 seq PLLLLLREELVTG/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 126:

```

ATATTAGTT CCTTTATTTT TTCTTTTCA AATGAATGGC TTTTAAAGTA CATGTTATGT   60
GAAGTATTCA CAAACACTGG TGCTTCCATG ATTATTGAGG AACATGTGAT TTATAAAATG  120
CCTCACTGTT TTCCAAGATA CACGATTGCG TCTGGGCACA GTTGATTICT CCTTGCCTAC  180
TCCCCCTCGC CCCTCACCOC C ATG AGT GAC AGA AAA AGA ACT AAA TTC TCA   231
                  Met Ser Asp Arg Lys Arg Thr Lys Phe Ser
                  -45                      -40

TAT GTC CAA CTC CCA TGC CCA ATC TCC CTT CTC CCA CGC AGT TTT AAA   279
Tyr Val Gln Leu Pro Cys Pro Ile Ser Leu Leu Pro Arg Ser Phe Lys
                  -35                      -30                      -25

AGG GGA CAA ATC CCA GGT CCC TCG GCT CCA CCA CTT CTT CTT CTT CTG   327
Arg Gly Gln Ile Pro Gly Pro Ser Ala Pro Pro Leu Leu Leu Leu Leu
                  -20                      -15                      -10

CGT GAG GAG TTG GTT ACC GGG GCC GTG                               354
Arg Glu Glu Leu Val Thr Gly Ala Val
                  -5                      1
  
```

(2) INFORMATION FOR SEQ ID NO: 127:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 248 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 12..134  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5  
 seq FCFPPAFLVXVS/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 127:

```

ACTCTTCGTT T ATG ACT CCG TTC CCC TCC GGC CCT CCT AGA GAG GCC TCC   50
Met Thr Pro Leu Gly Ser Gly Pro Pro Arg Glu Ala Ser
                  -40                      -35                      -30
  
```

ATA GCG CAG-GTT CGT GGG TTC TCG CGG ACC TTT TTC CGT GTA GCT TTC	98
Ile Ala Gln Val Arg Gly Phe Ser Arg Thr Phe Phe Arg Val Ala Phe	
-25 -20 -15	
TGC TTC TTC CCG GCA TTC CTT GTT WCG GTT TTM TCA CAG CCC TCT GGM	146
Cys Phe Phe Pro Ala Phe Leu Val Xaa Val Xaa Ser Gln Pro Ser Gly	
-10 -5 1	
TTT TCC ACC ACT GAG ACA CTT TGC GCT CAG GAC TTC AST GAC GTC ATC	194
Phe Ser Thr Thr Glu Thr Leu Cys Ala Gln Asp Phe Ser Asp Val Ile	
5 10 15 20	
TTT CTG CGG CGC GCR GAC ACC CGC CGG TGG AAG AAG AAA CAG CTC CGC	242
Phe Leu Arg Arg Ala Asp Thr Arg Arg Trp Lys Lys Lys Gln Leu Arg	
25 30 35	
CGC CGG	248
Arg Arg	

## (2) INFORMATION FOR SEQ ID NO: 128:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 242 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 93..137
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq CSALFPLLSLLSC/KE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 128:

ATGTGCTGAA ACTTAATCAG CAATGTGATG GTAATAGGTG GGGCCTTTAA AGGTGATTAA	60
GTCATGTGAG TGACCTTTAT AAAAAAGGCT TC ATG CGT TGT TCA GCT CTC TTT	113
Met Arg Cys Ser Ala Leu Phe	
-15 -10	
CCC CTT CTA TCT CTT TTG TCA TGC AAA GAG AGG ATR TGG TGT TTG TCC	161
Pro Leu Leu Ser Leu Leu Ser Cys Lys Glu Arg Xaa Trp Cys Leu Ser	
-5 1 5	
ACA TTG GAG GAT GCA GCG ACA DGG CGT CAC CTT GGA AST AGA GAG CAG	209
Thr Leu Glu Asp Ala Ala Thr Xaa Arg His Leu Gly Ser Arg Glu Gln	
10 15 20	
CCC TCA GGG GAT GCT GAG CCT GTG GAA GTA TGG	242
Pro Ser Gly Asp Ala Glu Pro Val Glu Val Trp	

25

30

35

## (2) INFORMATION FOR SEQ ID NO: 129:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 145 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 41..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq IISLLKLCSECFI/KD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 129:

```
AATTTAAGAT AATATCCAGT TCATGTAGAC ATGAATATAT ATG CTT TAT GAT CAA      55
                                     Met Leu Tyr Asp Gln
                                     -20

TAT TAC CTG ATA ATA TCA CTA CTA AAG CTA TGT TCT TTT TGC TTT ATT      103
Tyr Tyr Leu Ile Ile Ser Leu Leu Lys Leu Cys Ser Phe Cys Phe Ile
-15                               -10                               -5

AAA GAT TTT AAA GCC AGC AAC ATC ACT TTG GTA GTG ATA TTG              145
Lys Asp Phe Lys Ala Ser Asn Ile Thr Leu Val Val Ile Leu
1                               5                               10
```

## (2) INFORMATION FOR SEQ ID NO: 130:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 295 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 71..265
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5

seq LCSFLSLRFCTLS/FM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 130:

```

ASGAGACCGT GCCACCCCT AGATTSTCT TAAGCTCTTT TTTCATCTT TTAATTGCGT    60
AGACTCTGAA ATG GCT AAC TGT TTC CTA TCA CAT AAG AGC CAA ACT ATT    109
      Met Ala Asn Cys Phe Leu Ser His Lys Ser Gln Thr Ile
      -65                -60                -55

CTA ATT TCA AAG CCT GCT CTG ACT CAG TCC CAT TTT ACC TCT CCA GCC    157
Leu Ile Ser Lys Pro Ala Leu Thr Gln Ser His Phe Thr Ser Pro Ala
      -50                -45                -40

GGC TTG TTT CTA ACT GTT GAG AAA TCA CAC CTT TTG ACA AGG CTG TTT    205
Gly Leu Phe Leu Thr Val Glu Lys Ser His Leu Leu Thr Arg Leu Phe
      -35                -30                -25

TTT CAC TGG CTT TCG TTA GTG CTG TGC TCG TTT CTG TCT TTG AGA TTT    253
Phe His Trp Leu Ser Leu Val Leu Cys Ser Phe Leu Ser Leu Arg Phe
      -20                -15                -10                -5

TGC ACA TTA TCT TTT ATG TGC TCT TTT GCC CTT TTC CAC CTG          295
Cys Thr Leu Ser Phe Met Cys Ser Phe Ala Leu Phe His Leu
              1              5              10

```

## (2) INFORMATION FOR SEQ ID NO: 131:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 298 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 20..73
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq LTYLLFLPDWAAV/FE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 131:

```

AACGGACAGA TTTATTGGA ATG CAT GGA GCT GGT CTG ACC TAT TTA CTT TTC    52
      Met His Gly Ala Gly Leu Thr Tyr Leu Leu Phe
              -15                -10

CTT CCA GAC TGG GCT GCT GTA TTT GAA CTG TAC AAC TGT GAA GAT GAA    100
Leu Pro Asp Trp Ala Ala Val Phe Glu Leu Tyr Asn Cys Glu Asp Glu
      -5              1              5

```

```

CGC TGT TAC TTA GAC TTG GCC AGG CTG AGA GGC GTT CAC TAC ATC ACT    143
Arg Cys Tyr Leu Asp Leu Ala Arg Leu Arg Gly Val His Tyr Ile Thr
  10                      15                      20                      25

TGG CGA CGG CAG AAC AAA GTC TTT CCT CAG GAT AAG GGC CAC CAT CCA    196
Trp Arg Arg Gln Asn Lys Val Phe Pro Gln Asp Lys Gly His His Pro
          30                      35                      40

ACC CTG GGG GAG CAC CCG AAG TTC ACC AAC TAC TCT TTC GAT GTA GAA    244
Thr Leu Gly Glu His Pro Lys Phe Thr Asn Tyr Ser Phe Asp Val Glu
          45                      50                      55

GAA TTT ATG TAT CTT GTC CTT CAG GCT GCA GAC CAC GTA TTG CAA CAC    292
Glu Phe Met Tyr Leu Val Leu Gln Ala Ala Asp His Val Leu Gln His
          60                      65                      70

CCC GGG
Pro Gly
  75

```

## (2) INFORMATION FOR SEQ ID NO: 132:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 148 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 26..70
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9  
seq CLSATLAFSGSFL/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 132:

```

ACGCCAAACT CCTCCAGCTG GCCCC ATG TGC TGC CTT TCT GCC ACG CTA GCC    52
Met Cys Cys Leu Ser Ala Thr Leu Ala
          -15                      -10

TTT TCA GGC TCT TTT CTG GCT CCC CAC CTC ATC TTT TGC TGT TTC TTC    100
Phe Ser Gly Ser Phe Leu Ala Pro His Leu Ile Phe Cys Cys Phe Ser
  -5                      1                      5                      10

CAC CTG AAT GTC ATC ATC CTC CTA TCC TCA TTA TCC CCT ATC CAC GGG    148
His Leu Asn Val Ile Ile Leu Leu Ser Ser Leu Ser Pro Ile His Gly
          15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 133:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 172 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 38..154
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9  
seq SGLRGLLLQEALG/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 133:

```

AACAGAAGAA AGACAGCCTA GGAGCAGAGC CTCCCAG ATG GCT GAG TTG GAT CTA      55
                                     Met Ala Glu Leu Asp Leu
                                     -35

ATG GCT CCA GGG CCA CTG CCC AGG GCC ACT GCT CAG CCC CCA GCC CCT      103
Met Ala Pro Gly Pro Leu Pro Arg Ala Thr Ala Gln Pro Pro Ala Pro
-30 -25 -20

CTC AGC CCA GAC TCT GGG TTG AGG GGG CTG CTG TTG CAG GAG GCC CTG      151
Leu Ser Pro Asp Ser Gly Leu Arg Gly Leu Leu Leu Gln Glu Ala Leu
-15 -10 -5

GGA GCA GTG CCG GAC CCC AGG      172
Gly Ala Val Pro Asp Pro Arg
1 5

```

## (2) INFORMATION FOR SEQ ID NO: 134:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 370 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 203..286
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9  
seq FLVACPLFGVCLX/FF



## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 134:

```

ACTTCAGTAA ATCTATTATT GATGTAATAC TTTTGTCTAA TTASCATTCA TATTCTAATT    60
TTGTCACTTG TTCAATAATA TCCTTTTTGA CAATTTTTC TCCAGTGAGG GATCAAGTCT    120
AGGGCTGGAT ATTGTGTTTC ATTGTCAATG ATCTTGAGTC CCGTTTAATC TGSGAGAGTT    180
CCTCAGCTTT GCTTTGTGTC TT ATG ACA TTA ACA CAT GGG AAT AAT ATC CTC    232
           Met Thr Leu Thr His Gly Asn Asn Ile Leu
           -25                               -20

CAC CTC GCC AAC TTT TTT TTA GTA GCA TGT CCT TTA TTT GGG GTT TGC    280
His Leu Ala Asn Phe Phe Leu Val Ala Cys Pro Leu Phe Gly Val Cys
           -15                               -10                               -5

CTG AWR TTT TTC ATT CTT AGA TTC AGG TTA TAC ATT CAA GGC CCA AAT    328
Leu Xaa Phe Phe Ile Leu Arg Phe Arg Leu Tyr Ile Gln Gly Pro Asn
           1                               5                               10

GTC ACA CAA GTG ATA TTG CAT CTG TCT CAG GGA ACC TTG AGC    370
Val Thr Gln Val Ile Leu His Leu Ser Gln Gly Thr Leu Ser
           15                               20                               25

```

## (2) INFORMATION FOR SEQ ID NO: 135:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 181..222
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9  
seq VLRWLPWPRGSHS/DS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 135:

```

AAGTATCCAG CCTCAACATT CAGCAGAGGC CCCAGATCAG CGTCTGAGCC AGGCCAACAA    60
TGACCAAGGA GGATGGGATC CTGGGTGCAG CTCATCAGAA GCGTCGGGTG AGTCCGAGGC    120
CCCAGCTCTC TGCCCTCCTG KTCCTCTGCT CTCTCCTGCT CCTCCCACTT CTACTGGCTC    180
ATG GTG TTG AGA TGG TTG CCT TGG CCT AGG GGG TCA CAC AGC GAC TCG    228
Met Val Leu Arg Trp Leu Pro Trp Pro Arg Gly Ser His Ser Asp Ser
           -10                               -5

```

## (2) INFORMATION FOR SEQ ID NO: 136:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 166 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 50..121
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq FSFLGTLFHKNS/ED

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 136:

```

ATAGTATTGA TGCTGGGTCA AACTAGTTAG GAGGATTTTC AGTTCTCCC ATG AAA GCA   58
                               Met Lys Ala

AGG CTC TCT GGT AAT CTG ATT TGT TTT TCT TTT CTA GGA ACC CTC TTT   106
Arg Leu Ser Gly Asn Leu Ile Cys Phe Ser Phe Leu Gly Thr Leu Phe
-20                               -15                               -10

CAT AAA TCA AAC TCA GAA GAC AGC TCT GTA GGA AAA GGA GAC TGG AAG   154
His Lys Ser Asn Ser Glu Ile Ser Ser Val Gly Lys Gly Asp Trp Lys
-5                               1                               5                               10

AAG AAA AAT AAG   166
Lys Lys Asn Lys
                               15

```

## (2) INFORMATION FOR SEQ ID NO: 137:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 217 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 107..154

..(C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.8  
 seq VCLVPQTPLCLG/KG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 137:

```

AATGAACGAA CGGGGAAAGT GCATGTTGTA GTTCTCAAAA CCCAAAAAAA TCTAAGAGAA   60
ACCCAGCAGC AAGAAACACA GAGGTTTGGG TGTGAGCATC GGAGGA ATG TCT CAC   115
                                     Met Ser His
                                     -15

GTC TGC CTT GTC CCC CAG ACC CCG TCC CTG TGT CTG GGC AAA GGC ACG   163
Val Cys Leu Val Pro Gln Thr Pro Ser Leu Cys Leu Gly Lys Gly Thr
      -10                      -5                      1

CCC CGC TCC AGG TCG GCC CCA TTT CAG AGC AGT GGC CCT CAT AGG CTT   211
Pro Arg Ser Arg Ser Ala Pro Phe Gln Ser Ser Gly Pro His Arg Leu
      5                      10                      15

TGT GCG                                     217
Cys Ala
20

```

(2) INFORMATION FOR SEQ ID NO: 138:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 296 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 93..179
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq VLTSVNLFIGING/SV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 138:

```

ACTGCTTCCA GCAKAAGTCC TATGTGTCTT CCACCAATCT GCCTGTGCTA GCCCTTCTAC   60
TTTIGCTGTA TGGGTGGTCA ATCACACCTC TC ATG TAC CCA GCC TCC TTT GTG   113
                                     Met Tyr Pro Ala Ser Phe Val
                                     -25

TTC AAG ATC CCC ACC ACA GCC TAT GTG GTG CTC ACC AGC GTG AAC CTC   161
Phe Lys Ile Pro Ser Thr Ala Tyr Val Val Leu Thr Ser Val Asn Leu
      -20                      -15                      -10

```

```

TTC ATT GGC ATT AAT GGC AGC GTG GCC ACC TTT GTG CTG GAG CTG TTC      204
Phe Ile Gly Ile Asn Gly Ser Val Ala Thr Phe Val Leu Glu Leu Phe
   -5              1              5              10

ACC GAC AAT AAG CTG AAT AAT ATC AAT GAT ATC CTG AAG TCC GTG TTC      257
Thr Asp Asn Lys Leu Asn Asn Ile Asn Asp Ile Leu Lys Ser Val Phe
          15              20              25

TTG ATC TTC CCA CAT TTT TGC CTG GGA CGA GGG CAG ACG      296
Leu Ile Phe Pro His Phe Cys Leu Gly Arg Gly Gln Thr
          30              35

```

## (2) INFORMATION FOR SEQ ID NO: 139:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 290 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 165..254
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq RSSLWVTAPLVSA/CP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 139:

```

AAGAGCCTCT TGSATCCCCA CAGGGYAATG GGTGTBCCGA TCTCGCGGGG GACTCTGTGA      60
TCCGTGTTCC CCTGACCCTC CTAGTGACACA ACTTGCCCGG GCTCACTGGG CTCCTGCACC      120
ACTGCCTGTC AGGTCCGCTG CCAGCCCCAA GCCCCCACC AGCC ATG AGC TCC TCC      176
                               Met Ser Ser Ser
                               -30

AGA AAG GAC CAC CTC GGC GCC ASA GCT CAG AGC CCC TCC CGG TCA TCA      224
Arg Lys Asp His Leu Gly Ala Xaa Ala Gln Ser Pro Ser Arg Ser Ser
   -25              -20              -15

TTG TGG GTA ACG GCC CCT CTG GTA TCT GCC TGT CCT ACC TGC TCT CCS      272
Leu Trp Val Thr Ala Pro Leu Val Ser Ala Cys Pro Thr Cys Ser Pro
   -10              -5              1              5

GCT ACA CAC CCT ACC GGC      290
Ala Thr His Pro Thr Gly
          10

```

## (2) INFORMATION FOR SEQ ID NO: 140:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 397 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 230..286
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq ATYLVQSSACCPA/IV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 140:

```

ACCACGGGGA CAAGGACTGC KCCCACGATG GTGCTCCTGC CAVGCCCCAG CTGBACGGGG      60
AGTCCTGTGG GGGCCAGGCC TTGAACAGCC ACATGCCTGC TGAGACCGAG GAGCTGGGAC     120
GGTGGGGACC ACAGAGAGCA ACCTGATTAC CTCCTTGCTT GGGCTGTGCC AGAGCAAGAA     180
GAGTCGGGTG GCCTTGAAGG CCCAGGAGAA CCTGCTGCTC CTGGTGAGC ATG GCC TCC     238
                                   Met Ala Ser
CCA GCA GCT GCC ACC TAC CTG GTA CAG AGC AGC GCC TGC TGC CCT GCG     286
Pro Ala Ala Ala Thr Tyr Leu Val Gln Ser Ser Ala Cys Cys Pro Ala
  -15                               -10                               -5
ATC GTC CGG CAC CTT TGC CAG TBG TAC CGG TCC ATG CCT GTC TTC CTG     334
Ile Val Arg His Leu Cys Gln Xaa Tyr Arg Ser Met Pro Val Phe Leu
   1                               5                               10                               15
GAC CCC GCA GAS ATT GCC ACC TTA GAG GGC ATC AGC TGG AGG TTA CCC     382
Asp Pro Ala Xaa Ile Ala Thr Leu Glu Gly Ile Ser Trp Arg Leu Pro
      20                               25                               30
AGT GCC CCG TCT GAT                                           397
Ser Ala Pro Ser Asp
      35

```

## (2) INFORMATION FOR SEQ ID NO: 141:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 378 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 172..354  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.7  
seq LLPCNLHXS WLHS/SP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 141:

```

AGATTGGCTG GGCAGATGGG CTGACTGGCT GGGCAGATGG GTGGGTGAGT TCCCTCTCCC      60
CAGAGCCATC GGCCAGGTAC CAAAGCTCAG CTGTATGGAT TCCCAACAGG AGGACCTGCG      120
CTTCCCTGGG ACCCATTGTT GTACTGGATT AACAAGCGAC GGCCTACGG C ATG AAT      177
                                         Met Asn
                                         -60

GCA GCC ATC AAC ACG GGC CCT GCC CCT GCT GTC ACC AAG ACT GAG ACT      225
Ala Ala Ile Asn Thr Gly Pro Ala Pro Ala Val Thr Lys Thr Glu Thr
          -55                      -30                      -45

GAG GTC CAG AAT CCA GAT GTT CTG TGG GAT TTG GAC ATC CCC GAA GCC      273
Glu Val Gln Asn Pro Asp Val Leu Trp Asp Leu Asp Ile Pro Glu Ala
          -40                      -35                      -30

AGG AGC CAT GCT GAC CAA GAC AGC AAC CCB NMG GCG GAA GCC CTG CTC      321
Arg Ser His Ala Asp Gln Asp Ser Asn Pro Xaa Ala Glu Ala Leu Leu
          -25                      -20                      -15

CCC TGC AAC CTG CAC TGM AGC TGG CTC CAC AGC AGC CCC AGG CCA GAT      369
Pro Cys Asn Leu His Xaa Ser Trp Leu His Ser Ser Pro Arg Pro Asp
          -10                      -5                      1                      5

CCC CAT TCC      378
Pro His Ser

```

## (2) INFORMATION FOR SEQ ID NO: 142:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 362 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 130..308  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.7

seq GIFLVIFCSESFS/LL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 142:

```

AATGGAMTTC TGGGTGACA RATGTTTGT TGTGTTTGT TTATWCCTCA TCTGTTCTCT 60
ATTGTTTCTA GTTTGTAGTC AGCATTCATA GGTGTACTTG ATTCTCCTA TRWTATTAGN 120
NTCTAGCTGT TTTCAGGRAT TTCTCTTTKA TTTTGTAGTT CCAGTAGITT GACTATAAT 179

ATG ATA AAC CTA CTT GTG GGT AAC TGC ATT TAT CTG CTT GGA GCT ATT 227
Met Ile Asn Leu Leu Val Gly Asn Cys Ile Tyr Leu Leu Gly Ala Ile
      -40                -35                -30

AGA GCT TCT TGC ATG TGT AGA TKB ATG TCT TTC GCC AAA TTT GGG ATT 275
Arg Ala Ser Cys Met Cys Arg Xaa Met Ser Phe Ala Lys Phe Gly Ile
      -25                -20                -15

TTT CTT GTA ATA TTT TGT TCT GAA TCA TTT TCT CTT CTC CTC TGG AAC 323
Phe Leu Val Ile Phe Cys Ser Glu Ser Phe Ser Leu Leu Leu Trp Asn
      -10                -5                1                5

TTC AGT TCA ATA TAT GTT AAG ACC TTT TGG CCA GTG GGG 362
Phe Ser Ser Ile Tyr Val Lys Thr Phe Trp Pro Val Gly
      10                15

```

(2) INFORMATION FOR SEQ ID NO: 143:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 171 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 16..72
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq LRFLLRDPGCLLA/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 143:

```

AAGACGGCGG TCGGC ATG CTC TGT TGC GGT CCG CTT CGG TTT CTG TTG CGG 51
      Met Leu Cys Cys Gly Pro Leu Arg Phe Leu Leu Arg
                        -15                -10

GAC CCG GGG TGT CTC CTA GCG CAA CCG GAA CTA GCC TTC TGG GGG CCG 99
Asp Pro Gly Cys Leu Leu Ala Gln Pro Glu Leu Ala Phe Trp Gly Pro
      -5                1                5

```

GCT TCC TTT ATC TCT GGC GGC CTT GTA GTC GTC TCC GAG ACT CCC CAC 147  
 Ala Ser Phe Ile Ser Gly Gly Leu Val Val Val Ser Glu Thr Pro His  
 10 15 20 25

CCC TCC TTC CCT CTT GAC CCC CCG 171  
 Pro Ser Phe Pro Leu Asp Pro Pro  
 30

## (2) INFORMATION FOR SEQ ID NO: 144:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 360..416
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq ILLRMTVLPTLWT/RR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 144:

AAGAGAGAAA CTTGGCGATC ACGTTTTAC ATGATGCTCA CGCTCAGGGC GCTTCAATTA 60  
 TCCCTCCCCA CAAAGATAGG TGGCGCGTGT TTCAGGGTCT CTCGTCTCTC TCCTACAGAA 120  
 AAGAAAAAGA AAAAAATGTC ATTAGAAGAG GCGTAACACG TCAGTCCGTC CCCAGATCGA 180  
 GCCTGCGTGC TGCCGAAGCA GGGCGCCGAG TCCATGCGAA CTGCCACCTG ATCCGCTCTT 240  
 ATCAATGAAG CAGCCGATCA TGGCGGATGG CCCCCGGTGC AAGAGGCGCA AACAAGCCAA 300  
 TCCCAGGAGG AAAAACGTGG TGAACATGA CAATGTAGTG GACACAGGTT CTGAAACAG 359  
 ATG AGG AAG ACA AGC TTC ATA TTG CTG AGG ATG ACG GTA TTG CCA ACC 407  
 Met Arg Lys Thr Ser Phe Ile Leu Leu Arg Met Thr Val Leu Pro Thr  
 -15 -10 -5

CTC TGG ACC AGG AGA CGA GTC CAG CTA GTG 437  
 Leu Trp Thr Arg Arg Arg Val Gln Leu Val  
 1 5

## (2) INFORMATION FOR SEQ ID NO: 145:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 153 base pairs
- (B) TYPE: NUCLEIC ACID



..(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 31..99  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq VRVGLVVLVXRALC/LX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 145:

```

AGGGAAGGGA GGGCAGGCGG KGCTGGACTB ATG TGG TGG AAA CCT GCT CCT GAG      54
                               Met Trp Trp Lys Pro Ala Pro Glu
                               -20

GAA GGG GTC CGG GTG GGG TTG GTG CTT GTG TSA AGG GCT CTG TGC CTC      102
Glu Gly Val Arg Val Gly Leu Val Leu Val Xaa Arg Ala Leu Cys Leu
-15                      -10                      -5                      1

TKT GTA CTC TCT CGG TTC ATG TTC ASA AAT CCT GGC CTT GGT GGC ATG      150
Xaa Val Leu Ser Arg Phe Met Phe Xaa Asn Pro Gly Leu Gly Gly Met
                    5                      10                      15

GGG                      153
Gly

```

(2) INFORMATION FOR SEQ ID NO: 146:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 454 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 374..415  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq FNELLGNSSCVYQ/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 146:

```

AATTCTGGGC CAGCTTCTTC CCGAGCTCTA TCCTGCTTCC CTCATCTCC TATAGGATTC      60

```

```

TCCTTAGAGT TCTCCCTCCA TTAGTAGTTG TCTTAGGGTC TGTTCCTGGG GAGCCCTGCC 120
TAAGACTCAT GCTACAAGAA GTTAAATAAG TTTCCCGAAG TCACACAGCT AGCCTCTCAT 180
CCCTTTTCTA CTGAGAGGAA GTGGAATGCA CTCCGACAAG GATAAGGTTT TATTGTGAGC 240
TGGCCTTGSA ATTAAACCAC CACCAACACA CTTTTGGATT ATCAGNNGGT GGAAGGAGTG 300
CAAATGCCAG TTACGGTGAT GCGTTCAACA TCCTTATTTT CAGTTCAGAA TTTCCCTGGA 360
GCTCCAAATT TTT ATG TTT AAT TTC TTA CTG GGC AAT TCC AGT TGT GTA 409
          Met Phe Asn Phe Leu Leu Gly Asn Ser Ser Cys Val
                    -10                    -5

TAT CAA AGG CCC ATC AGA TTA AAA CTC ATT ATC TTC CCA TCA GGG 454
Tyr Gln Arg Pro Ile Arg Leu Lys Leu Ile Ile Phe Pro Ser Gly
      1             5             10

```

## (2) INFORMATION FOR SEQ ID NO: 147:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 413 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 57..182
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq LDPVLSAPAFSA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 147:

```

DRAAACCGGA GCCACAGAGG ACAGGGTAGA GTCGCAGAAA GGAGAGACAC ACATAC ATG 59
                                     Met

AAA AGA GGA GCT TTC TCC AAT CTT AAT GAT TCC CAG CTC TCA GCC TCG 107
Lys Arg Gly Ala Phe Ser Asn Leu Asn Asp Ser Gln Leu Ser Ala Ser
-40             -35             -30

TTT CTG CAA CCC AGC CTG CAA GCA AAC TGT CCT GCT TTG GAC CCT GCT 155
Phe Leu Gln Pro Ser Leu Gln Ala Asn Cys Pro Ala Leu Asp Pro Ala
-25             -20             -15             -10

GTG TCA CTC TCC GCA CCA GCC TTT GCC TCT GCT CTT CGC TCT ATG AAG 203
Val Ser Leu Ser Ala Pro Ala Phe Ala Ser Ala Leu Arg Ser Met Lys
      -5             1             5

TCC TCC CAG GCT GCA CGG AAG GAC GAC TTT CTC AGG TCT CTT AGT GAT 251

```

Ser Ser Gln Ala Ala Arg Lys Asp Asp Phe Leu Arg Ser Leu Ser Asp  
 10 15 20  
 GGA GAC TCA GGG ACA TCA GAA CAC ATC TCA GCG GTG GTG ACT AGC CCT 299  
 Gly Asp Ser Gly Thr Ser Glu His Ile Ser Ala Val Val Thr Ser Pro  
 25 30 35  
 CGG ATT TCC TGC CAT GGT GCT GCC ATT CCC AMM GCM MGT GCC CWC TGM 347  
 Arg Ile Ser Cys His Gly Ala Ala Ile Pro Xaa Ala Xaa Ala Xaa Xaa  
 40 45 50 55  
 MTA GGC TGT TCC TGC TGM ACC GAA CGM MTC CTC MTG MCA CCG CCC TCC 395  
 Xaa Gly Cys Ser Cys Xaa Thr Glu Arg Xaa Leu Xaa Xaa Pro Pro Ser  
 60 65 70  
 CTC CTT TCT TTA GAA GCC 413  
 Leu Leu Ser Leu Glu Ala  
 75

## (2) INFORMATION FOR SEQ ID NO: 148:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 271 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 32..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq FFIFCSLNTLLLG/GV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 148:

AACAACATATC CTGCCTGCTG CTTGCTGCAC C ATG AAG TCT GCC AAG CTG GGA 52  
 Met Lys Ser Ala Lys Leu Gly  
 -20  
 TTT CTT CTA AGA TTC TTC ATC TTC TGC TCA TTG AAT ACC CTG TTA TTG 100  
 Phe Leu Leu Arg Phe Phe Ile Phe Cys Ser Leu Asn Thr Leu Leu Leu  
 -15 -10 -5  
 GGT GGT GTT AAT AAA ATT GCG GAG AAG ATA TGT GGA GAC CTC AAA GAT 148  
 Gly Gly Val Asn Lys Ile Ala Glu Lys Ile Cys Gly Asp Leu Lys Asp  
 1 5 10 15  
 CCC TGC AAA TTG GAC ATG AAT TTT GGA AGC TGC TAT GAA GTT CAC TTT 196  
 Pro Cys Lys Leu Asp Met Asn Phe Gly Ser Cys Tyr Glu Val His Phe  
 20 25 30

(2) INFORMATION FOR SEQ ID NO: 149:

```
(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 150 base pairs
    (B) TYPE: NUCLEIC ACID
    (C) STRANDEDNESS: DOUBLE
    (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:
    (A) ORGANISM: Homo Sapiens
    (F) TISSUE TYPE: Testis

(ix) FEATURE:
    (A) NAME/KEY: sig_peptide
    (B) LOCATION: 31..75
    (C) IDENTIFICATION METHOD: Von Heijne matrix
    (D) OTHER INFORMATION:  score 4.6
                           seq ILFPLHSVIGSHP/QC
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 149:

AAGTTGCCTG AGGACAGCAG TSCAGTTGAC ATG GAT ATT CTC TTT CCT CTG CAC	54
Met Asp Ile Leu Phe Pro Leu His	
-15 -10	
AGT GTT ATT GGG AGC CAT CCT CAG TGC CTC CCA GAG AGG WGG ACA GCG	102
Ser Val Ile Gly Ser His Pro Gln Cys Leu Pro Glu Arg Xaa Thr Ala	
-5 1 5	
AGA ATG ATC AAG CTG AAG TGG GGG AAT GGC TCA GGA TCG GAT TTC GGG	150
Arg Met Ile Lys Leu Lys Trp Gly Asn Gly Ser Gly Ser Asp Phe Gly	
10 15 20 25	

(2) INFORMATION FOR SEQ ID NO: 150:

```
(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 430 base pairs
      (B) TYPE: NUCLEIC ACID
      (C) STRANDEDNESS: DOUBLE
      (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:
      (A) ORGANISM: Homo Sapiens
      (E) TISSUE TYPE: Testis
```

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 275..355
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq FGILILLSQRQWS/KN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 150:

```

GTTGGAAAAC AGTTTGGCT CTGAGGACCC AGCAGTTGAC AACAGGAGG CCTGGGACAA   60
GAGCAGTATG AGAAGTCAGA TCGCCTCTTT TAATGTCCT AGTCAGTACA GGCCTCGCCA  120
GACAAGTCTC TCCTCARMNT CACTTGAAG AACAGGCCSD CTCTTCATGA TCCTGGGTTT  180
CCTAGACWTA TTTCCAGGAC TGTTATGGGG ATTAGGGCCA ACTGTAAAAG TGGCTGAGGA  240
GACTAGGTAA AGAGTGTGT CTCACCTTAG AACA ATG CTG AAG GTG TTT AGA GCC   295
                               Met Leu Lys Val Phe Arg Ala
                               -25

TGM CAT CCT AAA ATA TGC CAC TTT GGC ATA CTG ATT CTT CTG AGC CAG   343
Xaa His Pro Lys Ile Cys His Phe Gly Ile Leu Ile Leu Leu Ser Gln
-20                               -15                               -10                               -5

AGG CAA TGG AGC AAA AAC AGA TGC AGG GAA GGC TGT CTG ACC ACC CTC   391
Arg Gln Trp Ser Lys Asn Arg Cys Arg Glu Gly Cys Leu Thr Thr Leu
                               1                               5                               10

TTT CTG TTT GAA GCG GAA CAT AAA AGT TCC CTT GTG AAA               430
Phe Leu Phe Glu Ala Glu His Lys Ser Leu Val Lys
    15                               20                               25

```

## (2) INFORMATION FOR SEQ ID NO: 151:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 353 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 219..320
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq LXWRKLAASWTLS/QE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 151:

ACAATTAAAG CACACAGAAA ATGATGTGAC TCATCTTCAA AAGGAAATGA GCAATTGTAS 60  
 AGCAGGTGAA AACGCTGSCA TGGGTAGGTT CACTAAGGTG GGTGAGCAAG AAAGGACAGT 120  
 GGACACCCCTG CCGTCCCCCC AGCACCCCGT GGCTCATTGC TGCAGTCAGC TGGAGGAGAG 180  
 GTGGCAGAGG TTGCAGAGCC AGGTCATCTC GGAGCTGG ATG CTT GTA AGG AAT GCA 236  
 Met Leu Val Arg Asn Ala  
 -30  
 CGC AGG GGG TCC AGA GGG AGG TCT CCA TGG TGG AGG GCA GGG TGT CTM 294  
 Arg Arg Gly Ser Arg Gly Arg Ser Pro Trp Trp Arg Ala Gly Cys Leu  
 -25 -20 -15  
 RTA TGG AGA AAA CTT GCA GCA AGC TGG ACT CTA TCT CAG GAA ATC TTC 332  
 Xaa Trp Arg Lys Leu Ala Ala Ser Trp Thr Leu Ser Gln Glu Ile Phe  
 -10 -5 1  
 AGA GGA TCA AGG AAG GGC TCG 353  
 Arg Gly Ser Arg Lys Gly Ser  
 5 10

## (2) INFORMATION FOR SEQ ID NO: 152:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 61..147
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq FTLGLGYPIPTRL/QP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 152:

GATGGCGGCG ASGSGGACGG TSAAGGTTGC CTCGCGCCCG TCCGGGCTCT GATCCTCCCC 60  
 ATG ACT AAA GGG CAT CAC CAC CAG CAT CCC CTG CAT CCC CAC CCA CTC 108  
 Met Thr Lys Gly His His His Gln His Pro Leu His Pro His Pro Leu  
 -25 -20 -15  
 TTC ACC CTG GGC TTG GGA TAC CCC ATA CCC ACT CGC CTG CAA CCA TGC 156  
 Phe Thr Leu Gly Leu Gly Tyr Pro Ile Pro Thr Arg Leu Gln Pro Cys  
 -10 -5 1  
 ACA TTA AGC TCA GAC CCC CTT CTG GAC ATT ACC TGT TCC CTG AGA AGC 204  
 Thr Leu Ser Ser Asp Pro Leu Leu Asp Ile Thr Cys Ser Leu Arg Ser  
 5 10 15

CCA AGC TCT GGG  
Pro Ser Ser Gly  
20

216

## (2) INFORMATION FOR SEQ ID NO: 153:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 236 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 162..230
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq RLHILFIVCLARG/KG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 153:

```

AGCTGCTTAG TTTGCTAATT CTAGTGGTTC AAACCAGATT TCAAAATCTG GGCTAAATCT   60
CTGTCATGCT ATGACATGGC ATTTGACAGT AATTCCTGAA TATTTAATTG ATAGAAAAAC   120
AGAAAGCATG CATATTGTTT AGTACAATTG TGTGAAGTGC T ATG ACA TAT CAT KRC   176
                                   Met Thr Tyr His Xaa
                                   -20

ATA CAG TTT TCT GAA AGA CTG CAT ATT TTA TTC ATT GTA TGC CTA GCA   224
Ile Gln Phe Ser Glu Arg Leu His Ile Leu Phe Ile Val Cys Leu Ala
      -15                      -10                      -5

CCG GGA AAA GGG                               236
Arg Gly Lys Gly
      1

```

## (2) INFORMATION FOR SEQ ID NO: 154:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 230 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

-- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 9..145  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.5  
 seq LIYCGLSQPLTLG/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 154:

```

ATTGTATC ATG TCT CAA TTT CCT CTC TGC AGC CCT CCG TGG AAA CCA CTT      50
Met Ser Gln Phe Pro Leu Cys Ser Pro Pro Trp Lys Pro Leu
      -45                      -40                      -35

GTC AAG GTC TCC AGA AAC CTG AAA ATA AGG ATG TCC ATT CCA TGG CCA      98
Val Lys Val Ser Arg Asn Leu Lys Ile Arg Met Ser Ile Pro Trp Pro
      -30                      -25                      -20

CTC TCA GTC CTG ATT TAC TGT GGT CTC TCG CAG CCT TTG ACC CTG GGG     146
Leu Ser Val Leu Ile Tyr Cys Gly Leu Ser Gln Pro Leu Thr Leu Gly
      -15                      -10                      -5

GTC ACC TCT CCT TCC TTC CCC CAA AAC TCT TTC TTC CCT TGG CTT CCA     194
Val Thr Ser Pro Ser Phe Pro Gln Asn Ser Phe Phe Pro Trp Leu Pro
      1                      5                      10                      15

GAA CAC CCC ACT CAC CTG GTC TCC TCT ACC CCA CAG                       230
Glu His Pro Thr His Leu Val Ser Ser Thr Pro Gln
      20                      25
  
```

(2) INFORMATION FOR SEQ ID NO: 155:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 445 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 26..100  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.4  
 seq AMGFLLMFDLTSQ/QS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 155:

```

AAAAGGACAT GTGGTGTGC AATCC ATG TTC CGG AGT CTC ACC ACT GCA TT      52
Met Phe Arg Ser Leu Thr Thr Ala Phe
      -25                      -20
  
```



```

TTC AGA GAC GCC ATG GGC TTC TTA TTA ATG TTT GAC CTC ACC AGT CAA 100
Phe Arg Asp Ala Met Gly Phe Leu Leu Met Phe Asp Leu Thr Ser Gln
-15 -10 -5

CAG AGC TTC TTA AAT GTC AGA AAC TGG ATG AGC CAA CTG CAA GCA AAT 148
Gln Ser Phe Leu Asn Val Arg Asn Trp Met Ser Gln Leu Gln Ala Asn
1 5 10 15

GCT TAT TGT GAA AAT CCA GAT ATA GTA TTA ATT GGC AAC AAG GCA GAC 196
Ala Tyr Cys Glu Asn Pro Asp Ile Val Leu Ile Gly Asn Lys Ala Asp
20 25 30

CTA CCA GAT CAG AGG GAA GTC AAT GAA CGG CAA GCT CGG GAA CTG GCT 244
Leu Pro Asp Gln Arg Glu Val Asn Glu Arg Gln Ala Arg Glu Leu Ala
35 40 45

GAC AAA TAT GGC ATA CCA TAT TTT GAA ACA AGT GCA GCA ACT GGA CAG 292
Asp Lys Tyr Gly Ile Pro Tyr Phe Glu Thr Ser Ala Ala Thr Gly Gln
50 55 60

AAT GTG GAG AAA GCT GTA GAA ACC CTT TTG GAC TTA ATC ATG NRG CGA 340
Asn Val Glu Lys Ala Val Glu Thr Leu Leu Asp Leu Ile Met Xaa Arg
65 70 75 80

ATG GAA CAG TGT GTG GAG AAG ACA CAA ATC CCT GAT ACT GTC AAT GGT 388
Met Glu Gln Cys Val Glu Lys Thr Gln Ile Pro Asp Thr Val Asn Gly
85 90 95

GGA AAT TCT GGA AAC TTG GAT GGG GAA AGC CAC CAG AGA AGA AAT GTA 436
Gly Asn Ser Gly Asn Leu Asp Gly Glu Ser His Gln Arg Arg Asn Val
100 105 110

TCT GCT AGA 445
Ser Ala Arg
115

```

## (2) INFORMATION FOR SEQ ID NO: 156:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 319 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 185..295
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4  
seq LSYASSALSPCPX/AP

## (x) SEQUENCE DESCRIPTION: SEQ ID NO: 156:

```

ATCACCTTCT TCTCCATCCT TSTCTGGGCC ASTCCCCARC CCACTCCCTC TCCTGACCTG   60
CCCAGCCCCAA GTCAGCCTTC AGCAGCGGCT TTTCTGCACA CAGATATTCC AGGCCTACCT   120
GGCATTCCAG GACCTCCGMA ATGATGCTCC AGTCCCTTAC AAGCGCTTCC TGGATGAGGS   180
TGGC ATG GTG CTG ACC ACC CTC CCC TTG CCC TCT GCC AAC AGC CCT GTG   229
Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val
      -35                -30                -25

AAC ATG CCC ACC ACT GGC CCC AAC AGC CTG AGT TAT GCT AGC TCT GCC   277
Asn Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala
      -20                -15                -10

CTG TCC CCC TGT CTG AHC GCT CCA AAG TCC CCC CGA CTT GGG   319
Leu Ser Pro Cys Leu Xaa Ala Pro Lys Ser Pro Arg Leu Gly
      -5                1                5

```

## (2) INFORMATION FOR SEQ ID NO: 157:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 270 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 106..195
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq LLPTLPWLPSTRL/LS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 157:

```

AGCACAGCCG TGRRTGCCA GGTTCGGGTA GGAGGCCCTT TGGGGGRMNR ATTCTTTAGG   60
AAATTCCTTT AGAAGVAAAC AACTTGGGAC TGGATAGCGT GCGAT ATG CAG AGA AAT   117
Met Gln Arg Asn
      -30

GCA ACT TTC ATT CAT TTG CAG TTA GCG ATC CSC CCT TCC CTG CTC CCC   165
Ala Thr Phe Ile His Leu Gln Leu Ala Ile Arg Pro Ser Leu Leu Pro
      -25                -20                -15

ACC CTT GGT TGG CTC CCC AGT ACC CGC CTG CTG TCG CCC ACA CCC TTA   213
Thr Leu Phe Trp Leu Pro Ser Thr Arg Leu Leu Ser Pro Thr Pro Leu
      -10                -5                1                5

GGA CAG CTT CGT GGT CCC CCG GGA DCG CAG AGG GGC ATG CCT ACC GCT   261

```

Gly Gln Leu Arg Gly Pro Pro Gly Xaa Gln Arg Ala Met Pro Thr Ala  
 10 15 20

CAT TTA AGA  
 His Leu Arg  
 25

270

## (2) INFORMATION FOR SEQ ID NO: 158:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 109 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 50..94
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq ILFCFHSFHPLEFQ/DT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 158:

ACATATATCT ATCCTGACAA TATTTAGCAG TTCAAAAGGT AATAAGATT ATG AAT ATA 58  
 Met Asn Ile  
 -15

TTA TTT TGC TTT CAT TCT TTT CAC CCT CTA TTT CAA GAC ACT ATC GAA 106  
 Leu Phe Cys Phe His Ser Phe His Pro Leu Phe Gln Asp Thr Ile Glu  
 -10 -5 1

TTT 109  
 Phe  
 5

## (2) INFORMATION FOR SEQ ID NO: 159:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 371 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (1X) FEATURE:

- (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 198..257  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.3  
 seq FNFLFLVQLCIL/CD

## (X1) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

```

AAGATAAATT GGGGAATTCT AGGGAAACCC TTGAATACCA AGATAGAAAA CTAAAGTTTT    60
TACTTCATTT GGTTCATGGGA AACTTGCCTT GAGCATGGGA GTCAATAATT AGAAGCAAGT   120
KAAATTCAAA AAGTCGAACC CCATTCATAA AACCAGCTGA TAGTCTGAAA ATACGCTTTG   180
AGCTAAGCAA AGAATAC ATG TTG ACA AAT CGT AAC TAC TTT AAC TTC CTT       230
      Met Leu Thr Asn Arg Asn Tyr Phe Asn Phe Leu
      -20                      -15                      -10

TTT CTT STA CAA TTG TGC ATC CTG GCT TGT GAC AAT GCA TAC CTT CAG       278
Phe Leu Val Gln Leu Cys Ile Leu Ala Cys Asp Asn Ala Tyr Leu Gln
      -5                      1                      5

TCG TGT CCC CTC ACC TCA AAG ACT CCT CTG TTA CAA ACC CAC TCT GCT       326
Ser Cys Pro Leu Thr Ser Lys Thr Pro Leu Leu Gln Thr His Ser Ala
      10                      15                      20

CTT TTC TAT AAT AGT ACA TAT GGG ATT TTC CTA CTC CTA GGA GTG         371
Leu Phe Tyr Asn Ser Thr Tyr Gly Ile Phe Leu Leu Leu Gly Val
      25                      30                      35

```

## (2) INFORMATION FOR SEQ ID NO: 160:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 363 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

## (11) MOLECULE TYPE: CDNA

## (11) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

## (1X) FEATURE:

- (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 190..267  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.3  
 seq ALCRFVGMQPCTA/QT

## (11) SEQUENCE DESCRIPTION: SEQ ID NO: 160:

```

AATTCTTAAAC AGTTCATTG GAATGTAAGG TCAGGGCACC ACTGAGTTCA GTACTTCAAA    60
ATTCTCTTAA TCTACCTCTC CCCAGTGCAC AAAAACACTC TCCACACCAA GCTGCTGCTG   120

```

```

CTGGGATGGA GGGATGGCGT CAGGATTCAA GACTTTTTT CCTACCTGTT CAGCACTTCT 180
TTCAGCGAT ATG AAG TTA AAT CCA GGC CAA GTT CCC ACC TGG TGG GAA GCA 231
      Met Lys Leu Asn Pro Gly Gln Val Pro Thr Trp Trp Glu Ala
      -25                -20                -15

CTG TSC AGG TTC GTG GGG ATG CAG CCC TSC ACA GCC CAG ACT GGA CTC 279
Leu Cys Arg Phe Val Gly Met Gln Pro Cys Thr Ala Gln Thr Gly Leu
      -10                -5                1

CTT CCC CAT GGA ACT CAC AAC ACA CGG GAG AGG CAG AGA GAT CCA AGC 327
Leu Pro His Gly Thr His Asn Thr Arg Glu Arg Gln Arg Asp Pro Ser
      5                10                15                20

SCA CAG AAA AAC ACA AGA AGA TTC AGC CCT GTT GGG 363
Ala Gln Lys Asn Thr Arg Arg Phe Ser Pro Val Gly
      25                30

```

## (2) INFORMATION FOR SEQ ID NO: 161:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 186 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 97..177
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq LCLNLCPCSSSLL/SP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 161:

```

ACTCTTGTAG TGGMGCCGGC TTGCATCCCA GGTCGTSSCG GTTTGGTGC CTGAAGCAGG 60
GAGCGCGGAG TCGTTCCCGA GAGAGGCGGC CAGGCT ATG CTC GCC GGT TTC CGG 114
      Met Leu Ala Gly Phe Arg
      -25

CGT TCC GGT CCG GCC AGC CAG AGT CTC TGT CTC AAC CTG TGT CCG TGC 162
Arg Ser Ala Pro Ala Ser Gln Ser Leu Cys Leu Asn Leu Cys Pro Cys
      -20                -15                -10

TCC AGG AGT CTC CTC AGC CCG GCG 186
Ser Ser Ser Leu Leu Ser Pro Ala
      -5                1

```

## (2) INFORMATION FOR SEQ ID NO: 162:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 311 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 237..290
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq SFYLLFFLNDVPP/CP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 162:

```

ACAGRACAGG TTAAAGAGAT AATCATTTGG GACTCAAATG TCTCTCCCC CGGGCACTTG      60
CATATGGGAC ATTGAGTCCT TTTGTTTTCC CTTGATCTAT AGCTCTTACC CCTCTGCCCCA 120
GTAATTCCTT GAGGAAGAGG TAAAGATCAR AGTTGRTACT TTGTCCITTC CTTCKTCTT   180
CCCTTATTTT TAAAGCTGTC RSCCACACTG ATTCCTGCTC TAATAGCAGA GCAGAG ATG    239
                                     Met
AAG GAA GGA GCT TCC TTC TAT CTG CTT TTC TTT CTC AAT GAT GTC CCA      287
Lys Glu Gly Ala Ser Phe Tyr Leu Leu Phe Phe Leu Asn Asp Val Pro
      -15                      -10                      -5

CCA TGT CCC CCT CAC ACC CCC GGG                                      311
Pro Cys Pro Pro His Thr Pro Gly
      1                      5

```

## (2) INFORMATION FOR SEQ ID NO: 163:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 305..391
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.3  
seq ETLCLKLSSQSR/NR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 163:

```

ATATTTTCAGC TTCCACATTT TATTTCAACA ACATTAGTCA TCGTAGCTGC GTATTCCTGT 60
THTCAGTGTA GTACCGTGA GCAHTTATGT TCCTAGCACT GTTCCAGGTA CCCTGTGCGT 120
TATGAGGCAG GCACATCTCT CCTGAAAGAA TTTATATTCT TGTCAGGGAA ATAAGGCTTC 180
AGATAAGAAA AAATTCGGGG GAAAGTGCCT AATTCCTTCT ACCCTAACCT GCCTCCATTT 240
CCTCCCTCCT CCGAGTTGAG ATGATTGGGT CAGAGCCAGC TCTTCCTGGG CTTGGGAAGA 300
GGAG ATG GGG CTT GAG TGC TGC TGC CCC CCT CAT AAC CTC AGA GTC TAT 349
      Met Gly Leu Glu Cys Cys Cys Pro Pro His Asn Leu Arg Val Tyr
              -25                      -20                      -15

ATT GAG ACT CTC TTG CTC AAA CTC TCC TCG CAG AGT AGA ACG AAC AGG 397
Ile Glu Thr Leu Leu Lys Leu Ser Ser Gln Ser Arg Thr Asn Arg
              -10                      -5                      1

CTG 400
Leu

```

(2) INFORMATION FOR SEQ ID NO: 164:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 376 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 275..337
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq VLSIAASLLQCRL/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 164:

```

AATCTGAAC AGTCTAGTC TCAAGCATTT TGGATGAGGG ATACCCATCC TCTATTTTAT 60
CAGCATTTTC ATGCTGTATT AAAATGAAAT TGCCAACTCA GTTCAAAGGA ATTTTCTTTC 120
TTAGCTTTAC ATTGTTGATT CATGGTGGAG GCCAACAAAC TATCGACTGG TGGGTGGAT 180
AATTTGGTCC AGAGAGGTCC TTGTGACATA TCTCATGGCC CATTACCTAG GTGATGTGAG 240

```

TCTTCCCTTC TGCCTGCAAT AAAGTTTTGT TGGA ATG CAG CTA TGC CCA TTT ACT 295  
 Met Gln Leu Cys Pro Phe Thr  
 -20 -15  
 AST GTA TTG TCC ATA GCT GCT TCT CTG CTA CAA TGT AGA TTA GCA GTT 343  
 Ser Val Leu Ser Ile Ala Ala Ser Leu Leu Gln Cys Arg Leu Ala Val  
 -10 -5 1  
 GTA ACA GAG ACT ATA TGG CCC CCC CAG VNT TGG 376  
 Val Thr Glu Thr Ile Trp Pro Pro Gln Xaa Trp  
 5 10

## (2) INFORMATION FOR SEQ ID NO: 165:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 354 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 139..270
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq QLLFKLNSTWCRA/LQ

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 165:

AAGCCYAGAA GTTATAAGGA AAGCCCTTCC AACTTGATAC AGTTGCTTTT CTTTCCTGAA 60  
 TCCCTGTITT ACTGGAAATT TCATTGGATT TTGGGAGGAG AGAGGTCTGA AGGAAGGAAA 120  
 GGCCTGTTTT CTGCTGTA ATG GAT GTA ACA TGC TGC TTT GAT GCA GTT GAA 171  
 Met Asp Val Thr Cys Cys Phe Asp Ala Val Glu  
 -40 -35  
 GGA AGT GAC TTC AGG GTT TGC TGT CAT GGA TGC GTG TCT TGG CTG TGT 219  
 Gly Ser Asp Phe Arg Val Cys Cys His Gly Cys Val Ser Trp Leu Cys  
 -30 -25 -20  
 TTC CAG ATG CTG CAG CTT TTA TTC AAG CTT AAT AGC ACT TGG TGC AGA 267  
 Leu Gln Met Leu Gln Leu Leu Phe Lys Leu Asn Ser Thr Trp Cys Arg  
 -15 -10 -5  
 GTA CTC CAG AGT GAA ACC TCA TTG GCT TCC CGG CGC CTG TGG ATG TGG 315  
 Ala Leu Gln Ser Glu Thr Ser Leu Ala Ser Arg Arg Leu Trp Met Trp  
 1 5 10 15  
 TTT TCT CAT CTG AYG GAG TTC TTT ACT GTG ACC CCC TGG 354  
 Val Ser His Leu Xaa Glu Phe Phe Thr Val Thr Pro Trp  
 20 25



## (2) INFORMATION FOR SEQ ID NO: 166:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 84 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 7..72
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq HCFCFTLFSYSSS/FF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 166:

```
AAGGAC ATG AGA CAA GGA CCT GGG GCC CCA CTC CAT TGC TTC TGT TTC      48
    Met Arg Gln Gly Pro Gly Ala Pro Leu His Cys Phe Cys Phe
          -20                      -15                      -10

ACC CTT TTT TCC TAC TCC TCC TCC TTT TTT TTT TTT      84
Thr Leu Phe Ser Tyr Ser Ser Ser Phe Phe Phe Phe
          -5                      1
```

## (2) INFORMATION FOR SEQ ID NO: 167:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 72..116
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq ITLLGIWLTXRLQ/FP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 167:

```

ACAAAGCCCCC GGCTTGCTCA TTTCATCCAG GTGAGGAGTC TGGAGTAGAG CAGGGGCTTCT      60
GAAATGGTGA C ATG CAC ATC ACT CTC CTG GGC ATC TGG TTA ACA KGC AGG      110
      Met His Ile Thr Leu Leu Gly Ile Trp Leu Thr Xaa Arg
      -15 . -10 -5
CTC CAG TTC CCC AGG TCT GGG CGG GCT GGG      140
Leu Gln Phe Pro Arg Ser Gly Arg Ala Gly
      1 5

```

(2) INFORMATION FOR SEQ ID NO: 168:

- ```
(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 316 base pairs
      (B) TYPE: NUCLEIC ACID
      (C) STRANDEDNESS: DOUBLE
      (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:
      (A) ORGANISM: Homo Sapiens
      (F) TISSUE TYPE: Spleen

(ix) FEATURE:
      (A) NAME/KEY: sig_peptide
      (B) LOCATION: 245..295
      (C) IDENTIFICATION METHOD: Von Heijne matrix
      (D) OTHER INFORMATION:  score 4.2
                               seq SWVCLLSAGTAFE/DY
```

(xi) SEQUENCE DESCRIPTION: SEO ID NO: 168:

|                                                                  |            |            |            |            |            |    |     |
|------------------------------------------------------------------|------------|------------|------------|------------|------------|----|-----|
| ATTTAGGATT                                                       | TAGACITTA  | GGGATTTTGA | TCTTTGGGGA | TTTCAATATT | TGGGATTATG |    | 60  |
| GTATTTTGAGA                                                      | TGGTCTCTTT | TAGGATTATG | ATCCAAACCC | ATCTCAGGAA | TGTGTGAAAT |    | 120 |
| TTACAGTAGT                                                       | CCATCCCCAT | CCCGGGCTGT | AGAAATGTAG | GACCCACAAG | CCTTCGTTAC |    | 180 |
| AGAGCCACTT                                                       | ACTGCCCAT  | GGAGTTCCCA | GGTAGATGAC | AGTAGCGGGG | AGGATACATG |    | 240 |
| GCAC ATG TTA TAT GGC TCT TGG GTG TGC CTT CTC TCA GCA GGC ACT GCC |            |            |            |            |            |    | 289 |
| Met Leu Tyr Gly Ser Trp Val Cys Leu Leu Ser Ala Gly Thr Ala      |            |            |            |            |            |    |     |
| -15                                                              |            |            | -10        |            |            | -5 |     |
| TTT GAA GAT TAT CAT TTG GGG GGT ACG                              |            |            |            |            |            |    | 316 |
| Phe Glu Asp Tyr His Leu Gly Gly Thr                              |            |            |            |            |            |    |     |
| 1                                                                |            |            | 5          |            |            |    |     |

## (2) INFORMATION FOR SEQ ID NO: 169:

- (1) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 208 base pairs
  - (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: DOUBLE
  - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 59..154  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.2  
seq XXXXFLLGRVVG/ES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

```

ACTATTTTTC CCTTCATTGT CTTTACTTTG CTTCAGAAGA ATAGTGTGTG ATGACGCC      53
ATG TTA TTT TTT CCC CTT CTT TCT TTC CGA TTT CTA CCC TCA GAG AGT      106
Met Leu Phe Phe Pro Leu Leu Ser Phe Arg Phe Leu Pro Ser Glu Ser
      -30                      -25                      -20

TTG TTG AAA GKC BTA WTG SYT TTT TTG CTG GGG AGG AGG GTA GTA GGA      154
Leu Leu Lys Xaa Xaa Xaa Xaa Phe Leu Leu Gly Arg Arg Val Val Gly
      -15                      -10                      -5

GAA TCA CNT TTT ATT TTC ACA TGT GGA AAT TTG CTT TTA ATT TGG CCT      202
Glu Ser Xaa Phe Ile Phe Thr Cys Gly Asn Leu Leu Leu Ile Trp Pro
      1              5              10              15

TAC GGG      208
Tyr Gly

```

(2) INFORMATION FOR SEQ ID NO: 170:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 197 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 113..160  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.2  
seq WAILGCWGTLSRG/HL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 170:

```

ATGAAAATTT TCCATGCTGT YTGAGGAATC ATATTTTCTC CCAGCAGATC ATTATCATGA      60

```

GAGTTGAAAC AAAAAAATA CATGGAGGTG GAACCTSCCA GCCCAGTGGT GG ATG CCA 113.  
Met Pro  
-15

GTC TGG GCC ATA CTG GGC TGC TGG GGC ACA CTC AGC AGG GGA CAT CTG 156  
Val Trp Ala Ile Leu Gly Cys Trp Gly Thr Leu Ser Arg Gly His Leu  
-10 -5 1

CCT GTG TCC TTG GAC CCA AAG 187  
Pro Val Ser Leu Asp Pro Lys  
5

## (2) INFORMATION FOR SEQ ID NO: 171:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 253 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 134..247
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1  
seq GILCGSLPGPSLC/PP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 171:

ACAAGTTCAC CTGGCCTCCT CTTCTCCAGC CTCAGTCACC TTCTGCTGAA CAGCTCCACC 50

TTGGCCTTGC TTACTCAGAG ACTAAGCCAG ATGACCTGCC TGCAGAGCCT CAGGTGAGTG 120

ACCGAGCGGC CCC ATG GGA ATG AGT GGG AAG AAA CAC TTC CCA CTC AGT 159  
Met Gly Met Ser Gly Lys Lys His Phe Pro Leu Ser  
-35 -30

TGG GAC CAC ATC CAG GGA AGC ACT GAG GCC ACC TCC CAG GGG ATC CTT 217  
Trp Asp His Ile Gln Gly Ser Thr Glu Ala Thr Ser Gln Gly Ile Leu  
-25 -20 -15

TGC GGA TCC CTC CCA GGC CCA TCC CTG TGC CCT CCG 253  
Cys Gly Ser Leu Pro Gly Pro Ser Leu Cys Pro Pro  
-10 -5 1

## (2) INFORMATION FOR SEQ ID NO: 172:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 362 base pairs

- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 141..251
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq PLSLDCGHSLCRA/CI

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 172:

```

AACACCCACC CTGGCTTTTC TTCACCTCTT CAACCAGGAG CCGAGATTTT TGTGCTCTG   60
AAGCCATCCA GGGGTCTTTA ACCAGAAGAG AGAGGAGAGC CTCAGGAGTT AGGACCAGAA  120
GAAGCCAGGG AAKCAGTGCA ATG GCT TCA AAA ATC TTG CTT AAC GTA CAA GAG   173
          Met Ala Ser Lys Ile Leu Leu Asn Val Gln Glu
          -35                      -30

GAG GTG ACC TGT CCC ATC TGC CTG GAG CTG TTG ACA GAA CCC TTG AGT   221
Glu Val Thr Cys Pro Ile Cys Leu Glu Leu Leu Thr Glu Pro Leu Ser
  -25                      -20                      -15

CTA GAC TGT GGC CAC AGC CTC TGC CGA GCC TGC ATC ACT GTG AGC AAC   269
Leu Asp Cys Gly His Ser Leu Cys Arg Ala Cys Ile Thr Val Ser Asn
  -10                      -5                      1                      5

AAG GAG GCA GTG ACC AGC ATG GGA GGA AAA AGC AGC TGT CCT GTG TGT   317
Lys Glu Ala Val Thr Ser Met Gly Gly Lys Ser Ser Cys Pro Val Cys
      10                      15                      20

GGT ATC AGT KAC TCA KTT GAA CAT CTA CAG GCT AAT CAG CAT CGG       362
Gly Ile Ser Xaa Ser Xaa Glu His Leu Gln Ala Asn Gln His Arg
      25                      30                      35

```

(2) INFORMATION FOR SEQ ID NO: 173:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 173:

(2) INFORMATION FOR SEQ ID NO: 174:

(A) LENGTH: 158 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

```
(A) NAME/KEY: sig_peptide
(B) LOCATION: 15..122
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4
                        seq KLAGLWSPGLVPA/AP
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 174:

|                                                                 |                                                 |                                                 |     |
|-----------------------------------------------------------------|-------------------------------------------------|-------------------------------------------------|-----|
| AAGTGTGCGC                                                      | ATAA                                            | ATG GGC GCC GGC GGA SGG AGG GAG ATC CGA CCG GCG | 50  |
|                                                                 | Met Gly Ala Gly Gly Xaa Arg Glu Ile Arg Ala Ala | -35 -30 -25                                     |     |
| GCG GCA AGC TGG CTG CGA GCG GCT GAG CAC TCC AAG CTC GCC GGC CTT |                                                 |                                                 | 98  |
| Ala Ala Ser Trp Leu Arg Ala Ala Glu His Ser Lys Leu Ala Gly Leu | -20                                             | -15                                             | -10 |
| TGG TCT CCA GGA CTT GTC CCA GCA GCC CCT CGA ACT GAG AAT TAC ACC |                                                 |                                                 | 146 |
| Trp Ser Pro Gly Leu Val Pro Ala Ala Pro Arg Thr Glu Asn Tyr Thr | -5                                              | 1                                               | 5   |
| ATC GGA CCC CTG                                                 |                                                 |                                                 | 158 |
| Ile Gly Pro Leu                                                 | 10                                              |                                                 |     |

## (2) INFORMATION FOR SEQ ID NO: 175:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 291 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 52..231
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq LVRRTLLVAALRA/WM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 175:

```

AAGGAAACAG CAACCAGAGG GAGATGATCA CCTGAACCAC TGCTCCAAAC C ATG GGC      57
                                     Met Gly
                                     -60

AGT AAA TGC TGT AAA GGT GGT CCA GAT GAA GAT GCA GTA GAA AGA CAG      105
Ser Lys Cys Cys Lys Gly Gly Pro Asp Glu Asp Ala Val Glu Arg Gln
      -55                      -50                      -45

AGG CGG CAG AAG TTG CTT CTT GCA CAA CTG CAT CAC AGA AAA AGG GTG      153
Arg Arg Gln Lys Leu Leu Leu Ala Gln Leu His His Arg Lys Arg Val
      -40                      -35                      -30

AAR GCA GCT GGG CAG ATC CAG GCC TGG TGG CGT GGG CTC CTG GTG CGC      201
Lys Ala Ala Gly Gln Ile Gln Ala Trp Trp Arg Gly Val Leu Val Arg
      -25                      -20                      -15

AGG ACC CTG CTG GTT GCT GCC CTC AGG GCC TGG ATG ATT CAG TGC TGG      249
Arg Thr Leu Leu Val Ala Ala Leu Arg Ala Trp Met Ile Gln Cys Trp
      -10                      -5                      1                      5

TGG AGG ACG TTG GTG CAG AGA CGG ATC CGT CAG CGG CGG CAG      291
Trp Arg Thr Leu Val Gln Arg Arg Ile Arg Gln Arg Arg Gln
      10                      15                      20

```

## (2) INFORMATION FOR SEQ ID NO: 176:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 192 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 103..180  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4  
seq SIHSWQLLTSAP/QQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 176:

```
CGGCAACGCG CGGCGGCTCA ACGGCCTGGC AGAGGTTTCA GCGCTGAGCA GGCCTGAGTT    60
CCGTCATGGC CCTCTATTAT GACCACCAGA TAGAAGCCCC GG ATG CAG CAG GGT    114
                                     Met Gln Gln Gly
                                     -25
CAC CCT CAT TTA TCA GCT GGC ACC CTG TCC ATC CAT TCT TGG CAG TTG    162
His Pro His Leu Ser Ala Gly Thr Leu Ser Ile His Ser Trp Gln Leu
      -20                      -15                      -10
CTT ACA TCA GCA CAA CCT CAA CAG GCA GGG    192
Leu Thr Ser Ala Gln Pro Gln Gln Ala Gly
      -5                      1
```

(2) INFORMATION FOR SEQ ID NO: 177:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 174 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 1..147  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4  
seq ATCCLSLFQWCAV/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 177:

```
ATG TCT AGA TAT GAG TMA GGA TCC TCC TTA TTG CCA TTT CCA GAC CAT    48
Met Ser Arg Tyr Glu Xaa Gly Ser Ser Leu Leu Pro Phe Pro Asp His
      -45                      -40                      -35
TTC TCT GTT TAC TCC TTT AAA ASA RAT AGT TTT TTT GAA CCG TAC ACC    96
```



Phe Ser Val Tyr Ser Phe Lys Xaa Xaa Ser Phe Phe Glu Ala Tyr Ser  
 -30 -25 -20

ATT TCA GAT TAT GCC ACC TGC TGT CTC TCC TTA TTT CAG TGG TGT GCA 144  
 Ile Ser Asp Tyr Ala Thr Cys Cys Leu Ser Leu Phe Gln Trp Cys Ala  
 -15 -10 -5

GTT CTG AGA TTC CTG TCT CTG CCC CTT CCG 174  
 Val Leu Arg Phe Leu Ser Leu Pro Leu Pro  
 1 5

## (2) INFORMATION FOR SEQ ID NO: 178:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 226 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 140..211
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq LLLHHYLLLFITT/SR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 178:

ACAGTTGTGG CTCTCAACTC TCCTTTTGT GTACTGCTAT ACTTGAGTAG CACACAGCCA 60

TACCAATTTC CAGGGTGCTC AGATTCATTC TACCCTTTCC TACTGGAAGA GGTA AAAAAG 120

CAACACCTTA GAATCTGAT ATG ATT TAT TTT ATC AAA ATA AAC AAT AAG CTA 172  
 Met Ile Tyr Phe Ile Lys Ile Asn Asn Lys Leu  
 -20 -15

CTG CTT TTG CAC CAT TAC TTG CTT CTA TTT ATA ACA ACC TCT CGC CCC 220  
 Leu Leu Leu His His Tyr Leu Leu Leu Phe Ile Thr Thr Ser Arg Pro  
 -10 -5 1

ACA GGG 226  
 Thr Gly  
 5

## (2) INFORMATION FOR SEQ ID NO: 179:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 129 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

-- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 28..108

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.9  
seq LSWALCLSQSGYY/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 179:

```
AGGGTATATT TCNTGTCCCC TAGGAGC ATG GAG CTT TTG TAC CTT AAA GTT AAG      54
                               Met Glu Leu Leu Tyr Leu Lys Val Lys
                               -25                               -20

AGA GGA CAA AAG GAT CTG AGC TGG GCT TTG TGC CTT TCC CAG AGT GGT      102
Arg Gly Gln Lys Asp Leu Ser Trp Ala Leu Cys Leu Ser Gln Ser Gly
      -15                               -10                               -5

TAT TAC CAC CCT TCC CAC CCC CAT TGG                                  129
Tyr Tyr His Pro Ser His Pro His Trp
      1                               5
```

(2) INFORMATION FOR SEQ ID NO: 180:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 base pairs

(B) TYPE: NUCLEIC ACID

(C) STRANDEDNESS: DOUBLE

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 36..77

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.9  
seq TLAVTLSALGATG/LF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 180:

```
AGAGGCCAGA TTMTGCAGGC CTGTGGGCTG ACACA ATG ACT TTG GCT GTT ACT      53
                               Met Thr Leu Ala Val Thr
                               -10

CTG AGT GCA TTG GGG GCC ACC GGA TTG TTT AAG GAG GCT TGT GAT CTA      101
```

## (2) INFORMATION FOR SEO ID NO: 181:

```
(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 330 base pairs
    (B) TYPE: NUCLEIC ACID
    (C) STRANDEDNESS: DOUBLE
    (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:
    (A) ORGANISM: Homo Sapiens
    (F) TISSUE TYPE: Uterus

(ix) FEATURE:
    (A) NAME/KEY: sig_peptide
    (B) LOCATION: 115..237
    (C) IDENTIFICATION METHOD: Von Heijne matrix
    (D) OTHER INFORMATION: score 3.9
                                seq CRCLITLPRSCR/ST

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 181:
```

[illegible]

## (2) INFORMATION FOR SEQ ID NO: 182:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 207 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 64..144
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq QLXLILVHFPAYS/VE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 182:

```

AAACTGCCAT CYGCAACTGA ACTTTGGCAG TAAACACAGC TTAGTTGTCT CAGAGGATTC      60
ACA ATG GGA AAT GTT TGT AGT TGC TGC CTC AGA GCA AGA TAT CAR CAG      108
  Met Gly Asn Val Cys Ser Cys Cys Leu Arg Ala Arg Tyr Gln Gln
    -25                      -20                      -15

TTG DCT TTA ATT TTA GTT CAT TTC CCA GCA TAT TCT GTT GAA GAT CAA      156
Leu Xaa Leu Ile Leu Val His Phe Pro Ala Tyr Ser Val Glu Asp Gln
    -10                      -5                      1

AGA GTG GAT CCT GGG GTG CCA GGG GAA TCC ACC GTC TGC CAC CAC AAT      204
Arg Val Asp Pro Gly Val Pro Gly Glu Ser Thr Val Cys His His Asn
   5                      10                      15                      20

CGG
Arg
   207

```

## (2) INFORMATION FOR SEQ ID NO: 183:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 130 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 8..70
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq PRCVISCIHGVWC/EE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 183:

```

ACTCTGC ATG CTT TAT GGC CTT GGC TCT GGG CCA AGG TGT GTG ATC TCC      49
  Met Leu Tyr Gly Leu Gly Ser Gly Pro Arg Cys Val Ile Ser
    -20                -15                -10

TGC ATT CAT GGT GTG TGG TGT GAG GAG GGG GAT GGG TCC CTG CCC CGT      97
  Cys Ile His Gly Val Trp Cys Glu Glu Gly Asp Gly Ser Leu Pro Arg
    -5                1                5

CTG CAC GTG GCC CTC ATG ATT CCC GCG CTA GGG      130
  Leu His Val Ala Leu Met Ile Pro Ala Leu Gly
    10                15                20

```

(2) INFORMATION FOR SEQ ID NO: 184:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 298 base pairs
  - (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: DOUBLE
  - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis

- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 62..187
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 3.8  
seq VTPLDSCPPSAHS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 184:

```

ACAGCTTCCA CTCTTGCTCTC CCTAAACCCCT GTTTTCCTCA CAGTAACTAG AATGTCCCTT      60

A ATG CAT AGA ATC ATG ACT CTC CTT CAT CTC AAA GCT CTC CAA CAG CTT      109
  Met His Arg Ile Met Thr Leu Leu His Leu Lys Ala Leu Gln Gln Leu
    -40                -35                -30

CAG AAT AAA ATC CAT GTC CCC AGG ATG CTC CCA GGG CCT GTG ACC CCT      157
  Gln Asn Lys Ile His Val Pro Arg Met Leu Pro Gly Pro Val Thr Pro
    -25                -20                -15

CTG GAC TCA TGC CCT CCT TCT GCT CAT TCT GCT CCA TCA CTG CTC ACT      205
  Leu Asp Ser Cys Pro Pro Ser Ala His Ser Ala Pro Ser Leu Leu Thr
    -10                -5                1                5

```

TCC CAG GTA CCC CTC CAA CAC ACC AAT GCG CCC CCA CCT CAC GGC CTC 253  
 Ser Gln Leu Pro Leu Gln His Thr Asn Ala Pro Pro Pro His Gly Leu  
                   10                  15                  20

TCC CTG CGC CGT GCC CTC CAC TGG ATT GCC CTT CCC TTG ATG GGG 298  
 Ser Leu Arg Arg Ala Leu His Trp Ile Ala Leu Pro Leu Met Gly  
                   25                  30                  35

## (2) INFORMATION FOR SEQ ID NO: 135:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 149 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 93..131
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq MLFLVLFYSAIFL/FT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 185:

AAACAACAAA AAAAAGTTTA AAAATTGGAA ACCACCAAAA GGTAGTATTA AAAGGGAAAT 60

AAAAATTACT CATAATCCCA GAACGCAGTC AT ATG CTA TTT TTA GTC TTA TTT 113  
                   Met Leu Phe Leu Val Leu Phe  
                                           -10

TAT TCA GCC ATT TTT CTC TTT ACA CTA ACT TTT TTT 149  
 Tyr Ser Ala Ile Phe Leu Phe Thr Leu Thr Phe Phe  
       -5                          1                          5

## (2) INFORMATION FOR SEQ ID NO: 186:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 180 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 133..174
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq VSLCVAALFPLQA/YG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 186:

```

AAAGAGTACC TGAAAACCTT AGAGAACCCT GGGGAAATAT TTATAGCCAG GCTTCTTGGA    60
GACTCTGGGA ACAGGAAAGT CAGGAACCCT GCCTTTCAGG AACTGCTGTA TCTCAGTCGM    120
MTTCTTCATT TC ATG GTT TCT CTC TGT GTA GCT GCT TTA TTT CCT CTT CAG    171
           Met Val Ser Leu Cys Val Ala Ala Leu Phe Pro Leu Gln
                        -10                               -5

GCT TAC GGG                                180
Ala Tyr Gly
  1

```

(2) INFORMATION FOR SEQ ID NO: 187:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 283 base pairs
  - (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: DOUBLE
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 218..268
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 3.7  
seq LFYIPSILTLLLA/CR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 187:

```

AAAATTCTTC CASAATGCTA ATGTAAATCT AATCAGCCTT TAGAATTTAA AGGCTTAAAA    60
AAGACTAAAG AAAAGTAACA ACCAAATGCA ATATGTAGAA CTTATATGGA GCCTGATTCC    120
AACATCAAGT ATAAAGAGAT ATTTTGTAGA AAATTGAGAA ATTTTAAAAC ATGAMATBAG    180
TATTATATGA TATTGAMGAC TGCTGCTTTT TCAMGAC ATG TCC TCA AAT TTA TTT    235
           Met Ser Ser Asn Leu Phe
                        -15

TAC ATT CCT TCC ATA CTA ACT CTT CTC CTT GCA TGT MGA CAG ACA GGG    283
Tyr Ile Pro Ser Ile Leu Thr Leu Leu Leu Ala Cys Arg Gln Thr Gly
-10                -5                1                5

```

## (2) INFORMATION FOR SEQ ID NO: 188:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 121 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 2..106
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq IKQFILCLGTCRG/EM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 188:

```

T ATG GGG CTT TTG AGA AAG TGT TTT CCC GTG ATG CTG GGG GGA AAC ACA   49
Met Gly Leu Leu Arg Lys Cys Phe Pro Val Met Leu Gly Gly Asn Thr
-35                      -30                      -25                      -20

CAT ATT CAA ATT ACT TGT ATA AAA CAG TTT ATT CTG TGT TTA GGA ACT   97
His Ile Gln Ile Thr Cys Ile Lys Gln Phe Ile Leu Cys Leu Gly Thr
-15                      -10                      -5

TGT AGG GGT GAA ATG CTG ACC AGG                                   121
Cys Arg Gly Glu Met Leu Thr Arg
      1                      5

```

## (2) INFORMATION FOR SEQ ID NO: 189:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 148 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 56..97
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq MLPLFCSPWESGG/RT



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 189:

```

TAAGCCGAGA AACTCCGTA CTGTGTTAAA AACTGTTTGA GGAACACTGG ATTAA ATG      58
   Met
ATG CTT CCA CTG TTC TSC TCT CCC TGG GAA AGC GGA GGC AGA ACG GTG      106
Met Leu Pro Leu Phe Cys Ser Pro Trp Glu Ser Gly Gly Arg Thr Val
      -10              -5              1
AAG CAG AGT GAA GGN YCT TGT TWA TTC CAG GCC CCC CAT GGG      149
Lys Gln Ser Glu Gly Xaa Cys Xaa Phe Gln Ala Pro His Gly
      5              10              15

```

(2) INFORMATION FOR SEQ ID NO: 190:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 27..71
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq KLLSDLVDSARC/KP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 190:

```

ATCTTAACAG AACTTTACAG ACTAGC ATG GCA AAG CTT CTC TCC GAT CTT AGT      53
   Met Ala Lys Leu Leu Ser Asp Leu Ser
   -15              -10
GTG GAC AGT GCT CGC TGC AAG CCT GGG AAT AAC CTT ACC AAA TCA CTC      101
Val Asp Ser Ala Arg Cys Lys Pro Gly Asn Asn Leu Thr Lys Ser Leu
      -5              1              5              10
TTG AAC ATT CAT GAT AAA CAA CTT CAA CAT GAC CCA CCG      140
Leu Asn Ile His Asp Lys Gln Leu Gln His Asp Pro Arg
      15              20

```

(2) INFORMATION FOR SEQ ID NO: 191:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 417 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 199..252  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.7  
seq VCWGHLLPARVST/RS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 191:

```

AGAGAYCAAC TCTATTTGAG CAASAGTKAG GAAGATTTCC CTGTCTCCCA GCTGAGTAAC   60
CACTCAGGTT TATTTAAATC CAGTTTAAAT ATGGTTTCAG TAATGATTTT CCAATGGTCT   120
ACAGCAAAGA ATGGTGCTCC AAGCCTGAAC ATTGAGCACG ACCCAGGTCA TATGCACAAC   180
ACGACAGGTT GAGCGTCC ATG TGT GGC TAC TGG GTT TGC TGG GGA CAC CTC   231
                Met Cys Gly Tyr Trp Val Cys Trp Gly His Leu
                -15                               -10

TTG CCT GCC AGG GTG AGC ACA CGC AGC AGT GAG CAG CCC CGT GTG ACC   279
Leu Pro Ala Arg Val Ser Thr Arg Ser Ser Glu Gln Pro Arg Val Thr
      -5                      1                      5

CCA CGG GAT GAG GAT GCC ATG ATG TCA GCA TCC CTT CTG ACT TGG AGG   327
Pro Arg Asp Glu Asp Ala Met Met Ser Ala Ser Leu Leu Thr Trp Arg
  10                15                20                25

TAT GTG ACA TTC ATG GTG CCA ATG CCA CTG TCA CCT TGC AGA TCA GTC   375
Tyr Val Thr Phe Met Val Pro Met Pro Leu Ser Pro Cys Arg Ser Val
          30                35                40

TGG GTT TGC TTC AGA CAG AAG ATC CTG GAA TAT GTT CAN GCA   417
Trp Val Cys Phe Arg Gln Lys Ile Leu Glu Tyr Val Xaa Ala
      45                50                55

```

(2) INFORMATION FOR SEQ ID NO: 192:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 167 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

-- (B) LOCATION: 66..137  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.7  
 seq AILGLSTFLNLLS/IN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 192:

```

AGCTGCGCAC AGATSATTGA ATTGCGGGGT TGCTGTAGGA ACCGCTGCTA TTGCCGCAGS   60
AGGAG ATG AAG TTA TCT TGT GCA GGC TGT GCA GAC ACA GGC ATT TTG GGA   110
    Met Lys Leu Ser Cys Ala Gly Cys Ala Asp Thr Ala Ile Leu Gly
                -20                -15                -10

CTC AGC ACT TTC CTT AAT TTA CTT TCC ATC AAC CTG CTC GGA ATG ATT   158
Leu Ser Thr Phe Leu Asn Leu Leu Ser Ile Asn Leu Leu Gly Met Ile
                -5                1                5

TCT TTC TCT                               167
Ser Phe Ser
                10
  
```

(2) INFORMATION FOR SEQ ID NO: 193:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 248 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (E) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 75..137  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.7  
 seq FSLGSCPAGPLSA/CV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 193:

```

ATGTCACATT TAASNAGAGG CCAGAGCTTG TCCAAATGG CTGTCCRWAM ACGACCCAC   60
ACTTGCGTTA GAAG ATG ATA CCT TTT TCA GGG ACA GTT TTC TCT CTT GGC   110
    Met Ile Pro Phe Ser Gly Thr Val Phe Ser Leu Gly
                -20                -15                -10

TCC TGT CCC GCT GGC CCT CTG TCT GCC TGT GTC CCT GAC CAT GGC TCC   158
Ser Cys Pro Ala Gly Pro Leu Ser Ala Cys Val Pro Asp His Gly Ser
                -5                1                5

CTG CAG TAC CCT TTA ACG ATT TAT CAG CAA GAC TGT KGA ACG CAT ARS   206
Leu Gln Tyr Pro Leu Thr Ile Tyr Gln Gln Asp Cys Xaa Thr His Xaa
                10                15                20
  
```

TGC CCA AGA TGC CTG TCC CTC CCC CTC CAG CAC CCC CGA CAG  
 Cys Pro Arg Cys Leu Ser Leu Pro Leu Gln His Pro Arg Gln  
 25 30 35

248

## (2) INFORMATION FOR SEQ ID NO: 194:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 360 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 70..174
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq PAVSLSAPAFASA/LR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 194:

AGAGGACAGG GTAGAGTCGC AGAAAGGAGA GACACACATA CATGGAAAGA GGAGCYTTCT 60  
 CCAATCTTA ATG ATT CCC AGC TCT CAG CCT CGT TTC TGM AAC CCA GCC TGC 111  
 Met Ile Pro Ser Ser Gln Pro Arg Phe Xaa Asn Pro Ala Cys  
 -35 -30 -25  
 AAG CAA ACT GTC CTG CTT WGG GAC CCT GCT GTG TCA CTC TCC GCA CCA 159  
 Lys Gln Thr Val Leu Leu Xaa Asp Pro Ala Val Ser Leu Ser Ala Pro  
 -20 -15 -10  
 GCC TTT GCC TCT GCT CTT CGC TCT ATG AMG TCC TCC CAG GCT GCA CGG 207  
 Ala Phe Ala Ser Ala Leu Arg Ser Met Xaa Ser Ser Gln Ala Ala Arg  
 -5 1 5 10  
 AAG GAC GAC TTT CTC AGG TCT CTT AGT GAT GCA GAC TCA GGG ACA TCA 255  
 Lys Asp Asp Phe Leu Arg Ser Leu Ser Asp Gly Asp Ser Gly Thr Ser  
 15 20 25  
 GAA CAC ATC TCA GCG GTG GTG ACT AGC CCT CGG ATT TCC TGC CAT GGT 303  
 Glu His Ile Ser Ala Val Val Thr Ser Pro Arg Ile Ser Cys His Gly  
 30 35 40  
 GCT GCC ATT CCC ACC GCC CGT GCC CTC TGC CTA YGC TGT TCC TGC TGC 351  
 Ala Ala Ile Pro Thr Ala Arg Ala Leu Cys Leu Xaa Cys Ser Cys Cys  
 45 50 55  
 ACC GAA CGC 360  
 Thr Glu Arg  
 60

## (2) INFORMATION FOR SEQ ID NO: 195:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 226 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 161..205
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
seq PTFLLISDSFLTQ/QP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 195:

```

ATTGGCCTGC TCTTCCTGAT ACCTACTTGG TCACTACTTA ATTACATTTT GTTTGTGTAT   60
CTTTTTTCTT CAGGCTGTAA ATTCTCTAAA GGCATTTTGC TTATTTTGGT GTCACAATTG  120
TTTAGGCCAT GCGCCTAGGT CTTCTTAAAA CACCTCTCTC ATG GCT CCT ACT TTT   175
                                   Met Ala Pro Thr Phe
                                   -15

CTA CTT ATT TCT GAT TCT TTT CTG ACT TCT CAG CCT TCT TTT TTT TTT   223
Leu Leu Ile Ser Asp Ser Phe Leu Thr Ser Gln Pro Ser Phe Phe Phe
-10                -5                1                5

TTT                               226
Phe

```

## (2) INFORMATION FOR SEQ ID NO: 196:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 362 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 219..275
- (C) IDENTIFICATION METHOD: Von Heijne matrix

-- (D) OTHER INFORMATION: score 3.6  
seq LSLGIIQWCLS/EN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 196:

```

AAAAAATACA CKGAGATTAT GTACGATTTA GTGATTGGT GGGATAATTA TAAATCGTGG    60
AATAATTTAT ATATGTGGAG TAAGAAGAGA GGGGTCAAAC CTTTGGTAC AAGCAACATC    120
TTGTTGCCAC CACCTTGATT TTCTCATAGG TGCTATTGTG TCCTAAGAGT RGRACAGRSR    180
RGRAACAAA GATAATTAAA CACAAGTCAG GTTACAAC ATG ATA TCT TTA ATT GTA    236
                               Met Ile Ser Leu Ile Val
                               -15

CTT TCT CTG CTT GGT ATC AAG ATT CAG TGG TGC TTG TCA GAA AAT ACC    284
Leu Ser Leu Leu Gly Ile Lys Ile Gln Trp Cys Leu Ser Glu Asn Thr
      -10                      -5                      1

TTG TTC TGT GAC TCT GAC TAT CTC TTG AGT CCC AAG GCT CCA ATT GAG    332
Leu Phe Cys Asp Ser Asp Tyr Leu Leu Ser Pro Lys Ala Pro Ile Glu
      5                      10                      15

CCT TTA TCT TTC AAC CTT ACC ACC CAG GGG    362
Pro Leu Ser Phe Asn Leu Thr Thr Gln Gly
      20                      25

```

(2) INFORMATION FOR SEQ ID NO: 197:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 263 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 129..257
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
seq LLYFNTFLPRKVA/RV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 197:

```

ATTAAGTCCT GCATTTTGTA AGAGGCAAAT GGAGASTAAC AGAAGAGTGT CTTTCTCCT    60
GGTTTTGGAG TCTTGCACTG GCCATGAGTG TTGKGACTGA TGGTCRACCC AGGCGGGCAT    120
TTTAATAA ATG GCC TGT GAT TCT TTT TTG AAA GAT GCT CTT CCA CAA GAG    170
      Met Ala Cys Asp Ser Phe Leu Lys Asp Ala Leu Pro Gln Glu
      -40                      -35                      -30

```

TTA AGC CAG TTA GRR TTC CTG TTT CCT CTT GTT GAT ATG AGA GAA GAT 218  
 Leu Ser Gln Leu Xaa Phe Leu Phe Pro Leu Val Asp Met Arg Glu Asp  
                   -25                                  -20                                  -15

CTC CTG TAT TTT AAC ACG TTT TTA CCA AGA AAG GTG GCA CGG GTG 263  
 Leu Leu Tyr Phe Asn Thr Phe Leu Pro Arg Lys Val Ala Arg Val  
                   -10                                  -5                                  1

## (2) INFORMATION FOR SEQ ID NO: 198:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 13..171
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
seq FLILHFFPQQIRK/KI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 198:

ATTCTCAATC AA ATG TTG CTA TTG AAT GAA AAC CTC AAA GCA GAA ATT CAA 51  
                   Met Leu Leu Leu Asn Glu Asn Leu Lys Ala Glu Ile Gln  
                                   -50                                  -45

AAG AAT GAA GCT CAA GGC AGC TGT ATC TTG TTT CTG TTT TGC TTT GAG 99  
 Lys Asn Glu Ala Gln Gly Ser Cys Ile Leu Phe Leu Phe Cys Phe Glu  
 -40                                  -35                                  -30                                  -25

AGT CAG AAT ATG CGA TCA AAA TCT ATC TTC CCC TTC CTT ATC CTT CAT 147  
 Ser Gln Asn Met Arg Ser Lys Ser Ile Phe Pro Phe Leu Ile Leu His  
                                   -20                                  -15                                  -10

TTT TTT CCC CAG CAG ATA AGA AAA AAA ATA GTC GTG CTT CTT TTA GGA 195  
 Phe Phe Pro Gln Gln Ile Arg Lys Lys Ile Val Val Leu Leu Leu Gly  
                                   -5                                  1                                  5

CTT AAT TCT CAG AAG GCA GGG 216  
 Leu Asn Ser Gln Lys Ala Gly  
                   10                                  15

## (2) INFORMATION FOR SEQ ID NO: 199:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 base pairs

- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 6..83
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq LLPFTFLSLKAFL/QX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 199:

```
AGCTG ATG ATA AGT AAG TAT GTG CAT TAT AGC TTG ACT GAC TTA CTA TTA   50
Met Ile Ser Lys Tyr Val His Tyr Ser Leu Thr Asp Leu Leu Leu
   -25                -20                -15

CCT TTT ACA TTC TTA AGC CTT AAA GCC TTT CTG CAG YYA AGA GTT TTA   98
Pro Phe Thr Phe Leu Ser Leu Lys Ala Phe Leu Gln Xaa Arg Val Leu
   -10                -5                1                5

ATG TCT CTT CCT CAA CAC AAG CCC TGG                               125
Met Ser Leu Pro Gln His Lys Pro Trp
               10
```

(2) INFORMATION FOR SEQ ID NO: 200:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 194 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 42..122
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq CSLLSSEFCALHFG/LK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 200:

```
AATATGTGAT CAAACGCCCA GGAGCCAGCT GGTGASAAAG A ATG GCG AGC ACA ATG   56
Met Ala Arg Thr Met
               -25
```



```

GGA GTT CCC AGA GCA TGC AAG GCC TTC TGT AGC CTC CTC TCC AGC TTC    104
Gly Val Pro Arg Ala Cys Lys Ala Phe Cys Ser Leu Leu Ser Ser Phe
      -20                      -15                      -10

TGT GCA TTA CAC TTT GGG CTC AAG AAA CAG TAT GGT ACT TCT TAC CTC    152
Cys Ala Leu His Phe Gly Leu Lys Lys Gln Tyr Gly Thr Ser Tyr Leu
      -5                      1                      5                      10

CAT GCC TGT GCT TAT GCT AGC CCC TTG ACC TGG GGT CCC TGG          194
His Ala Cys Ala Tyr Ala Ser Pro Leu Thr Trp Gly Pro Trp
      15                      20

```

## (2) INFORMATION FOR SEQ ID NO: 201:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 262..306
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq LCFLPHHRLQEA/RX

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 201:

```

ATTTCGCGGC GCTCGCBGMA CYHSGWTGTT CAGCACCTTC GGTCCGGTTG AGGTTGTCAA    60
GTCGGMCCAA ACAGGTTGTT TCTCTGCAGT TTCCAACATG GCAGGGMSGT TTAATAGACA    120
TGGATAAGAA GTCCACTCAC AGAAATCCTG AAGATGCCAG GGCTGGCAAA TATGAAGGTA    180
AACACAAACG AAAGAAAAGA AGAAAGCAAA ACCAAAACCA GCACCGATCC CGACATAGAT    240
CASTGACGTC TTTTCTTCA G ATG ATC CTA TGT TTC CTT CTT CCT CAT CAT    291
      Met Ile Leu Cys Phe Leu Leu Pro His His
      -15                      -10

CGT CTT CAG GAA GCC AGA YAG ATT CAA GTA TTG AAG ATK CTT CCA AGG    339
Arg Leu Gln Glu Ala Arg Xaa Ile Gln Val Leu Lys Xaa Leu Pro Arg
      -5                      1                      5                      10

GAA AAA TTA          348
Glu Lys Leu

```

## (2) INFORMATION FOR SEQ ID NO: 202:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 255 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 7..84
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq QCFFVCFSFKIYG/VI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 202:

```

AATAAC ATG CAA GAC TAC GTT TCA CAT GCA GTA CGG CGC CAC TGT CAG      48
    Met Gln Asp Tyr Val Ser His Ala Val Arg Arg His Cys Gln
      -25                -20                -15

TGT TTT TTT GTT TGT TTT TCC CCC AAG ATT TAT GGC GTA ATA ACA TGG      96
Cys Phe Phe Val Cys Phe Ser Pro Lys Ile Tyr Gly Val Ile Thr Trp
    -10                -5                1

ACC GTC CTG ATA ACT GGA GCC CGG GTT CTG TCA GAG CCC CAG AGG TTG      144
Thr Val Leu Ile Thr Gly Ala Arg Val Leu Ser Glu Pro Gln Arg Leu
    5                10                15                20

TGG GTT AGA CTT GAT GAC ATA ACA GCA AAT GCA GCG TGT GGT TAC AGA      192
Trp Val Arg Leu Asp Asp Ile Thr Ala Asn Ala Ala Cys Gly Tyr Arg
      25                30                35

AAG CAA GAG CCG CGG AAG ACG TTT GAA AAC AAT TGG GAA AAT TTG TAT      240
Lys Gln Glu Pro Arg Lys Thr Phe Glu Asn Asn Trp Glu Asn Leu Tyr
      40                45                50

ACG GAC TGG AAC TGG  255
Thr Asp Trp Asn Trp
      55

```

## (2) INFORMATION FOR SEQ ID NO: 203:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 224 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 120..212  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
 seq VLLNLALSHFNNC/GL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 203:

```

AGATYTTATA ATCTTGCTAC AAAGAAAGTA GGACAGTCTC AGCCTTTAAG AATGTCACTA   60
TAACAGTTTT TTTTTCCTT AAGGATATTT TAAACAGGAA AGTAGACAAC CGGGTAAGC   119
ATG GAG TTT GCT CAT GCT GCC GAA TGT GTG TCT TTT GCC CTA AAT GAA   167
Met Glu Phe Ala His Ala Ala Glu Cys Val Ser Phe Ala Leu Asn Glu
  -30                -25                -20
ACG CAC GTT CTT CTA AAT TTA GCC CTA TCA CAT TTT AAC AAT TGT GGC   215
Thr His Val Leu Leu Asn Leu Ala Leu Ser His Phe Asn Asn Cys Gly
  -15                -10                -5                1
CTC GCA GTG                               224
Leu Ala Val
  
```

(2) INFORMATION FOR SEQ ID NO: 204:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 276 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 133..222  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
 seq LLAASWLPRDAPC/EA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 204:

```

ACAGATTGCT TTCCAAGCTG AACATATGCA ACTGTATTGC TAAACTTACC AATTTCAAGG   60
AATCTGGGCG TCAAAAGCAT CCACATCCCT GCAGCAGGCC CCTGGGGAGG TAGGCAGGGT   120
GACAGCTGGG AA ATG GCR AAC CAG GGC TTT CCA TAC CTG TCT CCT TCT CTC   171
Met Gly Asn Gln Gly Phe Pro Tyr Leu Ser Pro Ser Leu
  -30                -25                -20
  
```

```

AGT GTC CAG GAT CTT CTT GCT GCT TCA TGG CTG CCC CGA GAT GCT CCC   219
Ser Val Gln Asp Leu Leu Ala Ala Ser Trp Leu Pro Arg Asp Ala Pro
      -15                      -10                      -5

TGT GAG GCC CCC CCG GGC CTG CCT TCA CAG ACA ATG CTC TGT GCC CCT   267
Cys Glu Ala Pro Pro Gly Leu Pro Ser Gln Thr Met Leu Cys Ala Pro
      1                      5                      10                      15

GGA CCA AGG   276
Gly Pro Arg

```

## (2) INFORMATION FOR SEQ ID NO: 205:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 196 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 68..133
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq AQLASPLLPGATP/VA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 205:

```

AACAAATTGA TCTGTGTGA TGAGTGTAAT AAAGCCTTCC ACCTGTTTGT TCTGAGGCCG   60

GCCCTCT ATG AAG TAC CAG ATG GTG AGT GGC AGT GCC CAG CTT GCC AGC   109
Met Lys Tyr Gln Met Val Ser Gly Ser Ala Gln Leu Ala Ser
      -20                      -15                      -10

CCG CTA CTG CCA GGC GCA ACT CCC GTG GCA GGA ACT ATA CTG AAG AGT   157
Pro Leu Leu Pro Gly Ala Thr Pro Val Ala Gly Thr Ile Leu Lys Ser
      -5                      1                      5

CTG CTT CTG AGG ACA GTG AAG ATG ATG AGA GTG TAT GGG   196
Leu Leu Leu Arg Thr Val Lys Met Met Arg Val Tyr Gly
      10                      15                      20

```

## (2) INFORMATION FOR SEQ ID NO: 206:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 145 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 55..94
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..40  
id AAL34726  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 99..121
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 34..66  
id AAL34726  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 72..140
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 1..69  
id R17226  
est

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 41..103
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 12.7  
seq ILFLLSWGPLQG/QQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 206:

```

AGTACGTTCC TTCTACTCTG GCACCACTCT CCAGGCTGCC ATG GGG CCC AGC ACC      55
                               Met Gly Pro Ser Thr
                               -20

CCT CTC CTC ATC TTG TTC CTT TTG TCA TGG TCG GGA CCC CTC CAA GGA      103
Pro Leu Leu Ile Leu Phe Leu Leu Ser Trp Ser Gly Pro Leu Gln Gly
-15                               -10                               -5

CAG CAG CAC CAC CTT GTG GAG TAC ATG GAA CGC CGA CAC GGG      145
Gln Gln His His Leu Val Glu Tyr Met Glu Arg Arg His Gly
1                               5                               10

```

(2) INFORMATION FOR SEQ ID NO: 207:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 172 base pairs

- (B) TYPE: NUCLEIC ACID
  - (C) STRANDEDNESS: DOUBLE
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Ovary
- (ix) FEATURE:
  - (A) NAME/KEY: other
  - (B) LOCATION: 73..169
  - (C) IDENTIFICATION METHOD: blastn
  - (D) OTHER INFORMATION: identity 100  
region 73..169  
id W25639  
est
- (ix) FEATURE:
  - (A) NAME/KEY: other
  - (B) LOCATION: 37..81
  - (C) IDENTIFICATION METHOD: blastn
  - (D) OTHER INFORMATION: identity 91  
region 38..82  
id W25639  
est
- (ix) FEATURE:
  - (A) NAME/KEY: other
  - (B) LOCATION: 42..169
  - (C) IDENTIFICATION METHOD: blastn
  - (D) OTHER INFORMATION: identity 100  
region 1..128  
id AA040016  
est
- (ix) FEATURE:
  - (A) NAME/KEY: other
  - (B) LOCATION: 34..169
  - (C) IDENTIFICATION METHOD: blastn
  - (D) OTHER INFORMATION: identity 96  
region 23..158  
id R72515  
est
- (ix) FEATURE:
  - (A) NAME/KEY: other
  - (B) LOCATION: 47..169
  - (C) IDENTIFICATION METHOD: blastn
  - (D) OTHER INFORMATION: identity 99  
region 1..123  
id T84313  
est
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 86..145
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 7.4  
seq LVFCVGLLTMAKA/ES

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 207:

```

AAGGGGGTCC AAGTGCTCA GCGGGGGG CACAGCAGGA CGTTTGSGGG CTTTCTTTCA . 60
GCAGG3GACA GCGGATTGG GGACA ATG GCG TCT CTT GGC CAC ATC TTG GTT      112
                               Met Ala Ser Leu Gly His Ile Leu Val
                               -20                               -15
TTC TGT GTG GGT CTC CTC ACC ATG GCC AAG GCA GAA AGT CCA AAG GAA      160
Phe Cys Val Gly Leu Leu Thr Met Ala Lys Ala Glu Ser Pro Lys Glu
   -10                               -5                               1                               5
CAC GAC CCG AGG  172
His Asp Pro Arg

```

## (2) INFORMATION FOR SEQ ID NO: 208:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 193 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 46..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 5..151  
id R14826  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 46..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 129..275  
id W55137  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 37..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 1..136  
id W64115  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 57..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 1..136  
id W75505  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 78..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 1..115  
id W20303  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 53..121
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.3  
seq ALSLLLVSGSLLP/GP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 208:

```
ACTCAATAAA TGTTTCCGC ATTAAGACGC TTCTTAGGAG TCTTCATGGA GG ATG TCG      58
                                     Met Ser

GGT TCG TCG CTG CCC AGC GCC CTG GCC CTC TCG CTG TTG CTG GTC TCT      106
Gly Ser Ser Leu Pro Ser Ala Leu Ala Leu Ser Leu Leu Leu Val Ser
-20                               -15                               -10

GGC TCC CTC CTC CCA GGC CCA GGC GCC GCT CAG AAC GAG CCA AGG ATT      154
Gly Ser Leu Leu Pro Gly Pro Gly Ala Ala Gln Asn Glu Pro Arg Ile
-5                               1                               5                               10

GTC ACC AGT GAA GAG GTC ATT ATT CGA GAC AGC CCC GTG                  193
Val Thr Ser Glu Glu Val Ile Ile Arg Asp Ser Pro Val
15                               20
```

## (2) INFORMATION FOR SEQ ID NO: 209:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 247 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 71..207



-- (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..137  
 id R73005  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 80..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..128  
 id N26942  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 86..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..122  
 id W02954  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 112..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..96  
 id T24907  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 137..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..71  
 id AA130938  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 53..223  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 7.1  
 seq VGLAVVSLGSGRG/SG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 209:

```

AACTGACAAG ACSTGGGSCCA AGAGGGGTCA CCCCCCCCCG ASCGGCGCGN AS  ATG  ATG    58
                                     Met  Met

GAA GTC GTA GTA GGA AAT GGC GTC GTG GCA TTG AGG GGC ATC CCT CCT    106
Glu Val Val Val Gly Asn Gly Val Val Ala Leu Arg Gly Ile Pro Pro
-55                -50                -45                -40

AGA ACC TCC AGG AAA AGC TCG CCG AAG ACS AGG TTC TCC GGA GAG AGA    154
Arg Thr Ser Arg Lys Ser Ser Arg Lys Thr Arg Phe Cys Gly Glu Arg

```

|                                                                 |     |     |     |     |
|-----------------------------------------------------------------|-----|-----|-----|-----|
|                                                                 | -35 | -30 | -25 |     |
| GGC TCC AAG CAG TCT GGG AAG TGT AGT CCA GTT GGC TTA GCA GTA GTT |     |     |     | 202 |
| Gly Ser Lys Gln Ser Gly Lys Cys Ser Pro Val Gly Leu Ala Val Val |     |     |     |     |
|                                                                 | -20 | -15 | -10 |     |
| TCG TTG GGG GGG AGC CGA GGT TCC GGG AAG GGG CTA GGC CGA CTG     |     |     |     | 247 |
| Ser Leu Gly Gly Ser Arg Gly Ser Gly Lys Gly Leu Gly Arg Leu     |     |     |     |     |
|                                                                 | -5  | 1   | 5   |     |

## (2) INFORMATION FOR SEQ ID NO: 210:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 373 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 252..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..124  
id AA081350  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 318..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..58  
id AA046671  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 200..247
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.7  
seq CFSLVLLLSIWT/TR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 210:

|                                                                   |     |
|-------------------------------------------------------------------|-----|
| AATTTTCCC CCAAGTACCT TGACAACTCA GAAGCTTGAA AGCAGGGAAA TCCGGATGTC  | 60  |
| TCGGTTATGA AGTGGAGCAG TGAGTGTGAG CCTCAACATA GTTCCAGAAC TCTCCATCCG | 120 |
| GACTAGTTAT TGAGCATCTG CCTCTCATAT CACCAGTGGC CATCTGAGGT GTTCCCTGG  | 180 |
| CTCTGAAGGG GTAGACAGC ATG GGC AGG TGC TTC AGC CTG GTG TTG CTT CTC  | 232 |

```

      Met Ala Arg Cys Phe Ser Leu Val Leu Leu Leu
      -15                               -10

ACT TCC ATC TGG ACC ACG AGG CTC CTG GTC CAA GGC TCT TTG CGT GCA   230
Thr Ser Ile Trp Thr Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala
  -5              1              5              10

GAA GAG CTT TCC ATC CAG GTG TCA TGC AGA ATK ATG GGG ATC ACC CTT   328
Glu Glu Leu Ser Ile Gln Val Ser Cys Arg Xaa Met Gly Ile Thr Leu
      15              20              25

GTG AGC AAA AAG GCG AAC CAG CAG CTG AAT TTC ACA GAA GCT AAG       373
Val Ser Lys Lys Ala Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys
      30              35              40

```

## (2) INFORMATION FOR SEQ ID NO: 211:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 438 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..355
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..207  
id R16604  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 354..407
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 207..260  
id R16604  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..352
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..214  
id N99553  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 380..429

(C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 237..285  
 id N99558  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 31..93  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.6  
 seq CLSCLLIPLALWS/II

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 211:

|                                                                  |     |
|------------------------------------------------------------------|-----|
| GAAATCTCCC GCAGTTCTAA GCAGGGCAAA ATG GGG TCT CGG AAG TGT GGA GGC | 54  |
| Met Gly Ser Arg Lys Cys Gly Gly                                  |     |
| -20 -15                                                          |     |
| TGC CTA AGT TGT TTG CTG ATT CCG CTT GCA CTT TGG AGT ATA ATC GTG  | 102 |
| Cys Leu Ser Cys Leu Leu Ile Pro Leu Ala Leu Trp Ser Ile Ile Val  |     |
| -10 -5 1                                                         |     |
| AAC ATA TTA TTG TAT TTC CCG AAT GGG CAA ACT TCC TAT GCA TCC AGC  | 150 |
| Asn Ile Leu Leu Tyr Phe Pro Asn Gly Gln Thr Ser Tyr Ala Ser Ser  |     |
| 5 10 15                                                          |     |
| AAT AAA CTC ACC AAC TAC GTG TGG TAT TTT GAA GGA ATC TGT TTC TCA  | 198 |
| Asn Lys Leu Thr Asn Tyr Val Trp Tyr Phe Glu Gly Ile Cys Phe Ser  |     |
| 20 25 30 35                                                      |     |
| GGC ATC ATG ATG CTT ATA GTA ACA ACA GTT CTT CTG GTA CTG GAG AAT  | 246 |
| Gly Ile Met Met Leu Ile Val Thr Thr Val Leu Leu Val Leu Glu Asn  |     |
| 40 45 50                                                         |     |
| AAT AAC AAC TAT AAA TGT TGC CAG AGT GAA AAC TGC AGC AAA AAA TAT  | 294 |
| Asn Asn Asn Tyr Lys Cys Cys Gln Ser Glu Asn Cys Ser Lys Lys Tyr  |     |
| 55 60 65                                                         |     |
| GTG ACA CTG CTG TCA ATT ATC TTT TCT TCC CTC GGA ATT GCT TTT TCT  | 342 |
| Val Thr Leu Leu Ser Ile Ile Phe Ser Ser Leu Gly Ile Ala Phe Ser  |     |
| 70 75 80                                                         |     |
| GGA TAC TGC CTG GTC ATC TCT GCC TTG GGT CTT GTC CAA GGG CCA TAT  | 390 |
| Gly Tyr Cys Leu Val Ile Ser Ala Leu Gly Leu Val Gln Gly Pro Tyr  |     |
| 85 90 95                                                         |     |
| TGC CGC ACC CTT GAT GGC TGG GAG TAT GCT TTT GAA GGC ACT GCT GGA  | 438 |
| Cys Arg Thr Leu Asp Gly Trp Glu Tyr Ala Phe Glu Gly Thr Ala Gly  |     |
| 100 105 110 115                                                  |     |

## (2) INFORMATION FOR SEQ ID NO: 212:

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 378 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 251..376  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 1..126  
id R16604  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 251..376  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 1..126  
id N99558  
est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 133..195  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.6  
seq CLSCLLIPLALWS/II

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 212:

```

ATTGTTCTC CAAACAGTAA ACCAGTATTT CACACTGAGA TTGTCGGCTG CGGGTATATT    60
CCAATCCCC GTCTCCTCAT GAATATGAAG TGAAGGGCTC TGAMCCTKGG AAGTGGTTCT    120
AAGCAGGGCA AA ATG GGG TCT CGG AAG TGT GGA GGC TGC CTA AGT TGT TTG    171
      Met Gly Ser Arg Lys Cys Gly Gly Cys Leu Ser Cys Leu
      -20              -15              -10

CTG ATT CCG CTT GCA CTT TGG AGT ATA ATC GTG AAC ATA TTA TTG TAT    219
Leu Ile Pro Leu Ala Leu Trp Ser Ile Ile Val Asn Ile Leu Leu Tyr
      -5              1              5

TTC CCG AAT GGG CAA ACT TCC TAT GCA TCC AGC AAT AAA CTC ACC AAC    267
Phe Pro Asn Gly Gln Thr Ser Tyr Ala Ser Ser Asn Lys Leu Thr Asn
      10              15              20

TAC GTG TGC TAT TTT GAA GGA ATC TGT TTC TCA GGC ATC ATG ATG CTT    315
Tyr Val Trp Tyr Phe Glu Gly Ile Cys Phe Ser Gly Ile Met Met Leu
      25              30              35              40

ATA GTA ACA ACA GTT CTT CTG GTA CTG GAG AAT AAT AAC AAC TAT AAA    363
Ile Val Thr Thr Val Leu Leu Val Leu Glu Asn Asn Asn Asn Tyr Lys
      45              50              55

TGT TGC CAG AAT GGG    378
Cys Cys Ser Ser Gly

```

60

## (2) INFORMATION FOR SEQ ID NO: 213:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 230 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 53..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 190..364  
id AA043641  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 92..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 1..136  
id N98697  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 69..102
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 393..426  
id AA147010  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 53..119
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 435..501  
id AA142584  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 159..209
- (C) IDENTIFICATION METHOD: Von Haefl matrix
- (D) OTHER INFORMATION: score 5.3  
seq ILFGVSFVFLTHC/TI

(xi). SEQUENCE DESCRIPTION: SEQ ID NO: 213:

```

AAACTTTAGC ACATCATTTT GGCATTCTAG AATATTTTCAT CTACCATACT ATAATTACCT    60
GAAGACATCA GGAGAAATACA AACTTGCAGG TGTTTTTCTT GGAGGTCGTT CAATGGGCTC    120
AAGAGCAGCT GCTTCTGTAA TGTGTACAT TGAGCCAG ATG ATG GTG ATG ATT TTG    176
                               Met Met Val Met Ile Leu
                               -15

TTC GGG GTC TCA TTT GTA TTT CTT ACC CAC TGC ACC ATC CAA AGC AGC    224
Phe Gly Val Ser Phe Val Phe Leu Thr His Cys Thr Ile Gln Ser Ser
   -10                -5                1                5

TGC GGG                                230
Cys Gly

```

(2) INFORMATION FOR SEQ ID NO: 214:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 394 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 310..393
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 1..84  
id HUM426A07B  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 293..349
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4  
seq VLVSLPHPHALT/CC

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 214:

```

AAACCTTGTT GCTAGGGACC GGGCGGTTTG CGSCAACCGT GGGCACTGCT GAATTTGAAT    60
TGAGGGGGCGA GGGAAAAGTT TTCCTCAGGT GTGGTGGGA GAGGGAGGCG GATGCCGGNG    120
AAACCGTAGG KACCGGTCA GAAAGCGGAC GCGCTCTCGG AGTTGGAAAG GGACGGCTGG    180
TTTCCCGGCA ACCGAACCGG GATGGGAAT GACTTCAATG AGATTGAACT TCAGCTGGAT    240

```

TGAAAGAGAG GCTAGAAGTT CCGCTTGCCA SCAGCCTCCT TASTAGAGCG GA ATG AGT 293  
Met Ser

AAT ACC CAC ACG GTG CTT GTC TCA CTT CCC CAT CCG CAC CCG GCC CTC 346  
Asn Thr His Thr Val Leu Val Ser Leu Pro His Pro His Pro Ala Leu  
-15 -10 -5

ACC TGC TGT CAC CTC GGC CWC CCA CAC CCG GTC CGC GGT CCC CGC CCG 394  
Thr Cys Cys His Leu Gly Xaa Pro His Pro Val Arg Ala Pro Arg Pro  
1 5 10 15

## (2) INFORMATION FOR SEQ ID NO: 215:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 473 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 111..321
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 1..211  
id N41784  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 143..237
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 5..99  
id T7C115  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 99..416
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1  
seq IITLACVPMTSFT/RN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 215:

ACAGCATCCT TTCGAAGGAT ADCTGAACAG AACCTTCTAA GTCTCAGACA CGTAAACCCA 60

AGTGTGGCRA AGGAACTCAT TGCTCTCGAA ATGCATAT ATG TKG GTT TAT ACA CTC 116  
Met Xaa Val Tyr Arg Leu  
-105



|                                                                 |     |
|-----------------------------------------------------------------|-----|
| CAA ACT CAA GAR AAG CCC AAC ACT ACT GTK CAA GTT CCA GCC TTT CTT | 164 |
| Gln Thr Gln Glu Lys Pro Asn Thr Thr Val Gln Val Pro Ala Phe Leu |     |
| -100 -95 -90 -85                                                |     |
| CAA GAG CTG GTA GAT CGG GAT AAT TCC AAA TTT GAG GAG TGG TGT ATT | 212 |
| Gln Glu Leu Val Asp Arg Asp Asn Ser Lys Phe Glu Glu Trp Cys Ile |     |
| -80 -75 -70                                                     |     |
| GAA ATG GCT GAG ATG CGT AAS AAA GTG TGG ATA AAG GAA AAG CAA AAC | 260 |
| Glu Met Ala Glu Met Arg Xaa Lys Val Trp Ile Lys Glu Lys Gln Asn |     |
| -65 -60 -55                                                     |     |
| ACG AAG AGG TTA AGG AGC TGT ACC AAA GGT TAC CTG CTG GAG CTG AGC | 308 |
| Thr Lys Arg Leu Arg Ser Cys Thr Lys Gly Tyr Leu Leu Glu Leu Ser |     |
| -50 -45 -40                                                     |     |
| CCT ATG AGT TTG TCT CTC TGG AAT GGC TGC AAA AGT GGT TGG ATG AAT | 356 |
| Pro Met Ser Leu Ser Leu Trp Asn Gly Cys Lys Ser Gly Trp Met Asn |     |
| -35 -30 -25                                                     |     |
| CAG CAA NTA CCA AAC CTA TTG ATA ATC ACG CTT GCC TGT GTT CCC ATG | 404 |
| Gln Gln Xaa Pro Asn Leu Leu Ile Ile Thr Leu Ala Cys Val Pro Met |     |
| -20 -15 -10 -5                                                  |     |
| ACA AGC TTC ACC CGG AAT AAA ATA TCA ATT ATG AAG AGG ATA TCT GAA | 452 |
| Thr Ser Phe Thr Arg Asn Lys Ile Ser Ile Met Lys Arg Ile Ser Glu |     |
| 1 5 10                                                          |     |
| TAT GCA GCK GAC ATT TTC TAT                                     | 473 |
| Tyr Ala Ala Asp Ile Phe Tyr                                     |     |
| 15                                                              |     |

## (2) INFORMATION FOR SEQ ID NO: 216:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 134 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 63..133
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 132..222  
id AA043974  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 99..133
- (C) IDENTIFICATION METHOD: blastn

.. (D) OTHER INFORMATION: identity 103  
region 1..36  
id WC5501  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 54..116  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.9  
seq LIAVVIIILLIFT/SV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 216:

```

AGACTTTGCT GATTAGCTT ATGGAAGAGG AACCAGAAAT TTGTCCTTGA ATA ATG      56
                                     Met
TTT CCC GTG TTG GGC TGG ATC TTG ATA GCA GTW GTY ATC ATC ATT CTT      104
Phe Pro Val Leu Gly Trp Ile Leu Ile Ala Val Val Ile Ile Ile Leu
-20          -15          -10          -5
CTG ATT TTT ACA TCT GTC ACC CGA TGC CTG      134
Leu Ile Phe Thr Ser Val Thr Arg Cys Leu
          1          5

```

## (2) INFORMATION FOR SEQ ID NO: 217:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 202 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 153..199  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..47  
id R14297  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 8..64  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.9  
seq SVCLC?CLNKGQS/EN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 217:

```

AAGCAAG ATG TTC TCC TGC TGT ATC TCA GTT TGT CTA TGT CCT TGT CTC      49
Met Phe Ser Cys Cys Ile Ser Val Cys Leu Cys Pro Cys Leu
          -15                      -10

AAC AAA GGC CAA AGT GAG AAT CTT TCC AGA GAC TGC GGW CAT TGG CTG      97
Asn Lys Gly Gln Ser Glu Asn Leu Ser Arg Asp Cys Gly His Trp Leu
-5              1              5              10

AAC CCT CAC CAT CGA CGC CTC TGG CCA TTT GGC AGA AGG CAC CCA CAG      145
Asn Pro His His Arg Arg Leu Trp Pro Phe Gly Arg Arg His Pro Gln
          15              20              25

GAT TGT GGA CTC TTC CAA GAT TCA CAA TGR TAT GGT GAA TCC AAA GAC      193
Asp Cys Gly Leu Phe Gln Asp Ser Gln Xaa Tyr Gly Glu Ser Lys Asp
          30              35              40

TGG AAC GGG                                202
Trp Asn Gly
          45

```

## (2) INFORMATION FOR SEQ ID NO: 218:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 406 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 38..103  
id W78795  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 405..470  
id AA151030  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 6..71  
id H48640

est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 338..403  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 98  
 region 73..138  
 id R99176  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 338..403  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 48..113  
 id W79571  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 143..229  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
 seq LTYLLLLSPIKYP/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 218:

```

ATCGTATTGG CACAGTTCTC TATGTAAGCA ATTTGAGAGG GAAGCAAAGG GGAAAAGTTT   60
GAGTTAGCTG TTCTCTGTCC TAGAATTTCCT CTGCATTAAT CTTGTCCTTG AAAATATATA 120
TAATACTGGT CCCTTAAACT CC ATG AGG CTT TGT CTC ATT ATG TAT TGT TCT   172
                Met Arg Leu Cys Leu Ile Met Tyr Cys Ser
                -25                               -20

TTT GGT ACC CTT TCC CAC TTA ACT TAC CTT TTG CTC CTA AGT CCT ATA   220
Phe Gly Thr Leu Ser His Leu Thr Tyr Leu Leu Leu Ser Pro Ile
                -15                               -10                               -5

AAA TAC CCC TTG GAT CTG GAT TTT TTA TAC CCG ATT TTC TCC ACT GTG   268
Lys Tyr Pro Leu Asp Leu Asp Phe Leu Tyr Pro Ile Phe Ser Thr Val
                1                               5                               10

TAT AAA AGG TAT ATT GTG ACT GTA AAT TTT TGT ATA TCA TGT TCT GAG   316
Tyr Lys Arg Tyr Ile Val Thr Val Asn Phe Cys Ile Ser Cys Ser Glu
                15                               20                               25

AGC TTC TTA CTT TCT GAT CTC ATA GCA CTA TTC CTG ATC AGA GAA CTC   364
Ser Phe Leu Leu Ser Asp Leu Ile Ala Leu Phe Leu Ile Arg Glu Leu
                30                               35                               40                               45

CAG TTG CTT CAA CAC ACA GTA TCA GTA GTG CAG CCA CCC ACG   406
Gln Leu Leu Gln His Thr Val Ser Val Val Gln Pro Pro Thr
                50                               55

```

## (ix) INFORMATION FOR SEQ ID NO: 219:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 210 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(118..206)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 144..232  
id T77881  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(64..118)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 231..285  
id T77881  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(126..206)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 139..219  
id R01713  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 70..147
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.5  
seq LLLALLLPVQVSS/FV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 219:

```

AACCAGCCAG GAGCCACCCA TCCTCCAGCA CACTGAGCAG CAAGCTGGAC ACACGGCACA   60
CTGATCCAA ATG GGT AAG GGG ATG GTG GCG ATG CTC ATT CTG GGT CTG CTA   111
      Met Gly Lys Gly Met Val Ala Met Leu Ile Leu Gly Leu Leu
      -25                -20                -15

GTT CTG GCG CTG CTC CTA CCC GTG CAG GTT TCT TCA TTT GTT CCT TTA   159
Leu Leu Ala Leu Leu Leu Pro Val Gln Val Ser Ser Phe Val Pro Leu
      -10                -5                1

AAG AGT ATG CCC GAA CCT ACT GCA GCG GAA ACC ACA AAG CCC TCC AAT   207

```

Thr Ser Met Pro Glu Ala Thr Ala Ala Glu Thr Thr Lys Pro Ser Asn  
 5 10 15 20

GGG  
 Gly

210

## (2) INFORMATION FOR SEQ ID NO: 220:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 189 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 2..170
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
 region 4..172  
 id H56777  
 est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 25..87
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.5  
 seq LLVLFVLLANVQG/PG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 220:

AAGAAAGGCT GGCCTCTCTT CAAC ATG GGA TCT TCT GGA CTT TTG AGC CTC 51  
 Met Gly Ser Ser Gly Leu Leu Ser Leu  
 -20 -15

CTG GTG CTA TTC GTC CTC TTA GCG AAT GTC CAG GGA CCT GGT CTC ACT 99  
 Leu Val Leu Phe Val Leu Leu Ala Asn Val Gln Gly Pro Gly Leu Thr  
 -10 -5 1

GAT TGG TTA TTT CCC AGG AGA TGT CCC AAA ATC AGA GAA GAA TGT GAA 147  
 Asp Trp Leu Phe Pro Arg Arg Cys Pro Lys Ile Arg Glu Glu Cys Glu  
 5 10 15 20

TTC CAA GAA AGG GAT GTG TGT ACA AAG GAC AGA CAA TGC CGA 189  
 Phe Gln Glu Arg Asp Val Cys Thr Lys Asp Arg Gln Cys Arg  
 25 30

## (2) INFORMATION FOR SEQ ID NO: 221:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 323 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 52..253
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 10..216  
id R60167  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 235..319
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 194..278  
id R60167  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 143..258
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 90..205  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 55..145
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..91  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 235..316
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 183..264  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 141..258
- (C) IDENTIFICATION METHOD: blastn

171

.. (D) OTHER INFORMATION: identity 91  
region 85..202  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 55..144  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..89  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 235..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 180..264  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 58..257  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 1..200  
id AA039912  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 248..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 190..261  
id AA039912  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 50..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..270  
id R54127  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 90..194  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 8.3  
seq NLLLLHCVSRSHS/QN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 221:

AGGAAAGGCG TGACCACGCA GTTCTTGGGT CTGTCTCTCT GGCTGGGGT TGTGTTGAG 60



```

GCCGTGTCTC CGCTCCTGTG CCCGGGAAG ATG STG CTA GGT GGT TGC CCG GTT      113
                        Met Val Leu Gly Gly Cys Pro Val
                        -35                               -30

AGT TAC TTA CTT CTG TGC GGC CAG GCG GCT TTG CTG CTG GGG AAT TTA      161
Ser Tyr Leu Leu Leu Cys Gly Gln Ala Ala Leu Leu Leu Gly Asn Leu
      -25                               -20                               -15

CTT CTG CTG CAT TGT GTG TCT CGG AGC CAC TCG CAA AAT GCG ACC GCT      209
Leu Leu Leu His Cys Val Ser Arg Ser His Ser Gln Asn Ala Thr Ala
      -10                               -5                               1                               5

GAG CCT GAG CTC ACA TCC GCT GGC GCC CCC AGC CGG AGG GCC CCG GGG      257
Glu Pro Glu Leu Thr Ser Ala Gly Ala Pro Ser Arg Arg Ala Pro Gly
                        10                               15                               20

GTG CTG CGA GGT GGG AAT ATG GCG ACC CCC ACT CTC CGG TCA TCC TCT      305
Val Leu Arg Ala Gly Asn Met Ala Thr Pro Thr Leu Arg Ser Ser Ser
                        25                               30                               35

GCT CTT ACC TAC CTT GGG      323
Ala Leu Thr Tyr Leu Gly
      40

```

## (2) INFORMATION FOR SEQ ID NO: 222:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 165 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 31..143
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 271..383  
id W16767  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 34..87
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.3  
seq LLSLSSLPLVLLG/WE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 222:

```

AACGAGACTT CTGACCCCTT GGGCAACAGC CAG ATG GAG ACT GGT CGC GTT TTG      54

```

Met Glu Thr Gly Arg Leu Leu  
-15

AGC CTC AGC TCT CTT CCT CTT GTT CTC CTA GGG TGG GAG TAC AGC AGC 102  
Ser Leu Ser Ser Leu Pro Leu Val Leu Leu Gly Trp Glu Tyr Ser Ser  
-10 -5 1 5

CAA ACG CTG AAC TTA GTC CCA TCC ACT TCC ATC TTA TCC TTT GTG CCC 150  
Gln Thr Leu Asn Leu Val Pro Ser Thr Ser Ile Leu Ser Phe Val Pro  
10 15 20

TTC ATC CCC CGA GTG 165  
Phe Ile Pro Arg Val  
25

## (2) INFORMATION FOR SEQ ID NO: 223:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 201 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 24..203
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 1..180  
id HSC1PF091  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 114..203
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 73..167  
id H03709  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 35..107
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..73  
id H03709  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 25..81

.. (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8.2  
 seq QVLALVLVAALWG/GT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 223:

```

AAAGTAGAAG ACAGCGGCGT TGCC ATG GCG GCG TCT CTG GGG CAG GTG TTG      51
                Met Ala Ala Ser Leu Gly Gln Val Leu
                        -15

GCT CTG GTG CTG GTG GCC GCT CTG TGG GGT GGC ACG CAG CCG CTG CTG      99
Ala Leu Val Leu Val Ala Ala Leu Trp Gly Gly Thr Gln Pro Leu Leu
-10                -5                1                5

AAG CGG GCC TCC GCC GGC CTG CAG CGG GTT CAT GAG CCG ACC TGG GCC      147
Lys Arg Ala Ser Ala Gly Leu Gln Arg Val His Glu Pro Thr Trp Ala
                10                15                20

CAG CAG TTG CTA CAG GAG ATG AAG ACC CTC TTC TTG AAT ACT GAG TAC      195
Gln Gln Leu Leu Gln Glu Met Lys Thr Leu Phe Leu Asn Thr Glu Tyr
                25                30                35

CTG ATG  201
Leu Met
40
  
```

(2) INFORMATION FOR SEQ ID NO: 224:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 462 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 6..119
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 1..114  
id N83684  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 147..241
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 112..206  
id N83684  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 283..323  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 245..285  
id N83684  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 327..361  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 287..321  
id N83684  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 150..299  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 177..326  
id H94179  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 23..109  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 41..127  
id H94179  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 17..90  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 1..74  
id AA093069  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 144..194  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 123..173  
id AA093069  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 150..249  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 193..297  
id T67190

|                                                                    |     |
|--------------------------------------------------------------------|-----|
| ATTGTGTTCTC CAAACAGTAA ACCAGTATTT CACACTGAGA TTGTCGGCTG CGGGTATATT | 60  |
| CCAAATTCCCC GTCTCCTCAT GAATATGAAG TGAAGGGCTC TGAMCCTKGG AAGTGGTTCT | 120 |
| AAGCAGGGCA AA ATG GGG TCT CGG AAG TGT GGA GGC TGC CTA AGT TGT TTG  | 171 |
| Met Gly Ser Arg Lys Cys Gly Gly Cys Leu Ser Cys Leu                |     |
| -20 -15 -10                                                        |     |
| CTG ATT CCG CTT GCA CTT TGG AGT ATA ATC GTG AAC ATA TTA TTG TAT    | 219 |
| Leu Ile Pro Leu Ala Leu Trp Ser Ile Ile Val Asn Ile Leu Leu Tyr    |     |
| -5 1 5                                                             |     |
| TTC CCG AAT GGG CAA ACT TCC TAT GCA TCC ACC AAT AAA CTC ACC AAC    | 267 |
| Phe Pro Asn Gly Gln Thr Ser Tyr Ala Ser Ser Asn Lys Leu Thr Asn    |     |
| 10 15 20                                                           |     |
| TAC GTG TGG TAT TTT GAA GGA ATC TGT TTC TCA GGC ATC ATG ATG CTT    | 315 |
| Tyr Val Trp Tyr Phe Glu Gly Ile Cys Phe Ser Gly Ile Met Met Leu    |     |
| 25 30 35 40                                                        |     |
| ATA GTA ACA ACA GTT CTT CTG GTA CTG GAG AAT AAT AAC AAC TAT AAA    | 363 |
| Ile Val Thr Thr Val Leu Leu Val Leu Glu Asn Asn Asn Asn Tyr Lys    |     |
| 45 50 55                                                           |     |
| TGT TGC CAC AGT GGG                                                | 378 |
| Cys Cys Ser Ser Gly                                                |     |

(C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 237..265  
 id N99558  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 31..93  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.6  
 seq CLSCLLIPLALWS/II

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 211:

|                                                                  |     |
|------------------------------------------------------------------|-----|
| GAAATCTCCC GCAGTTCTAA GCAGGGCAAA ATG GGG TCT CGG AAG TGT GGA GGC | 54  |
| Met Gly Ser Arg Lys Cys Gly Gly                                  |     |
| -20 -15                                                          |     |
| TGC CTA AGT TGT TTG CTG ATT CCG CTT GCA CTT TGG AGT ATA ATC GTG  | 102 |
| Cys Leu Ser Cys Leu Leu Ile Pro Leu Ala Leu Trp Ser Ile Ile Val  |     |
| -10 -5 1                                                         |     |
| AAC ATA TTA TTG TAT TTC CCG AAT GGG CAA ACT TCC TAT GCA TCC AGC  | 150 |
| Asn Ile Leu Leu Tyr Phe Pro Asn Gly Gln Thr Ser Tyr Ala Ser Ser  |     |
| 5 10 15                                                          |     |
| AAT AAA CTC ACC AAC TAC GTG TGG TAT TTT GAA GGA ATC TGT TTC TCA  | 198 |
| Asn Lys Leu Thr Asn Tyr Val Trp Tyr Phe Glu Gly Ile Cys Phe Ser  |     |
| 20 25 30 35                                                      |     |
| GGC ATC ATG ATG CTT ATA GTA ACA ACA GTT CTT CTG GTA CTG GAG AAT  | 246 |
| Gly Ile Met Met Leu Ile Val Thr Thr Val Leu Leu Val Leu Glu Asn  |     |
| 40 45 50                                                         |     |
| AAT AAC AAC TAT AAA TGT TGC CAG AGT GAA AAC TGC AGC AAA AAA TAT  | 294 |
| Asn Asn Asn Tyr Lys Cys Cys Gln Ser Glu Asn Cys Ser Lys Lys Tyr  |     |
| 55 60 65                                                         |     |
| GTG ACA CTG CTG TCA ATT ATC TTT TCT TCC CTC GGA ATT GCT TTT TCT  | 342 |
| Val Thr Leu Leu Ser Ile Ile Phe Ser Ser Leu Gly Ile Ala Phe Ser  |     |
| 70 75 80                                                         |     |
| GGA TAC TGC CTG GTC ATC TCT GCC TTG GGT CTT GTC CAA GGG CCA TAT  | 390 |
| Gly Tyr Cys Leu Val Ile Ser Ala Leu Gly Leu Val Gln Gly Pro Tyr  |     |
| 85 90 95                                                         |     |
| TGC CGC ACC CTT GAT GGC TGG GAG TAT GCT TTT GAA GGC ACT GCT GGA  | 438 |
| Cys Arg Thr Leu Asp Gly Trp Glu Tyr Ala Phe Glu Gly Thr Ala Gly  |     |
| 100 105 110 115                                                  |     |

## (12) INFORMATION FOR SEQ ID NO: 212:

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 378 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

```

Met Ala Arg Cys Phe Ser Leu Val Leu Leu Leu
-15 -10

ACT TCC ATC TGG ACC ACG AGG CTC CTG GTC CAA GGC TCT TTG CGT GCA 230
Thr Ser Ile Trp Thr Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala
-5 1 5 10

GAA GAG CTT TCC ATC CAG GTG TCA TGC AGA ATK ATG GGG ATC ACC CTT 328
Glu Glu Leu Ser Ile Gln Val Ser Cys Arg Xaa Met Gly Ile Thr Leu
15 20 25

GTG AGC AAA AAG GCG AAC CAG CAG CTG AAT TTC ACA GAA GCT AAG 373
Val Ser Lys Lys Ala Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys
30 35 40

```

## (2) INFORMATION FOR SEQ ID NO: 211:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 438 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..355
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..207  
id R16604  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 354..407
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 207..260  
id R16604  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 149..352
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..214  
id N99555  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 350..429

```

          -35              -30              -25
GGC TCC AAG CAG TCT GGG AAG TGT AGT CCA GTT GGC TTA GCA GTA GTT      202
Gly Ser Lys Gln Ser Gly Lys Cys Ser Pro Val Gly Leu Ala Val Val
          -20              -15              -10
TCG TTG GGG GGG AGC CGA GGT TCC GGG AAG GGG CTA GGC CGA CTG      247
Ser Leu Gly Gly Ser Arg Gly Ser Gly Lys Gly Leu Gly Arg Leu
          -5              1              5

```

## (2) INFORMATION FOR SEQ ID NO: 210:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 373 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 252..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..124  
id AA081350  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 319..375
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 1..59  
id AA046671  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 200..247
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.7  
seq CFSLVLLLSIWT/TR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 210:

```

AATTTTCCC CCACTGAGCT TGACAAGTCA GAAGCTTGAA ASCAGGGAAA TCCGGATGTC      60
TCGGTTATGA AGTGGAGCAG TGAGTGTGAC CCTCAACATA GTTCCAGAAC TCTCCATCCG      120
GACTAGTTAT TGACATCTG CTTCTCATAT CACCAGTGSC CATCTGAGGT GTTCCCTGG      180
CTCTGAAGGG GTAAGGAGG ATG GGC AGG TGC TTC AGC CTA GTG TTG CTT CTC      232

```



(C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..137  
 id R73005  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 80..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..128  
 id N26942  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 86..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..122  
 id WU2954  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 112..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..96  
 id T24907  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 137..207  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 1..71  
 id PA130938  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 53..223  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 7.1  
 seq VGLAVVSLGGSRG/SG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 209:

```

AACTGACAAG ACGTGGGGCCA AGAGGGGTCA CCCCCCCCCG ACCGGCGCGN AS ATG ATG   58
                                     Met Met

GAA GTC GTA GTA GGA AAT GGC GTC GTG GCA TTG AGG GGC ATC CCT CCT   106
Glu Val Val Val Gly Asn Gly Val Val Ala Leu Arg Gly Ile Pro Pro
-55                      -50                      -45                      -40

AGA ACC TCC AGG AAA AGC TCG CCG AAG ACC AGC TTC TGC GGA GAG AGA   154
Arg Thr Ser Arg Lys Ser Ser Arg Lys Thr Arg Phe Cys Gly Glu Arg

```

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 207:

```

AAGGGGGTCC AAAGTGCTCA GCGGGGGGG CACAGCAGGA CGTTTG3GGG CTTTCTTTCA   60
GCAGGGGACA GCGCGATTGG GGACA ATG GCG TCT CTT GGC CAC ATC TTG GTT   112
                               Met Ala Ser Leu Gly His Ile Leu Val
                               -20                               -15

TTC TGT GTG GGT CTC CTC ACC ATG GCC AAG GCA GAA AGT CCA AAG GAA   160
Phe Cys Val Gly Leu Leu Thr Met Ala Lys Ala Glu Ser Pro Lys Glu
-10                               -5                               1                               5

CAC GAC CCG AGG   172
His Asp Pro Arg

```

## (2) INFORMATION FOR SEQ ID NO: 208:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 193 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 46..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 5..151  
id R14926  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 46..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 129..275  
id W55137  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 57..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 1..136  
id W64115  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 57..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 1..136  
id W75505  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 78..192
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 1..115  
id W20303  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 53..121
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.3  
seq ALSLLLVSGSLLP/GP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 208:

```

ACTCAATAAA TGTTCCTCCG ATTAAGACGC TTCTTAGGAG TCTTCATGGA GG ATG TCG      58
                                     Met Ser

GGT TCG TCG CTG CCC AGC GCC CTG GCC CTC TCG CTG TTG CTG GTC TCT      106
Gly Ser Ser Leu Pro Ser Ala Leu Ala Leu Ser Leu Leu Leu Val Ser
-20                               -15                               -10

GGC TCC CTC CTC CCA GGC CCA GGC GCC GCT CAG AAC GAG CCA AGG ATT      154
Gly Ser Leu Leu Pro Gly Pro Gly Ala Ala Gln Asn Glu Pro Arg Ile
-5                               1                               5                               10

GTC ACC AGT GAA GAG GTC ATT ATT CGA GAC AGC CCC GTG                  193
Val Thr Ser Glu Glu Val Ile Ile Arg Asn Ser Pro Val
                               15                               20

```

## (2) INFORMATION FOR SEQ ID NO: 209:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 247 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 71..207

60

## (2) INFORMATION FOR SEQ ID NO: 213:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 230 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 53..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 190..364  
id AA043641  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 92..227
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 1..136  
id N98697  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 69..102
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 393..426  
id AA147010  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 53..119
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 435..501  
id AA142584  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 159..209
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq ILFGVSFVFLTHC/TI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 213:

```

AAACTTTASC ACATCATTTT GSCATTCTAG AATATTTTCAT CTACCATACT ATAATTACCT   60
GAAGACATCA GGAGAATACA AACTTGCACG TGTITTTTCTT GGAGGTCGTT CAATGGGCTC   120
AAGAGCAGCT GCTTCTGTAA TGTGTACAT TGAGCCAG ATG ATG GTG ATG ATT TTG   176
                               Met Met Val Met Ile Leu
                               -15

TTC GGG GTC TCA TTT GTA TTT CTT ACC CAC TGC ACC ATC CAA AGC AGC   224
Phe Gly Val Ser Phe Val Phe Leu Thr His Cys Thr Ile Gln Ser Ser
   -10               -5               1               5

TGC GGG                               230
Cys Gly

```

(2) INFORMATION FOR SEQ ID NO: 214:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 394 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 310..393
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 1..84  
id HUM426A07B  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 293..349
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4  
seq VLVSLPHPHALT/CC

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 214:

```

AAACCTTGTT GCTAGGGACC GGGCGGTTTG CGSCAACCGT GGGCACTGCT GAATTTGAAT   60
TGAGGGGGCA GGGAAAAGTT TTCCTCAGGT GTGGTGGGGA GAGGGAGGCG GATGCCGCG   120
AAACCGTAGG KACCGGCTCA GAAAGCGGAC CGCCTCTCGG AGTTGGAAAG GGACGCGTGG   180
TTTCCCGGCA AGCGAACCGG GATGGGAAGT GACTTCAGTG AGATTGAACT TCAGCTGGAT   240

```

TGAAAGAGAG GCTAGAAGTT CCGCTTGCCA SCAGCCTCCT TAGTAGAGCG GA ATG AGT 293  
Met Ser

AAT ACC CAC ACG GTG CTT GTC TCA CTT CCC CAT CCG CAC CCG GCC CTC 346  
Asn Thr His Thr Val Leu Val Ser Leu Pro His Pro His Pro Ala Leu  
-15 -10 -5

ACC TGC TGT CAC CTC GGC CWC CCA CAC CCG GTC CGC GGT CCC CGC CCG 394  
Thr Cys Cys His Leu Gly Xaa Pro His Pro Val Arg Ala Pro Arg Pro  
1 5 10 15

## (2) INFORMATION FOR SEQ ID NO: 215:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 473 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 111..321
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 1..211  
id N41784  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 143..237
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 5..99  
id T70115  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 99..416
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1  
seq IITLACVPMTSFT/RN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 215:

ACAGCATCCT TTCAAGGAT ADCTGAACAG AACCTTCTAA GTCTCAGACA CGTAAACCCA 60

AGTGTGGCAA AGGAATCAT TGCTCTCGAA ATGCATAT ATG TKG GTT TAT ACA CTC 116  
Met Xaa Val Tyr Arg Leu  
-105

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| CAG ACT CAA GAG AAG CCC AAC ACT ACT GTG CAA GTT CCA GCC TTT CTT | 164 |
| Gln Thr Gln Glu Lys Pro Asn Thr Thr Val Gln Val Pro Ala Phe Leu |     |
| -100 -95 -90 -85                                                |     |
| CAG GAG CTG GTA GAT CGG GAT AAT TCC AAA TTT GAG GAG TGG TGT ATT | 212 |
| Gln Glu Leu Val Asp Arg Asp Asn Ser Lys Phe Glu Glu Trp Cys Ile |     |
| -80 -75 -70                                                     |     |
| GAA ATG GCT GAG ATG CGT AAS AAA GTG TGG ATA AAG GAA AAG CAA AAC | 260 |
| Glu Met Ala Glu Met Arg Xaa Lys Val Trp Ile Lys Glu Lys Gln Asn |     |
| -65 -60 -55                                                     |     |
| ACG AAG AGG TTA AGG AGC TGT ACC AAA GGT TAC CTG CTG GAG CTG AGC | 308 |
| Thr Lys Arg Leu Arg Ser Cys Thr Lys Gly Tyr Leu Leu Glu Leu Ser |     |
| -50 -45 -40                                                     |     |
| CCT ATG AGT TTG TCT CTC TGG AAT GGC TGC AAA AGT GGT TGG ATG AAT | 356 |
| Pro Met Ser Leu Ser Leu Trp Asn Gly Cys Lys Ser Gly Trp Met Asn |     |
| -35 -30 -25                                                     |     |
| CAG CAA NTA CCA AAC CTA TTG ATA ATC ACG CTT GCC TGT GTT CCC ATG | 404 |
| Gln Gln Xaa Pro Asn Leu Leu Ile Ile Thr Leu Ala Cys Val Pro Met |     |
| -20 -15 -10 -5                                                  |     |
| ACA AGC TTC ACC CGG AAT AAA ATA TCA ATT ATG AAG AGG ATA TCT GAA | 452 |
| Thr Ser Phe Thr Arg Asn Lys Ile Ser Ile Met Lys Arg Ile Ser Glu |     |
| 1 5 10                                                          |     |
| TAT GCA GCK GAC ATT TTC TAT                                     | 473 |
| Tyr Ala Ala Asp Ile Phe Tyr                                     |     |
| 15                                                              |     |

## (2) INFORMATION FOR SEQ ID NO: 216:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 134 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 63..133
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 152..222  
id AAD43974  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 99..133
- (C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 100  
region 1..36  
id WC5501  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 54..116  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.9  
seq LIAVVIIILLIFT/SV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 216:

```

AGACTTTGCT GATTTAGCTT ATGGAAGAGG AACCAGAAAT TTGTCCTTGA ATA ATG      56
   Met
TTT CCC GTG TTG GGC TGG ATC TTG ATA GCA GTW GTY ATC ATC ATT CTT      104
Phe Pro Val Leu Gly Trp Ile Leu Ile Ala Val Val Ile Ile Ile Leu
-20                               -15                               -10                               -5

CTG ATT TTT ACA TCT GTC ACC CGA TGC CTG                                134
Leu Ile Phe Thr Ser Val Thr Arg Cys Leu
      1                               5

```

## (2) INFORMATION FOR SEQ ID NO: 217:

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 202 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(E) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 153..199  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..47  
id R14297  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 8..64  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.9  
seq SVCLCPCLNKGQS/EN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 217:



```

AAGCAAG ATG TTC TGC TGT ATC TCA GTT TGT CTA TGT CCT TGT CTC      49
Met Phe Ser Cys Cys Ile Ser Val Cys Leu Cys Pro Cys Leu
      -15                      -10

AAC AAA GGC CAA AGT GAG AAT CTT TCC AGA GAC TGC GGW CAT TGG CTG      97
Asn Lys Gly Gln Ser Glu Asn Leu Ser Arg Asp Cys Gly His Trp Leu
      -5                      1                      5                      10

AAC CCT CAC CAT CGA CSC CTC TGG CCA TTT GGC AGA AGG CAC CCA CAG      145
Asn Pro His His Arg Arg Leu Trp Pro Phe Gly Arg Arg His Pro Gln
      15                      20                      25

GAT TGT GGA CTC TTC CAA GAT TCA CAA TGR TAT GGT GAA TCC AAA GAC      193
Asp Cys Gly Leu Phe Gln Asp Ser Gln Xaa Tyr Gly Glu Ser Lys Asp
      30                      35                      40

TGG AAC GGG  202
Trp Asn Gly
      45

```

## (2) INFORMATION FOR SEQ ID NO: 218:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 406 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 38..103  
id W78795  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 405..470  
id AA151030  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 338..403
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 6..71  
id H48640

est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 338..403  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 98  
                           region 73..138  
                           id R99176  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 338..403  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
                           region 48..113  
                           id W79571  
                           est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 143..229  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
                           seq LTYLLLLSPIKYP/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 218:

```

ATCGTATTGG CACAGTTCTC TATGTAAGCA ATTTGAGAGG GAAGCAAAGG GGAAAAGTTT   60
GAGTTAGCTG TTCTCTGTCC TAGAATTTCC CTGCATTAAT CTTGTCCTTG AAAATATATA   120
TAATACTGGT CCCTTAAACT CC ATG AGG CTT TGT CTC ATT ATG TAT TGT TCT   172
                Met Arg Leu Cys Leu Ile Met Tyr Cys Ser
                -25                               -20

TTT GGT ACC CTT TCC CAC TTA ACT TAC CTT TTG CTC CTA AGT CCT ATA   220
Phe Gly Thr Leu Ser His Leu Thr Tyr Leu Leu Leu Leu Ser Pro Ile
                -15                               -10                               -5

AAA TAC CCC TTG GAT CTG GAT TTT TTA TAC CCG ATT TTC TCC ACT GTG   268
Lys Tyr Pro Leu Asp Leu Asp Phe Leu Tyr Pro Ile Phe Ser Thr Val
                1                               5                               10

TAT AAA AGG TAT ATT GTG ACT GTA AAT TTT TGT ATA TCA TGT TCT GAG   316
Tyr Lys Arg Tyr Ile Val Thr Val Asn Phe Cys Ile Ser Cys Ser Glu
                15                               20                               25

AGC TTC TTA CTT TCT GAT CTC ATA GCA CTA TTC CTG ATC AGA GAA CTC   364
Ser Phe Leu Leu Ser Asp Leu Ile Ala Leu Phe Leu Ile Arg Glu Leu
                30                               35                               40                               45

CAG TTG CTT CAA CAC ACA GTA TCA GTA GTG CAG CCA CCC ACG   406
Gln Leu Leu Gln His Thr Val Ser Val Val Gln Pro Pro Thr
                50                               55

```

## (ix) INFORMATION FOR SEQ ID NO: 219:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 210 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: CDNA
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis
- (ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(118..206)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
                           region 144..232  
                           id T77881  
                           est
- (ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(64..118)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 98  
                           region 231..285  
                           id T77881  
                           est
- (ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: complement(126..206)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
                           region 139..219  
                           id R01713  
                           est
- (ix) FEATURE:  
 (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 70..147  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 10.5  
                           seq LLLALLLPVQVSS/FV
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 219:

```

AACCAGCCAG GAGCCACCCA TCCTCCAGCA CACTGAGCAG CAGGTGGAC ACACGGCACA   60
CTGATCCAA ATG GGT AAG GGG ATG GTG GCG ATG CTC ATT CTG GGT CTG CTA   111
Met Gly Lys Gly Met Val Ala Met Leu Ile Leu Gly Leu Leu
      -25              -20              -15

GTT CTG GCG CTG CTC CTA CCC GTG CAG GTT TCT TCA TTT GTT CCT TTA   159
Leu Leu Ala Leu Leu Leu Pro Val Gln Val Ser Ser Phe Val Pro Leu
      -10              -5              1

AAG AAT ATG CCG GAA GGT ACT GCA GCG GAA AGC ACA AAG CCG TCG AAT   217

```

Thr Ser Met Pro Glu Ala Thr Ala Ala Glu Thr Thr Lys Pro Ser Asn  
 5 10 15 20

GGG  
 Gly

210

## (2) INFORMATION FOR SEQ ID NO: 220:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 189 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 2..170
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
 region 4..172  
 id H56777  
 est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 25..87
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.5  
 seq LLVLFVLLANVQG/PG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 220:

AAGAAAGGCT GGCCTCTCTT CAAC ATG GGA TCT TCT GGA CTT TTG AGC CTC 51  
 Met Gly Ser Ser Gly Leu Leu Ser Leu  
 -20 -15

CTG GTG CTA TTC GTC CTC TTA GCG AAT GTC CAG GGA CCT CGT CTC ACT 99  
 Leu Val Leu Phe Val Leu Leu Ala Asn Val Gln Gly Pro Gly Leu Thr  
 -10 -5 1

GAT TGG TTA TTT CCC AGG AGA TGT CCC AAA ATC AGA GAA GAA TGT GAA 147  
 Asp Trp Leu Phe Pro Arg Arg Cys Pro Lys Ile Arg Glu Glu Cys Glu  
 5 10 15 20

TTC CAA GAA AGG GAT GTG TGT ACA AAG GAC AGA CAA TGC CGA 189  
 Phe Gln Glu Arg Asp Val Cys Thr Lys Asp Arg Gln Cys Arg  
 25 30

## (2) INFORMATION FOR SEQ ID NO: 221:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 323 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 52..253
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 10..216  
id R60167  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 235..319
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 194..278  
id R60167  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 143..253
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 90..205  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 55..145
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..91  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 235..316
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 183..264  
id R17888  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 141..253
- (C) IDENTIFICATION METHOD: blastn

171

.. (D) OTHER INFORMATION: identity 91  
region 85..202  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 56..144  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..89  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 235..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 180..264  
id N40052  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 58..257  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 1..200  
id AA039912  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 248..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 190..261  
id AA039912  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 50..319  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..270  
id R54127  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 90..194  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 8.3  
seq NLLLLHCVSRSHS/QN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 221:

AGGAAAGGCG TGACCACGCA GTTCTTGGGT CTGTCTGCT GCGCTGGGGT TGTGTTGAG 60

```

GCCGTGTCTC CGCTCCTGTG CCCGGGAAG ATG STG CTA GGT GGT TGC CCS GTT 113
Met Val Leu Gly Gly Cys Pro Val
-35 -30

AGT TAC TTA CTT CTG TGC GGC CAG GCG GCT TTG CTG CTG GGG AAT TTA 161
Ser Tyr Leu Leu Leu Cys Gly Gln Ala Ala Leu Leu Leu Gly Asn Leu
-25 -20 -15

CTT CTG CTG CAT TGT GTG TCT CGG AGC CAC TCG CAA AAT GCG ACC GCT 209
Leu Leu Leu His Cys Val Ser Arg Ser His Ser Gln Asn Ala Thr Ala
-10 -5 1 5

GAG CCT GAG CTC ACA TCC GCT GGC GCC CCC AGC CGG AGG GCC CCG GGG 257
Glu Pro Glu Leu Thr Ser Ala Gly Ala Pro Ser Arg Arg Ala Pro Gly
10 15 20

GTG CTG CGA GGT GGG AAT ATG GCG ACC CCC ACT CTC CGG TCA TCC TCT 305
Val Leu Arg Ala Gly Asn Met Ala Thr Pro Thr Leu Arg Ser Ser Ser
25 30 35

GCT CTT ACC TAC CTT GGG 323
Ala Leu Thr Tyr Leu Gly
40

```

## (2) INFORMATION FOR SEQ ID NO: 222:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 165 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 31..143
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 271..383  
id W16767  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 34..87
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.3  
seq LLSLSSLPLVLLG/WE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 222:

```

AACGAGACTT CTGACCCCTT GGGCAACAGC CAG ATG GAG ACT GGT GCG CTT TCG 34

```

(2) INFORMATION FOR SEQ ID NO: 223:

(A) LENGTH: 201 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(A) NAME/KEY: other  
(B) LOCATION: 24..203  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..180  
id HSC1PF091  
est

(A) NAME/KEY: other  
(B) LOCATION: 114..203  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 73..167  
id H03709  
est

(A) NAME/KEY: other  
(B) LOCATION: 35..107  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 1..73  
id H03709  
est

```
(A) NAME/KEY: sig_peptide
(3) LOCATION: 25..81
```



.. (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8.2  
 seq QVLALVLVAALWG/GT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 223:

```

AAAGTAGAAG ACAGCGGCGT TGCC ATG GCG GCG TCT CTG GGG CAG GTG TTG      51
Met Ala Ala Ser Leu Gly Gln Val Leu
-15

GCT CTG GTG CTG GTG GCC GCT CTG TGG GGT GGC ACG CAG CCG CTG CTG      99
Ala Leu Val Leu Val Ala Ala Leu Trp Gly Gly Thr Gln Pro Leu Leu
-10          -5          1          5

AAG CGG GCC TCC GCC GGC CTG CAG CGG GTT CAT GAG CCG ACC TGG GCC      147
Lys Arg Ala Ser Ala Gly Leu Gln Arg Val His Glu Pro Thr Trp Ala
          10          15          20

CAG CAG TTG CTA CAG GAG ATG AAG ACC CTC TTC TTG AAT ACT GAG TAC      195
Gln Gln Leu Leu Gln Glu Met Lys Thr Leu Phe Leu Asn Thr Glu Tyr
          25          30          35

CTG ATG      201
Leu Met
40

```

(2) INFORMATION FOR SEQ ID NO: 224:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 462 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 6..119
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 1..114  
id N83684  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 147..241
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 112..206  
id N83684  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 283..323
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 245..285  
id N83684  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 327..361
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 287..321  
id N83684  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 150..299
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 177..326  
id H94179  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 23..109
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 41..127  
id H94179  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 17..90
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..74  
id AA093069  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 144..194
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 123..173  
id AA093069  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 150..249
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 199..297  
id T67100

est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 211..387
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.1  
seq FLLGISNLSQVRA/SN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 224:

```

AGCGGGTGTT GAGAGCGGTG TGGTAGGTGT TGTAGCCGCT ATGGTGAAGT TCGCTTTGTA    60
GCGGCCCCGG CTAGAGAGTT GGCCTGTTC CTGCCTTTGT GACCCGGAGG AGCTTTTGGG   120
GTGCGTCAAG CCCCTGGCCT GAGGCAGCGA DCTGGTTTGT GGCCTGTTTG ATTCTGTCA   180
GAGGTTTGCT GACCCAAGAC AGTATCGAAA ATG CAT ATT AAG TCA ATT ATT CTA   234
                               Met His Ile Lys Ser Ile Ile Leu
                               -55

GAG GGA TTC AAG TCC TAT GCT CAG AGG ACC GAA GTC AAT GGT TTT GAC   282
Glu Gly Phe Lys Ser Tyr Ala Gln Arg Thr Glu Val Asn Gly Phe Asp
-50                      -45                      -40

CCC CTC TTC AAT GCT ATC ACT GGC TTA AAT GGT AGT GGG AAA TCC AAC   330
Pro Leu Phe Asn Ala Ile Thr Gly Leu Asn Gly Ser Gly Lys Ser Asn
-35                      -30                      -25                      -20

ATA TTG GAC TCC ATC TGC TTT TTG CTG GGC ATC TCC AAC CTG TCT CAG   378
Ile Leu Asp Ser Ile Cys Phe Leu Leu Gly Ile Ser Asn Leu Ser Gln
-15                      -10                      -5

GTT CGG GCT TCT AAT TTA CAA GAT TTA GTT TAC AAA AAT GGG CAG GCT   426
Val Arg Ala Ser Asn Leu Gln Asp Leu Val Tyr Lys Asn Gly Gln Ala
      1                      5                      10

GGT ATT ACC AAA GCC TCT GTG TCA ATC AMT TTT GAT   462
Gly Ile Thr Lys Ala Ser Val Ser Ile Xaa Phe Asp
15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 225:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 473 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 280..404

.. (C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 165..289  
id N45466  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 76..168  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 52..144  
id N46466  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 405..469  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 289..353  
id N46466  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(405..469)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 180..244  
id W86648  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(343..404)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 95  
region 244..305  
id W86648  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(297..347)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 302..352  
id W86648  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 273..358  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 39..123  
id W86523  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 357..404
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 123..170  
id W86523  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 405..436
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 170..201  
id W86523  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 285..341
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8  
seq WGFLCVLFTAVHP/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 225:

```

AAGMSAGGGG AAGCGCCCAA GGTCACACAG CTGGGATGTG GCAGAGCTGG GGTTCAGCT   60
CCTGTTCCCA TTGCTGGACA GCTGCCACAT CTGGCACCCA ATTAGGACC CCGCGGGGAG   120
GCCAAGCCC CGGGGGTGGC GGGGGATCCT AGAGGAAAGT GGCAAGGCCA GGACCCTGGA   180
GCAGAGCCAG AGTAGAAAC TGAGGCTCTG AGAGATGAAG CTACTTGCCA AGGTCACGCA   240
GCACAGTCAC ATCCTACTGA ACATCATCCT GTTCTCTGGG TGGA ATG TCA CCA TCG   296
                               Met Ser Pro Ser
CCC AGG TGG GGA TTT TTG TGT GTT TTG TTC ACT GCT GTA CAC CCA GCC   344
Pro Arg Trp Gly Phe Leu Cys Val Leu Phe Thr Ala Val His Pro Ala
-15          -10          -5          1
CCC AGC ACA GCG CCT GTC CAG GAC AAG TGC CCA GTA AAC ACT TGG GAA   392
Pro Ser Thr Ala Pro Val Gln Asp Lys Cys Pro Val Asn Thr Trp Glu
          5          10          15
GCA ATG CAB VMG GTC CTC CCA GCA GCT CCT GCA AAC AGA CCC CCG ACC   440
Ala Met Xaa Xaa Val Leu Pro Ala Ala Pro Ala Asn Arg Pro Pro Thr
          20          25          30
CAA GCC TTT CCT TCT GCM TCC ACT GCC ACA GGG   473
Gln Ala Phe Pro Ser Ala Ser Thr Ala Thr Gly
          35          40

```

## (2) INFORMATION FOR SEQ ID NO: 226:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 250 base pairs
- (B) TYPE: NUCLEIC ACID

-- (C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:  
(A) NAME/KEY: other  
(B) LOCATION: complement(1..189)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 99  
region 1..189  
id R47502  
est

(ix) FEATURE:  
(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 56..127  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.8  
seq FLLCLCIAYWAST/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 226:

|                                                                  |     |
|------------------------------------------------------------------|-----|
| AATTCTCATC GCGATTGCAC TCATCAAAGA AGCCAGCAGG GCTGTGGGAT ACGTC ATG | 53  |
| Met                                                              |     |
| TGC TCC TTG CTC TAC CCA CTG GTC ACC TTC TTC TTG CTG TGC CTC TGC  | 106 |
| Cys Ser Leu Leu Tyr Pro Leu Val Thr Phe Phe Leu Leu Cys Leu Cys  |     |
| -20 -15 -10                                                      |     |
| ATC GCC TAC TGG GCC AGC ACT GCT GTC TTC CTG TCC ACT TCC AAC GAA  | 154 |
| Ile Ala Tyr Trp Ala Ser Thr Ala Val Phe Leu Ser Thr Ser Asn Glu  |     |
| -5 1 5                                                           |     |
| GCG GTC TAT AAG ATC TTT GAT GAC AGC CCC TGC CCA TTT ACT GCG AAA  | 202 |
| Ala Val Tyr Lys Ile Phe Asp Asp Ser Pro Cys Pro Phe Thr Ala Lys  |     |
| 10 15 20 25                                                      |     |
| ACC TGC AAC CCA GAG ACC TTC CCC TCC TCC AAT GAG CCC CGC CAT GGG  | 250 |
| Thr Cys Asn Pro Glu Thr Phe Pro Ser Ser Asn Glu Pro Arg His Gly  |     |
| 30 35 40                                                         |     |

(2) INFORMATION FOR SEQ ID NO: 227:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 176 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens

.. (F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(51..119)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 404..472  
id AA099571  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(118..174)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 348..404  
id AA099571  
est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 24..71  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.6  
seq FLFFSTLFSSIFT/EA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 227:

```
AAAAGCTTTG GAGATATTGA ATC ATG TTA CCA TTT CTG TTT TTT TCC ACC CTG      53
Met Leu Pro Phe Leu Phe Phe Ser Thr Leu
-15                               -10

TTT TCT TCC ATA TTT ACT GAA GCT CAG AAG CAG TAT TGG GTC TGC AAC      101
Phe Ser Ser Ile Phe Thr Glu Ala Gln Lys Gln Tyr Trp Val Cys Asn
-5                               1           5           10

TCA TCC GAT GCA AGT ATT CAT ACA CCT ACT GTG ATA AAA TGC AAT ACC      149
Ser Ser Asp Ala Ser Ile His Thr Pro Thr Val Ile Lys Cys Asn Thr
15                               20           25

CAA TTT CAA TTA ATG TTA ACC CCT GGG      176
Gln Phe Gln Leu Met Leu Thr Pro Gly
30                               35
```

(2) INFORMATION FOR SEQ ID NO: 228:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 383 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 103..248  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 12..157  
id W56658  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 255..385  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 99  
region 164..294  
id W56658  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 18..248  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 1..231  
id AA127477  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 123..385  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 1..263  
id N40410  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(340..371)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 354..385  
id R93185  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 126..167  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.5  
seq VALNLLVPCCAA/WC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 228:

AATGTATGT TACGATGTTG TATTGATTTT TAAGAAAGTA ATTKRATTTG TAAAACTTCT 60  
JCTCGTTTAC ACTGCACATT GAATACAGGT AACTAATTGG WWGGAGAGGG GAGGTCACTC 120  
TTTTG ATG GTG GCC CTG AAC CTC ATT CTG GTT CCC TGC TGC GCT GCT TGG 170



```

Met Val Ala Leu Asn Leu Ile Leu Val Pro Cys Cys Ala Ala Trp
      -10                      -5                      1
TGT GAC CCA CGG AGG ATC CAC TCC CAG GAT GAC GTG CTC CGT AGC TCT    218
Cys Asp Pro Arg Arg Ile His Ser Gln Asp Asp Val Leu Arg Ser Ser
      5                      10                      15
GCT GCT GAT ACT GGG TCT GCG ATG CAG CGG CGT GAG GCC TGG GCT GGT    266
Ala Ala Asp Thr Gly Ser Ala Met Gln Arg Arg Gln Ala Trp Ala Gly
      20                      25                      30
TGG AGA AGG TCA CAA CCC TTC TCT GTT GGT CTG CCT TCT GCT GAA AGA    314
Trp Arg Arg Ser Gln Pro Phe Ser Val Gly Leu Pro Ser Ala Glu Arg
      35                      40                      45
CTC GAG AAC CAA CCA GGG AAG CTG TCC TGG AGG TCC CTG GTC GGA GAG    362
Leu Glu Asn Gln Pro Gly Lys Leu Ser Trp Arg Ser Leu Val Gly Glu
      50                      55                      60                      65
GGA CAT AGA ATC TGT GAC CTC    383
Gly His Arg Ile Cys Asp Leu
      70

```

## (2) INFORMATION FOR SEQ ID NO: 229:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 372 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 83..291
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 69..277  
id AA149265  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 12..57
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 1..46  
id AA149265  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 321..351
- (C) IDENTIFICATION METHOD: blastn

-- (D) OTHER INFORMATION: identity 93  
region 310..340  
id AA149265  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 81..372  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 53..344  
id W39570  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 27..57  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 2..32  
id W39570  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 81..372  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 55..346  
id N41332  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 24..57  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 94  
region 1..34  
id N41332  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 10..168  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.1  
seq IAVGLGVAALAF/GR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 229:

|                                                                   |     |
|-------------------------------------------------------------------|-----|
| AGCCTTGCC ATG GCT GCC CGT GGT GTC ATC GCT CCA GTT GCC GAG AGT YTG | 51  |
| Met Ala Ala Arg Gly Val Ile Ala Pro Val Gly Glu Ser Leu           |     |
| -50 -45 -40                                                       |     |
| CGC TAC GCT GAG TAC TTG CAG CCC TCG GCC AAA CGG CCA GAC GCC GAC   | 99  |
| Arg Tyr Ala Glu Tyr Leu Gln Pro Ser Ala Lys Arg Pro Asp Ala Asp   |     |
| -35 -30 -25                                                       |     |
| GTC GAC CAG CAG AGA CTG GTA AGA AGT TTG ATA GCT GTA GGA CTG GGT   | 147 |
| Val Asp Gln Gln Arg Leu Val Arg Ser Leu Ile Ala Val Gly Leu Gly   |     |

```

      -20              -15              -10
GTT GCA GCT CTT GCA TTT GCA GGT CGC TAC GCA TTT CGG ATC TGG AAA   135
Val Ala Ala Leu Ala Phe Ala Gly Arg Tyr Ala Phe Arg Ile Trp Lys
      -5              1              5
CCT CTA GAA CAA GTT ATC ACA GAA ACT GCA AAG AAG ATT TCA ACT CCT   243
Pro Leu Glu Gln Val Ile Thr Glu Thr Ala Lys Lys Ile Ser Thr Pro
      10              15              20              25
AGC TTT TCA TCC TAC TAT AAA GGA GGA TTT GAA CAG AAA ATG AGT AGG   291
Ser Phe Ser Ser Tyr Tyr Lys Gly Gly Phe Glu Gln Lys Met Ser Arg
      30              35              40
CGA GAA GCT GGT CTT ATT TTA GGT GTA AGC CCA TCT GCT GGC AAG GCT   339
Arg Glu Ala Gly Leu Ile Leu Gly Val Ser Pro Ser Ala Gly Lys Ala
      45              50              55
AAG ATT AGA ACA GCT CAT AGG AGA GTC ATG ATT   372
Lys Ile Arg Thr Ala His Arg Arg Val Met Ile
      60              65

```

## (2) INFORMATION FOR SEQ ID NO: 230:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 254 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..249
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..247  
id HUM225B05B  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 3..135
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 1..133  
id HUM224A06B  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 131..183
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92

region 128..180  
id HUM224A063  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 182..223  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 95  
region 178..219  
id HUM224A063  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(2..165)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 1..164  
id R81598  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 126..170  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.1  
seq KKKLLSLRPSLC/IP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 230:

```

AACACAAGCA AAACTTTTAA ATATTGGAAT TGACAGTTAC ATGTTTCATA ACTTTGTATG   60
TCTATTGGTT GTGCAGGTGT AATTTTTTCC CTTTTTGATT AGGGTTACAA AATTTAGAGA   120
CCAGT ATG ATT AAG TTG AAG CTC CTT AGC CTC CTT CGA CCT AGT CTC TGC   170
Met Ile Lys Leu Lys Leu Leu Ser Leu Leu Arg Pro Ser Leu Cys
  -15                -10                -5

ATA CCT CAA CTT TTA CGT ACC AAT GCT ACT CTG CTG TTC ACA ATT GCC   218
Ile Pro Gln Leu Leu Arg Thr Asn Ala Thr Leu Leu Phe Thr Ile Ala
   1                5                10                15

TCA TGT AAT CTG CAG ATT CCT GCC TCC CCA CGA CGG   254
Ser Cys Asn Leu Gln Ile Pro Ala Ser Pro Arg Arg
   20                25

```

## (2) INFORMATION FOR SEQ ID NO: 231:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 143 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

.. (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 100..144  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
                           region 95..139  
                           id T95183  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 56..105  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 90  
                           region 52..101  
                           id T95183  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 100..144  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
                           region 101..145  
                           id R48890  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 73..105  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 90  
                           region 75..107  
                           id R48890  
                           est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 13..77  
 (C) IDENTIFICATION METHOD: Von Haefne matrix  
 (D) OTHER INFORMATION: score 6.5  
                           seq GLCVLQLTTAVTS/AF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 231:

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| AACCTTCACA GTGTGAG ATG CCT AGT GTG AAC AGT GCT GGA TTA TGT GTC  | 50  |
| Met Pro Ser Val Asn Ser Ala Gly Leu Cys Val                     |     |
| -20 -15 -10                                                     |     |
| TTG CAG TTG ACA ACG GCA GTR ACC AGT GCC TTT TTA CTA GCA AAA GTG | 98  |
| Leu Gln Leu Thr Thr Ala Val Thr Ser Ala Phe Leu Leu Ala Lys Val |     |
| -5 1 5                                                          |     |
| AAT CCT TTC GAA RCT TTT CTC TCA AGG GGC TTT TGG CTA TGT GCT     | 143 |
| Asn Pro Phe Glu Xaa Phe Leu Ser Arg Gly Phe Trp Leu Cys Ala     |     |
| 10 15 20                                                        |     |

## (2) INFORMATION FOR SEQ ID NO: 232:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 178 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(118..179)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 296..357  
id T92237  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 86..145
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq ALFLLVSKYMIRS/GT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 232:

```

ACAGAGAMTA ACGATGTTTC TTATTTGAAT CCAAGTGAAG TACTCATGCT TTGTGTTCTT   60
GGGAATTACT GAGTTCAAAT TCCTA ATG ATG CTT GGG TTA CAC TTT GCT TTG   112
                Met Met Leu Gly Leu His Phe Ala Leu
                -20                      -15
TTT CTC CTA GTT TCT KTW TAT ATG ATC CGG AGT GGC ACT GGT AAT AAG   160
Phe Leu Leu Val Ser Xaa Tyr Met Ile Arg Ser Gly Thr Gly Asn Lys
-10                      -5                      5
ATT GAA GAA GGT GGG CGG   178
Ile Glu Glu Gly Gly Arg
                10

```

## (2) INFORMATION FOR SEQ ID NO: 233:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 181 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other  
(B) LOCATION: 32..178  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 2..148  
id H42383  
est

## (ix) FEATURE:

- (A) NAME/KEY: other  
(B) LOCATION: 35..178  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 5..148  
id R67703  
est

## (ix) FEATURE:

- (A) NAME/KEY: other  
(B) LOCATION: 145..178  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 91  
region 29..62  
id W90193  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 38..76  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.4  
seq MALLLSVLRVLLG/GF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 233:

```

ATCGGCGGGG CCAACCCACG GTGGGGGGGAG CGCGGCC ATG GCG CTC CTG CTT TCG      55
                                     Met Ala Leu Leu Leu Ser
                                     -10

GTG CTC CCT GTA CTG CTG GGC GGC TTC TTC GCG CTC GTG GGG TTG GCC      103
Val Leu Arg Val Leu Leu Gly Gly Phe Phe Ala Leu Val Gly Leu Ala
      -5                      1                      5

AAG CTC TCG GAG GAG ATC TCG GCT CCA GTD TCG GAG CGG ATG AAT GCC      151
Lys Leu Ser Glu Glu Ile Ser Ala Pro Val Ser Glu Arg Met Asn Ala
      10                      15                      20                      25

CTG TTC GTR MAG TTT GCT GAG GTG CTC GGG      181
Leu Phe Val Xaa Phe Ala Glu Val Leu Gly
      30                      35

```

## (2) INFORMATION FOR SEQ ID NO: 234:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 156 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 100..154
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 111..165  
id HSC2EB021  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 100..154
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 37..91  
id T31104  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 34..84
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2  
seq LWLSLVAWHWGEA/VL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 234:

```

ACTTTTTC A CGCTACTCCC CCGGAGTGCT TGG ATG TTG AAG AGT CTC TGG TTG      54
                               Met Leu Lys Ser Leu Trp Leu
                               -15

AGC CTT GTG GCC TGG CAC TGG GGT GAG GCT GTC CTC CTC TCC CCT CAT      102
Ser Leu Val Ala Trp His Trp Gly Glu Ala Val Leu Leu Ser Pro His
-10                -5                1                5

CTC CCT GCA GCG GCA GAA TGG CCC CGG GCA GCG TGT GAT TCG GGA GGT      150
Leu Pro Ala Ala Ala Glu Trp Pro Arg Ala Ala Cys Asp Ser Gly Gly
                10                15                20

GAA CCG  156
Glu Pro

```

## (2) INFORMATION FOR SEQ ID NO: 235:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 254 base pairs



- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 75..152
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 109..186  
id R38459  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 148..200
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 183..235  
id R38459  
est

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 183..227
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.1  
seq IVTWLLXSEFMSSA/EE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 235:

```

AATACTTTGG CAGCTTCTTC ACGTCGGTCC TCTCCGCGCG CGGGTAGGAA CCGTCCACGG      60
CCTTAAAGAA GCCTCCTCAC CAGCCATACT TCCATTGCC TCCAGCTGTT GCACGGAGGT      120
TTCACATCAT ATTTCCAGAA GGCTCCTGGA AACAGTGAAT ATGTGTGCGA TCCAGAGAGC      180
TG ATG GGG ATT GTG ACT TGG CTG CTG TMA TCC TTC ATG TCA AGC GCA      227
Met Gly Ile Val Thr Trp Leu Leu Xaa Ser Phe Met Ser Ser Ala
-15                -10                -5

GAA GAA TCT GTG TCA GCC CGC ACA CGG      254
Glu Glu Ser Val Ser Ala Arg Thr Arg
1                5

```

(2) INFORMATION FOR SEQ ID NO: 236:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 190 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 83..175
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 80..172  
id T62095  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 37..82
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 35..80  
id T62095  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 2..36
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..35  
id T62095  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 71..187
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 85..201  
id N43024  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 4..71
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 17..84  
id N43024  
est

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 37..187
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 26..176  
id W42796  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 86..187
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 114..215  
id AA030227  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 86..187
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 51..152  
id AA118270  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 80..163
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6  
seq IGLMFLMLGCALP/IY

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 236:

```
GTAGCGCGTC TTGGGTCTCC CGGCTGCCGC TGCTGCCGCC GCCGCCTCGG GTCGTGGAGC   60
CAGGAGCGAC GTCACCGCC ATG GCA GGC ATC AAA GCT TTG ATT AGT TTG TCC   112
              Met Ala Gly Ile Lys Ala Leu Ile Ser Leu Ser
              -25                               -20

TTT GGA GGA GCA ATC GGA CTG ATG TTT TTG ATG CTT GGA TGT GCC CTT   160
Phe Gly Gly Ala Ile Gly Leu Met Phe Leu Met Leu Gly Cys Ala Leu
      -15                -10                -5

CCA ATA TAC AAC AAA TAC TGG CCC CCC GGG   190
Pro Ile Tyr Asn Lys Tyr Trp Pro Pro Gly
      1                      5
```

## (2) INFORMATION FOR SEQ ID NO: 237:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 222 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(133..168)

(C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 270..299  
 id W73179  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: complement(139..168)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 295..324  
 id R59325  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: complement(113..144)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 218..249  
 id R06388  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 121..198  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.3  
 seq VKLVTLSPVPSLA/SS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 237:

```

ATTGTTCCTCA AAGGTTCCAA TTATTCAAGA CTGCCTTGGG CTTCCTTTTAC AACATGGATG    60
ATTCTATGTT ATGGGCACTG AAAGTAAAG AAAGTGTGGA AGGATTGGTA CATTAGAGAA    120
ATG AAA AAG CAA AAA CAT CAG AAA TTA TGG TGT ATT TCT GTA AAG TTA    168
Met Lys Lys Gln Lys His Gln Lys Leu Trp Cys Ile Ser Val Lys Leu
   -25               -20               -15

GTG ACA CTG AGT GTG CCC ACC TCT CTT GCC TCC TCT TTA ACC TCC CCT    216
Val Thr Leu Ser Val Pro Thr Ser Leu Ala Ser Ser Leu Thr Ser Pro
   -10               -5               1               5

ACA GGG    222
Thr Gly
  
```

## (2) INFORMATION FOR SEQ ID NO: 238:

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 417 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

## (11) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(227..414)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 94..1281  
id H53025  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 131..264
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 190..323  
id H52956  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 285..318
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 347..380  
id H52956  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 131..233
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 191..293  
id H53024  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 227..272
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 238..333  
id H53024  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 184..303
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq VLFALFVAFLLRG/KL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 238:

AAGTATCTCA CCAATTTCTTT CTCTTTCTGA ACCACATTGG GTGCCAACAG AASTTGCTCT 60

```

CTGTTCTCTT.TCAAAATTAC CAACATGGAC CCCACCCAAT TCTTCCCTTG GAACTAAGGA 120
ACGCCGTGACT GATCATCTGA TACAGCAGTK CCTGAGCAGA ACAAACAAAC AAAAACAGGA 180
CAG ATG GAT GGA ATA CCC ATG TCA ATG AAG AAT GAA ATG CCC ATC TCC 228
Met Asp Gly Ile Pro Met Ser Met Lys Asn Glu Met Pro Ile Ser
-40 -35 -30
CAA CTA CTG ATG ATC ATC GCC CCC TCC TTG GGA TTT GTG CTC TTC GCA 276
Gln Leu Leu Met Ile Ile Ala Pro Ser Leu Gly Phe Val Leu Phe Ala
-25 -20 -15 -10
TTG TTT GTG GCG TTT CTC CTG AGA GGG AAA CTC ATG GAA ACC TAT TGT 324
Leu Phe Val Ala Phe Leu Leu Arg Gly Lys Leu Met Glu Thr Tyr Cys
-5 1 5
TCG CAG AAA CAC ACA AGG CTA GAC TAC ATT GGA GAT AGT AAA AAT GTC 372
Ser Gln Lys His Thr Arg Leu Asp Tyr Ile Gly Asp Ser Lys Asn Val
10 15 20
CTC AAT GAC GTG CAG CAT GGA AGG GAA GAC GAA GAC GGC CAT GGG 417
Leu Asn Asp Val Gln His Gly Arg Glu Asp Glu Asp Gly His Gly
25 30 35

```

## (2) INFORMATION FOR SEQ ID NO: 239:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 293 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 246..293
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 90..137  
id H43824  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 246..293
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 93..110  
id B73173  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 246..293

.. (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
                           region 112..159  
                           id H26792  
                           est

## (1x) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 21..191  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.7  
                           seq LAICSCCLPGPGPA/LP

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 239:

|                                                                   |     |
|-------------------------------------------------------------------|-----|
| ACCTCGCTGC TCTTCATCCC ATG GGT GGA TTT TTG CAT CTC CCT GCT CTG TCT | 53  |
| Met Gly Gly Phe Leu His Leu Pro Ala Leu Ser                       |     |
| -55 -50                                                           |     |
| TCC TCC TGT CTT TGG ACA TTT CCA CCG ATG TGT GTT CGC ATC TTC TCC   | 101 |
| Ser Ser Cys Leu Trp Thr Phe Pro Pro Met Cys Val Arg Ile Phe Ser   |     |
| -45 -40 -35                                                       |     |
| TAT GTT CCT TTA CCT ATC CTG ACC CCC AAA ACC ATA AAT CTC ATC CCC   | 149 |
| Tyr Val Pro Leu Pro Ile Leu Thr Pro Lys Thr Ile Asn Leu Ile Pro   |     |
| -30 -25 -20 -15                                                   |     |
| GTT CTG GCC ATC TGT TCC TGT CTT CCT GGC CCC GGG CCG GCC CTT CCT   | 197 |
| Val Leu Ala Ile Cys Ser Cys Leu Pro Gly Pro Gly Pro Ala Leu Pro   |     |
| -10 -5 1                                                          |     |
| CTT CCT GCC TTC CCG ACC CTC CTT GTG TCT TGG TAC CAC TGC CCC CCA   | 245 |
| Leu Pro Ala Phe Pro Thr Leu Leu Val Ser Trp Tyr His Cys Pro Pro   |     |
| 5 10 15                                                           |     |
| CAG AAG AAG ACA GGC ATG ATG GAC ACG GAT GAT TTC CGC GCC TGC CCG   | 293 |
| Gln Lys Lys Thr Gly Met Met Asp Thr Asp Asp Phe Arg Ala Cys Pro   |     |
| 20 25 30                                                          |     |

## (2) INFORMATION FOR SEQ ID NO: 240:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 416 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (E) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 259..413  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 99

region 165..319  
id N46466  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 55..147  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 52..144  
id N46466  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 252..338  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 95  
region 38..124  
id W86523  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 336..413  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 123..200  
id W86523  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(322..413)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 214..305  
id W36648  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(276..326)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 302..352  
id W86648  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 264..320  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3  
seq WGFLCVLFYAVRP/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 240:

GTCCACACAGC TGGGATGTGG CAGAGSTGGG GTCCAGGTC CTGTTCCCAT TCTGGACAG 50



```

CTGCCACATC TGGCACCCAA TTTAGGACCC CGCGGGGAGG CCCAAGCCCC GGGGGTGGCG 120
GGGGATCCTA GAGGAAAGTG GCAAGGCCAG GACCCTGGAG CAGAGCCAGA GTAGAAAAC 180
GAGGCTCTGA GAGATGAAGC TACTTSCCAA GGTACGCAG CACAGTCACA TCCTACTGAA 240
CATCATCCTG TTCTCTGGGT GGA ATG TCA CCA TCG CCC AGG TGG GGA TTT TTG 293
                Met Ser Pro Ser Pro Arg Trp Gly Phe Leu
                -15                               -10

TGT GTB TTG TTC ACT GCT GTA CAC CCA GCC CCC AGC ACA GCG CCT GTC 341
Cys Val Leu Phe Thr Ala Val His Pro Ala Pro Ser Thr Ala Pro Val
                -5                               1                               5

CAG GAC AAG TGC CCA GTA AAC ACT TGG GAA GCA ATG CAA GCG TCC TCC 389
Gln Asp Lys Cys Pro Val Asn Thr Trp Glu Ala Met Gln Ala Ser Ser
                10                               15                               20

CAG CAG CTC CTG CAA ACA GAC CCC ATG 416
Gln Gln Leu Leu Gln Thr Asp Pro Met
                25                               30

```

## (2) INFORMATION FOR SEQ ID NO: 241:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 432 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 60..386
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 7..333  
id AA035203  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 400..429
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 349..373  
id AA035203  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 77..429
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99

region 1..353  
id H64963  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 50..328  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 10..288  
id R97144  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 56..393  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 3..340  
id N73170  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 14..300  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 2..288  
id H13072  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 154..381  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.3  
seq IILASASFSPNFT/QV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 241:

```

AGTAAAAAAA CACTGGAATA AGGAAGGGCT GATGACITTC AGAAGATGAA GGTAAGTAGA    60
AACCGTTGAT GGGACTGAGA AACCAGAGTK AAAACCTCTT TGGAGCTTCT GAGGACTCAG    120
CTGGAACCAA CGGGCACAGT TGGCAACACC ATC ATG ACA TCA CAA CCT GTT CCC    174
                               Met Thr Ser Gln Pro Val Pro
                               -75                               -70

AAT GAG ACC ATC ATA GTG CTC CCA TCA AAT GTC ATC AAC TTC TCC CAA    222
Asn Glu Thr Ile Ile Val Leu Pro Ser Asn Val Ile Asn Phe Ser Gln
                               -65                               -60                               -55

GCA GAG AAA CCC GAA CCC ACC AAC CAG GGG CAG CAT ACC CTG AAG AAA    270
Ala Glu Lys Pro Glu Pro Thr Asn Gln Gly Gln Asp Ser Leu Lys Lys
                               -50                               -45                               -40

CAT CTA CAC GCA GAR RTC AAA GTT ATT GCG ACT ATC CAG ATC TTG TGT    318
His Leu His Ala Glu Xaa Lys Val Ile Gly Thr Ile Gln Ile Leu Cys
                               -35                               -30                               -25

```

```

GGC ATG AFG GTA TTG AGC TTG GGG ATC ATT TTG GCA TCT GCT TCC TTC      366
Gly Met Met Val Leu Ser Leu Gly Ile Ile Leu Ala Ser Ala Ser Phe
  -20                      -15                      -10

TCT CCA AAT TTT ACC CAA GTG ACT TCT ACA CTG TTG AAC TCT GCT TAC      414
Ser Pro Asn Phe Thr Gln Val Thr Ser Thr Leu Leu Asn Ser Ala Tyr
  -5                      1                      5                      10

CCA TTC ATA GGA CCC GGG      432
Pro Phe Ile Gly Pro Gly
          15

```

## (2) INFORMATION FOR SEQ ID NO: 242:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 437 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 47..230
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 158..341  
id AA040813  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 229..395
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 341..507  
id AA040813  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 205..429
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 111..335  
id H34584  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(325..422)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 225..323

id AAC40149  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(215..269)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 381..435  
id AAC40149  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: complement(279..327)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 321..369  
id AAC40149  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 57..329  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.8  
seq IILRLPWLNRST/VV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 242:

|                                                                    |     |
|--------------------------------------------------------------------|-----|
| ACGCTTCGTC CTCTGCAGTC AAGACGCTGG GCGCGTCGAG GACTGGGATT TCAAAAT ATG | 59  |
| Met                                                                |     |
| CGT GCA TTA GAG AAT GAT TTT TTC AAT TCT CCC CCA AGA AAA ACT GTT    | 107 |
| Arg Ala Leu Glu Asn Asp Phe Phe Asn Ser Pro Pro Arg Lys Thr Val    |     |
| -90 -85 -30 -75                                                    |     |
| CGG TTT GGT GGA ACT GTG ACA GAA GTC TTG CTG AAG TAC AAA AAG GGT    | 155 |
| Arg Phe Gly Gly Thr Val Thr Glu Val Leu Leu Lys Tyr Lys Lys Gly    |     |
| -70 -65 -60                                                        |     |
| GAA ACA AAT GAC TTT GAG TTG TTG AAG AAC CAG CTG TTA GAT CCA GAC    | 203 |
| Glu Thr Asn Asp Phe Glu Leu Leu Lys Asn Gln Leu Leu Asp Pro Asp    |     |
| -55 -50 -45                                                        |     |
| ATA AAG GAT GAC CAG ATC ATC AAC TGG CTG CTA GAA TTC CGT TCT TCT    | 251 |
| Ile Lys Asp Asp Gln Ile Ile Asn Trp Leu Leu Glu Phe Arg Ser Ser    |     |
| -40 -35 -30                                                        |     |
| GTC ATG TAC TTG ACA AAA GAC TTT GAG CAA CTT ATC AGT ATT ATA TTG    | 299 |
| Val Met Tyr Leu Thr Lys Asp Phe Glu Gln Leu Ile Ser Ile Ile Leu    |     |
| -25 -20 -15                                                        |     |
| AGA TTG COT TGG TTG AAT AGA AGT CAA ACA GTA GTG GAA GAG TAT TTG    | 347 |
| Arg Leu Pro Trp Leu Asn Arg Ser Gln Thr Val Val Glu Glu Tyr Leu    |     |
| -10 -5 1 3                                                         |     |
| GCT TTT CTT GGT AAT CTT GTA TCA GCA GAG AGT GTT TTC CTC AAG CCG    | 395 |
| Ala Phe Leu Gly Asn Leu Val Ser Ala Glu Thr Val Phe Leu Arg Pro    |     |
| 10 15 20                                                           |     |

TGT CTC AGC ATG ATT GCT TCC CAT TTT GWG CCT CCC GAG CTG  
Cys Leu Ser Met Ile Ala Ser His Phe Xaa Pro Pro Glu Leu  
25 30 35

437

## (2) INFORMATION FOR SEQ ID NO: 243:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 244 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 54..242
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 12..200  
id R19497  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 78..242
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..165  
id H75597  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 84..242
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 1..159  
id H93398  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 122..243
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..122  
id HUM030E11B  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 74..166
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8  
seq WAFSCGTWLPSRA/EW

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 243:

```

ATAGAAGGGG GTGGGGCCAC GTTTGCGTCC GCGCCATCAG GCCCGAGATA GCGGCGAGGT    60
CGGCTTTTCAG TGT ATG GTT TTC CCT GCC AAA CGG TTC TGC TTG GTG CCA    109
      Met Val Phe Pro Ala Lys Arg Phe Cys Leu Val Pro
      -30                -25                -20

TCC ATG GAG GGC GTG CGC TGG GCC TTT TCC TGC GGC ACT TGG CTG CCG    157
Ser Met Glu Gly Val Arg Trp Ala Phe Ser Cys Gly Thr Trp Leu Pro
      -15                -10                -5

AGC CGA GCC GAA TGG CTG CTG GCA GTG CGA TCG ATT CAG CCC GAG GAG    205
Ser Arg Ala Glu Trp Leu Leu Ala Val Arg Ser Ile Gln Pro Glu Glu
      1                5                10

AAG GAG CGC ATT GGC CAG TTC GTC TTT GCC CGG GAC GGG    244
Lys Glu Arg Ile Gly Gln Phe Val Phe Ala Arg Asp Gly
      15                20                25

```

(2) INFORMATION FOR SEQ ID NO: 244:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 373 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 101..273
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 159..331  
id W57194  
est

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 95..340
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq LTCLADLFHSIAT/XK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 244:

```

AAGAGGAGAA GGCTGGGTAT GAGGAACTG CTGCTTTTC ACTGGATCAG ACCTAAATAT    60

```

```

TAACTGGACC-TCTCTTTAGA TTCTTTGCTC AATA ATG AAT TGT TTT CAG GGC ACC 115
Met Asn Cys Phe Gln Gly Thr
-80

AAT GCC TCT GCT CTG GAA AAA GAC ATT GGT CCA GAG CAG TTT CCA ATC 163
Asn Ala Ser Ala Leu Glu Lys Asp Ile Gly Pro Glu Gln Phe Pro Ile
-75 -70 -65 -60

AAT GAA CAC TAT TTC GGA TTG GTC AAT TTT GGA AAC ACA TGC TAC TGT 211
Asn Glu His Tyr Phe Gly Leu Val Asn Phe Gly Asn Thr Cys Tyr Cys
-55 -50 -45

AAC TCC GTG CTT CAG GCA TTG TAC TCC TGC CGT CCA TTC CGG GAG AAT 259
Asn Ser Val Leu Gln Ala Leu Tyr Ser Cys Arg Pro Phe Arg Glu Asn
-40 -35 -30

GTG TTG GCA TAC AAG GCC CAG CAA AAG AAG GAA AAC TTG CTG ACG 307
Val Leu Ala Tyr Lys Ala Gln Gln Lys Lys Lys Glu Asn Leu Leu Thr
-25 -20 -15

TGC CTG GCG GAC CTT TTC CAC AGC ATT GCC ACA SAG AAG AAG AAG GTT 355
Cys Leu Ala Asp Leu Phe His Ser Ile Ala Thr Xaa Lys Lys Lys Val
-10 -5 1 5

GBR TCA TCC CAC CTG GGG 373
Xaa Ser Ser His Leu Gly
10

```

## (2) INFORMATION FOR SEQ ID NO: 245:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 184 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 73..182
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 68..177  
id W60868  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 62..182
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 1..121  
id C17761  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: complement(150..182)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 96  
                           region 185..217  
                           id W60944  
                           est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 20..67  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.6  
                           seq ALRVRXXXFGTRA/CR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 245:

```

AATTTCGGAS CCGGGCAAG ATG GCA GCG GCG CTG CGC GTG CGT KGT TSA STG      52
      Met Ala Ala Ala Leu Arg Val Arg Xaa Xaa Xaa
      -15                               -10

TTC GGG ACG CGG GCC TGC AGG CGC CAT GGT CTT CCT CAC CGC GCA STC      100
Phe Gly Thr Arg Ala Cys Arg Arg His Gly Leu Pro His Arg Ala Xaa
-5                               1                               5                               10

TGG CTG CGG AAT CGC GTC ASC GAC CGC TAC TTT CGG ATC CAG GAG GTG      148
Trp Leu Arg Asn Arg Val Xaa Asp Arg Tyr Phe Arg Ile Gln Glu Val
15                               20                               25

CTG AAG CAS GCC AGG CAC TTC CGG GGA AGG AAA AGG                        184
Leu Lys Xaa Ala Arg His Phe Arg Gly Arg Lys Arg
30                               35

```

## (2) INFORMATION FOR SEQ ID NO: 246:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 190 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 35..186  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
                           region 1..182  
                           id AA058818  
                           est



## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 32..135
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..104  
id T50012  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 134..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 102..154  
id T50012  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 56..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 33..163  
id H79942  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 21..135
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 3..117  
id AA058605  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 134..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 115..167  
id AA058605  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 48..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 1..139  
id R37526  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 56..100
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq LLTHNLLSSHVRC/VC

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 246:

```

CTAATCGAAA AGGGGGATTT TCCGGTTCCG GCCTGGCGAG AGTTTGTGGG GCGAC ATG      58
                                     Met
                                     -15

AAA CTG CTT ACC CAC AAT CTG CTG AGC TCG CAT GTG CCG GGG GTG GGG      106
Lys Leu Leu Thr His Asn Leu Leu Ser Ser His Val Arg Gly Val Gly
          -10                      -5                      1

TCC CGT GGC TTC CCC CTG CGC CTC CAG GCC ACC GAG GTC CGT ATC TGC      154
Ser Arg Gly Phe Pro Leu Arg Leu Gln Ala Thr Glu Val Arg Ile Cys
          5                      10                      15

CCT GTG GAA TTC AAC CCC AAC TTC GTG GCG CGA CGG                      190
Pro Val Glu Phe Asn Pro Asn Phe Val Ala Arg Arg
      20                      25                      30

```

(2) INFORMATION FOR SEQ ID NO: 247:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 189 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 54..186
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 45..177  
id HSC2KH091  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 9..52
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 1..44  
id HSC2KH091  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 82..117
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 1..36  
id AA090704  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 129..186  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
                           region 36..93  
                           id AA126596  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 93..131  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 94  
                           region 1..39  
                           id AA126596  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 122..181  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 91  
                           region 40..99  
                           id AA090640  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 88..117  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 96  
                           region 8..37  
                           id AA090640  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 99..186  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 95  
                           region 1..88  
                           id T36119  
                           est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 7..129  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.6  
                           seq VSAGSLLLLPAPQA/EX

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 247:

|        |     |     |     |     |     |     |     |     |     |     |     |     |     |     |    |
|--------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|
| AACGGG | ATG | GGA | TWC | TTC | TCA | CGG | CGC | ACG | TTC | TGT | GGG | CGG | AGT | GGG | 48 |
|        | Met | Gly | Lys | Phe | Ser | Arg | Arg | Thr | Phe | Cys | Gly | Arg | Ser | Gly |    |
|        | -40 |     |     |     |     |     | -35 |     |     |     |     |     | -30 |     |    |
| CGG    | AGC | TGG | CGG | GGT | CAG | TTG | GTC | CAA | GTG | TCC | CGG | CCT | GAG | GTG | 96 |

```

Arg Ser Cys Arg Gly Gln Leu Val Gln Val Ser Arg Pro Glu Val Ser
   -25                -20                -15

GCC GGA TCC CTC CTT CTC CCG GCG CCT CAA GCG GAA GAS CAT TCC TCA    144
Ala Gly Ser Leu Leu Leu Pro Ala Pro Gln Ala Glu Xaa His Ser Ser
   -10                -5                1                5

WGR RTT TTG TAT CCA AGG CCC AAA AGT TTG TTA CCC AAG ATG GGG    189
Xaa Xaa Leu Tyr Pro Arg Pro Lys Ser Leu Leu Pro Lys Met Gly
          10                15                20

```

## (2) INFORMATION FOR SEQ ID NO: 248:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 237 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 132..235
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 94..197  
id R36207  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 37..110
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..74  
id R36207  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 41..194
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 94  
region 1..154  
id AA090796  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 107..235
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 15..143  
id AA091961

est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 141..193  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 96  
 region 33..85  
 id AA091520  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 190..237  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
 region 81..128  
 id AA091520  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 109..142  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 2..35  
 id AA091520  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 1..165  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.6  
 seq CALSLPDRPGASG/GR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 248:

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| ATG GAA GGA GGC GTT CGT CTA GAT TTG TCG GGT TGC GGG GAG ACT TCA | 48  |
| Met Glu Gly Gly Val Arg Leu Asp Leu Ser Ala Cys Gly Glu Thr Ser |     |
| -55 -50 -15 -40                                                 |     |
| GGA GTC GCT GTC TCT GAA CTT CCA GCC TCA GAG ACC GCC GCC CTT GTC | 96  |
| Gly Val Ala Val Ser Glu Leu Pro Ala Ser Glu Thr Ala Ala Leu Val |     |
| -35 -30 -25                                                     |     |
| CCC GAG GGC CAT GGG CCG GGT CTC AGG GCT TGT GCC CTC TCG CTT CCT | 144 |
| Pro Glu Gly His Gly Pro Gly Leu Arg Ala Cys Ala Leu Ser Leu Pro |     |
| -20 -15 -10                                                     |     |
| GAC GCT CCT GGC GCA TCT GGT GGT CGT CAT GAG CTT ATT CTG GTC CCG | 192 |
| Asp Ala Pro Gly Ala Ser Gly Gly Arg His His Leu Ile Leu Val Pro |     |
| -5 1 5                                                          |     |
| GGA CAG CAA CAT ACA GGC CTG CCT GCC TCT GAG GGT CAC CCC CAG     | 237 |
| Gly Gln Gln His Thr Gly Leu Pro Ala Ser His Val His Pro Gln     |     |
| 10 15 20                                                        |     |

## (2) INFORMATION FOR SEQ ID NO: 249:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 144..213
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 1..70  
id N53816  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 21..63
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..43  
id T34269  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 163..204
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq TLLSFAALTAAFS/VL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 249:

```

AAGCGCCGGA HGCGGTGAGG CACAGATGAG TAACGTGAAT TTGTCCGTCT CCGACTTCTG   60
GAGGTAAGGC GGTGTCAGC CTATCTCTTC TGCTGGCTGG GCTCAATGCC GCGGGTGAGC  120
GTTGCGCCGA GGCTGCTCCT ACCCTTGAGT GATGTGCCTT GA ATG ACG CTG CTT      174
   Met Thr Leu Leu
TCA TTC GCT GCT CTC ACG GCT GCT TTC TCC GTC CTC CCC AAG          216
Ser Phe Ala Ala Leu Thr Ala Ala Phe Ser Val Leu Pro Lys
-10              -5              1

```

## (2) INFORMATION FOR SEQ ID NO: 250:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 269 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE

-- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 46..271

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 98  
region 36..261  
id HSC3IF011  
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 11..44

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 100  
region 2..35  
id HSC3IF011  
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 50..271

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 98  
region 39..260  
id N28442  
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 18..234

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 98  
region 1..217  
id HUM517CG1B  
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 125..215

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 98  
region 1..91  
id T77607  
est

(ix) FEATURE:

(A) NAME/KEY: other

(B) LOCATION: 217..271

(C) IDENTIFICATION METHOD: blastn

(D) OTHER INFORMATION: identity 94  
region 92..146  
id T77607  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 36..98
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4  
seq GLSKLQFAPFSSA/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 250:

```

AGTGGTTGCN GGAAGTTGAG CGGCGGCAAG AAATA ATG GCG GCA GCT ACG GGG      53
                               Met Ala Ala Ala Thr Gly
                               -20

GAT CCT GGA CTC TCT AAA CTG CAG TTT GCC CCT TTT AGT AGT GCC TTG      101
Asp Pro Gly Leu Ser Lys Leu Gln Phe Ala Pro Phe Ser Ser Ala Leu
-15                      -10                      -5                      1

GAT GTT GGG TTT TGG CAT GAG TTG ACC CAG AAG AAG CTG AAC GAG TAT      149
Asp Val Gly Phe Trp His Glu Leu Thr Gln Lys Lys Leu Asn Glu Tyr
                    5                      10                      15

CGG CTG GAT GAA GCT CCC AAG GAC ATT AAG GGT TAT TAC TAC AAT GGT      197
Arg Leu Asp Glu Ala Pro Lys Asp Ile Lys Gly Tyr Tyr Tyr Asn Gly
                20                      25                      30

GAC TCT GCT GGG MTG CCA GCT CGC TTA ACA TTG GAG TTC AGT GCT TTT      245
Asp Ser Ala Gly Xaa Pro Ala Arg Leu Thr Leu Glu Phe Ser Ala Phe
    35                      40                      45

GAC ATG AGT GCT CCC ACC CCA AGC      269
Asp Met Ser Ala Pro Thr Pro Ser
    50                      55

```

## (2) INFORMATION FOR SEQ ID NO: 251:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 145 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 39..143
- (C) IDENTIFICATION METHOD: Blastn
- (D) OTHER INFORMATION: identity 96  
region 50..154  
id R50695  
est

## (ix) FEATURE:



```

.. (A) NAME/KEY: other
   (B) LOCATION: 3..45
   (C) IDENTIFICATION METHOD: blastn
   (D) OTHER INFORMATION:  identity 90
                           region 15..57
                           id R50695
                           est

```

## (ix) FEATURE:

```

(A) NAME/KEY: other
(B) LOCATION: 81..143
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION:  identity 96
                           region 104..166
                           id R94786
                           est

```

## (ix) FEATURE:

```

(A) NAME/KEY: other
(B) LOCATION: 81..143
(C) IDENTIFICATION METHOD: blastn
(D) OTHER INFORMATION:  identity 96
                           region 105..167
                           id T98442
                           est

```

## (ix) FEATURE:

```

(A) NAME/KEY: sig_peptide
(B) LOCATION: 50..130
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION:  score 4.4
                           seq LSKSLLLVPSXLS/LL

```

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 251:

```

AAGCTTCCCC TCCCCCGGCG CCCTCTGGGG CTCCGAGCCC GGCGGGACC ATG TTC ACC      58
                                   Met Phe Thr
                                   -25

AGC ACC GGC TCC AGT GGG CTC TAC AAG GCG CCT CTG TCG AAG AGC CTT      106
Ser Thr Gly Ser Ser Gly Leu Tyr Lys Ala Pro Leu Ser Lys Ser Leu
          -20                      -15                      -10

CTG CTG GTC CCC AGT RCC CTC TCC CTC CTG CSC GCC CAG                  145
Leu Leu Val Pro Ser Xaa Leu Ser Leu Leu Xaa Ala Gln
          -5                      1                      5

```

## (2) INFORMATION FOR SEQ ID NO: 252:

## (i) SEQUENCE CHARACTERISTICS:

```

(A) LENGTH: 427 base pairs
(B) TYPE: NUCLEIC ACID
(C) STRANDEDNESS: DOUBLE
(D) TOPOLOGY: LINEAR

```

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

-- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 137..291  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 99  
region 139..292  
id AA121372  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 6..91  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..86  
id AA121372  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 318..397  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 322..401  
id AA121372  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 95..132  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 94..131  
id AA121372  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 284..313  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 286..315  
id AA121372  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 2..102  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 15..115  
id T53974  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 150..253  
(C) IDENTIFICATION METHOD: blastn

(C) OTHER INFORMATION: identity 92  
region 167..275  
id T53974  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 95..171  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 111..187  
id T53974  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 2..102  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
region 15..115  
id R09314  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 95..171  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 111..187  
id R09314  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 150..222  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 167..239  
id R09314  
est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 179..298  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.3  
seq ITLVSAAPGKVIC/EM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 252:

```
AAAATCGCGG ACCACCGGGG CTGCCAKCTC GCCTSACTCC CGGCCTCTTG CGCTCCTAGG    60
GGCGGAGAAG GGTGCGGGCT CTTGCCCCTT TGTGTCTTC TTCTACTAAC TTCTGGACTT    120
TCCAGCTCTT CGAAGTTTCG TTCTTGCGCA AAGCCCAAAG GCTGGAAAAC CGTCCACG    178
ATG ACC AGC ATG ACT CAG TCT CTG CGG GAG GTG ATA AAG GCC ATG ACC    226
Met Thr Ser Met Thr Gln Ser Leu Arg Glu Val Ile Lys Ala Met Thr
-40          -35          -30          -25
```

```

AAG GCT CGC AAT TTT GAG AGA GTT TTG GGA AAG ATT ACT CTT GTC TCT      274
Lys Ala Arg Asn Phe Glu Arg Val Leu Gly Lys Ile Thr Leu Val Ser
      -20                      -15                      -10

GCT GCT CCT GGG AAA GTG ATT TGT GAA ATG AAA GTA GAA GAA GAG CAT      322
Ala Ala Pro Gly Lys Val Ile Cys Glu Met Lys Val Glu Glu Glu His
      -5                      1                      5

ACC AAT GCA ATA GGC ACT CTC CAC GGC GGT TTG ACA GGC ACG TTA GTA      370
Thr Asn Ala Ile Gly Thr Leu His Gly Gly Leu Thr Ala Thr Leu Val
      10                      15                      20

GAT AAC ATA TCA ACA ATG GCT CTG CTA TGC ACG GAA AGG GGA GCA CCC      418
Asp Asn Ile Ser Thr Met Ala Leu Leu Cys Thr Glu Arg Gly Ala Pro
      25                      30                      35                      40

GGA GTC AGT      427
Gly Val Ser

```

## (2) INFORMATION FOR SEQ ID NO: 253:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 332 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 2..285
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 8..291  
id T31110  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 278..331
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 285..338  
id T31110  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 2..329
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 6..333  
id T09244  
est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 2..329  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 98  
                           region 7..334  
                           id T35807  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 9..331  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 98  
                           region 1..323  
                           id T33763  
                           est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 15..331  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 99  
                           region 1..317  
                           id AA132848  
                           est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 75..293  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.3  
                           seq DIILSGLVPGSTT/LH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 253:

```

AAGCGAGCCC AGGCGGCAGT CTTGATTCCC TTTTGGCCAG CAGTTTTTAG GTCTGTCAGT   60
ACTGCACATGC AAGA ATG GCA GAT TTT GGG ATC TCT GCT GGC CAG TTT GTG   110
      Met Ala Asp Phe Gly Ile Ser Ala Gly Gln Phe Val
                        -70                      -65

GCA GTS GTC TGG GAT AAG TCA TCC CCA GTG GAG GCT CTG AAA GGT CTG   158
Ala Val Val Trp Asp Lys Ser Ser Pro Val Glu Ala Leu Lys Gly Leu
      -60                      -55                      -50

GTG GAT AAG CTT CAA GCG TTA ACC GGC AAT GAG GGC CGC GTG TCT GTG   206
Val Asp Lys Leu Gln Ala Leu Thr Gly Asn Glu Gly Arg Val Ser Val
      -45                      -40                      -35                      -30

GAA AAC ATC AAG CAG CTG TTG CAA TCT GCC CAC AAA GAA TCC AGC BTT   254
Glu Asn Ile Lys Gln Leu Leu Gln Ser Ala His Lys Glu Ser Ser Xaa
      -25                      -20                      -15

GAC ATT ATT TTG TCA GGT TTA GTC CCA GGA AGC ACC ACT CTG CAC AGT   302
Asp Ile Ile Leu Ser Gly Leu Val Pro Gly Ser Thr Thr Leu His Ser
      -10                      -5                      1

GCT GAG ATT TTG GCT GAA ATC GCG TGG GTC   332

```

(A) LENGTH: 131 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

[illegible]

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 84..125  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.2  
seq GILLGLLLLLGHLT/VR

AACAGACGCT GGCGGCCACC AGAAGTTTGA GCCTCTTTGG TAGCAGGAGG CTGGAAGAAA 60  
GGACAGAAGT AGCTCTGGCT GTG ATG GGG ATC TTA CTG GGC CTG CTA CTC CTG 113  
Met Gly Ile Leu Leu Gly Leu Leu Leu Leu  
-10 -5  
GGG CAC CTA ACA GTG AGA 131  
Gly His Leu Thr Val Arg  
1

(A) LENGTH: 486 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(v1) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 13..53  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 90  
region 1..41  
id AA063860  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 55..111  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.1  
seq LLLGQRCSLKVSG/QE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 255:

|                                                                 |     |     |
|-----------------------------------------------------------------|-----|-----|
| AAATCTTCAG GGCAGCTCCC AGAGCATGGA TCCCTCCTGA TTCCACTCAG CCCG     | ATG | 57  |
|                                                                 | Met |     |
| TTC CTC ACA GTC AAG CTG CTC CTG GGC CAG AGA TGC AGT CTG AAG GTG |     | 105 |
| Phe Leu Thr Val Lys Leu Leu Leu Gly Gln Arg Cys Ser Leu Lys Val |     |     |
| -15 -10 -5                                                      |     |     |
| TCA GGG CAA GAG AGT GTA GCC ACG CTG AAG AGA CTG GTG TCC AGG CGG |     | 153 |
| Ser Gly Gln Glu Ser Val Ala Thr Leu Lys Arg Leu Val Ser Arg Arg |     |     |
| 1 5 10                                                          |     |     |
| CTG AAG GTG CCT GAG GAG CAG CAG CAC CTG CTT TTC CGT GGC CAG CTC |     | 201 |
| Leu Lys Val Pro Glu Glu Gln Gln His Leu Leu Phe Arg Gly Gln Leu |     |     |
| 15 20 25 30                                                     |     |     |
| CTG GAG GAT GAC AAG CAC CTC TCT GAC TAC TGC ATT GGG CCC AAT GCC |     | 249 |
| Leu Glu Asp Asp Lys His Leu Ser Asp Tyr Cys Ile Gly Pro Asn Ala |     |     |
| 35 40 45                                                        |     |     |
| TCT ATC AAT GTC ATC ATG CAG CCC TTG GAG AAG ATG GCG CTA AAG GAG |     | 297 |
| Ser Ile Asn Val Ile Met Gln Pro Leu Glu Lys Met Ala Leu Lys Glu |     |     |
| 50 55 60                                                        |     |     |
| GCC CAC CAG CCG CAG ACC CAG CCC CTG TGG CAC CAG CTG GGA CTG GTC |     | 345 |
| Ala His Gln Pro Gln Thr Gln Pro Leu Trp His Gln Leu Gly Leu Val |     |     |
| 65 70 75                                                        |     |     |
| CTA GCT AAA CAC TTT GAA CCA CAG GAT GCC AAG GCC GTG CTG CAG CTG |     | 393 |
| Leu Ala Lys His Phe Glu Pro Gln Asp Ala Lys Ala Val Leu Gln Leu |     |     |
| 80 85 90                                                        |     |     |
| CTA AGG CAG GAG CAC GAR GAG CCC CTG CAG AAG ATA AGC CTG GAG CAC |     | 441 |
| Leu Arg Gln Glu His Glu Glu Arg Leu Gln Lys Ile Ser Leu Glu His |     |     |
| 95 100 105 110                                                  |     |     |
| CTG GAG CAG CTG GCC CAG TAC CTC CTG GCA GAG GAG CTC ACG TGG     |     | 486 |
| Leu Glu Gln Leu Ala Gln Tyr Leu Leu Ala Glu Glu Leu Thr Trp     |     |     |
| 115 120 125                                                     |     |     |

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 411 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

(v1) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: other

- (3) LOCATION: complement(195..411)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 99  
region 86..302  
id AA062591  
est

(A) NAME/KEY: sig peptide

- (B) LOCATION: 94..189  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.1  
seq RLSSLLLTMSNN/NP

CGCCG CCA TTTT GGA GTCTTCCCTA AGGATCCTCT ACCGGCTTTT CGAGTCAGTG 60

GGCGACGCCG CCATTTTGGA GTCTTCCTTA AGGATCCTCT ACCGGCTTTT CGAGTCAGTG 60

CTGCCGCCGC TSCCCGCGGC TTTGCAGAGC AGG ATG AAT GTG ATA GAC CAC GTG 114  
Met Asn Val Ile Asp His Val  
-30

CGG GAC ATG GCG GCC GCG GGG CTG CAC TCC AAC GTG CGG CTC CTC AGC 162  
Arg Asp Met Ala Ala Ala Gly Leu His Ser Asn Val Arg Leu Leu Ser  
-25 -20 -15 -10

AGC TTG TTA GTT ACA ATG AGT AAT AAC AAC CCT GAG TTA TTC TCC CCA 210  
Ser Leu Leu Leu Thr Met Ser Asn Asn Asn Pro Glu Leu Phe Ser Pro  
-5 1 5

CCT CAG AAG TAC CAG CTT TTG GTG TAT CAT GCA GAT TCT CTC TTT CAT 258  
Pro Gln Lys Tyr Gln Leu Leu Val Tyr His Ala Asp Ser Leu Phe His  
10 15 20

GAT AAG GAA TAT CGG AAT GCT GTS AST AAG TAT ACC ATG GCT TTA CAG 306  
Asp Lys Glu Tyr Arg Asn Ala Val Ser Lys Tyr Thr Met Ala Leu Gln  
25 30 35

CAG AAG AAG CCG CTA AGT AAA ACF TCA AAA GTG AGA CCT TCA ACT GGA 354  
Gln Lys Lys Ala Leu Ser Lys Thr Ser Lys Val Arg Pro Ser Thr Gly  
40 45 50 55



GTG AAA TAC 411  
Val Lys Tyr

(2) INFORMATION FOR SEQ ID NO: 257:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 232 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

- [illegible]

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 56..178  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.1  
seq RVLCPLLXAAAP/KR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 257:

|                                                                  |     |
|------------------------------------------------------------------|-----|
| AAGTAGCTCT CTAGGCCTGG GKRCGGGAGS GAGGGAGGCG GGCAGAGKWG GGGAG ATG | 58  |
|                                                                  | Met |
| GGC ACC CCC AST CTT TCC ATC CTC CTC ATA GGG GCA CCC GAA TCC CCT  | 106 |
| Gly Thr Pro Ser Leu Ser Ile Leu Leu Ile Gly Ala Pro Glu Ser Pro  |     |
| -40 -35 -30 -25                                                  |     |
| ATT CCT TAT TTC CCC TAT CAC TCA GGC ACT GGC AGG GTC CTT TGC CCA  | 154 |
| Ile Pro Tyr Phe Pro Tyr His Ser Gly Thr Gly Arg Val Leu Cys Pro  |     |
| -20 -15 -10                                                      |     |
| CTC CTG TWG GCC GCT GCG GCT CCA AAG CGA GAT GTG CCT GAG ACA GGT  | 202 |
| Leu Leu Xaa Ala Ala Ala Ala Pro Lys Arg Asp Val Pro Glu Thr Gly  |     |
| -3 1 5                                                           |     |
| TTG ACC ACC CAA CTG AAA AGA GAT CCT GGG                          | 232 |
| Leu Thr Arg Gln Leu Lys Asp His Pro Gly                          |     |
| 10 1                                                             |     |

## (2) INFORMATION FOR SEQ ID NO: 258:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 216 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 28..211
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 149..332  
id H15076  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 28..139
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 98  
region 147..258  
id R18367  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 138..179
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 258..299  
id R18367  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 46..123
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq HALFVLCCLLYAMS/HN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 258:

AAATAATTGA TTCCCTGTGT CTGGAATACC TGACCCTTCC TGGAT ATG GTG TAC CAC 57  
Met Val Tyr His  
-25

GCG CTC CAC ACC CCG GAT GAT GAT TAC CAT GCC CTG TTC GTG CTC TGC 105  
Ala Leu Asp Ser Pro Asp Asp Asp Tyr His Ala Leu Phe Val Leu Cys  
-20 -15 -10

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| CTC CTC TAT GCC ATG TCT CAT AAT AAA GGC ATG GAT CCT GAA AAA TTA | 153 |
| Leu Leu Tyr Ala Met Ser His Asn Lys Gly Met Asp Pro Glu Lys Leu |     |
| -5 1 5 10                                                       |     |
| GAG CGA ATC CAG CTC CCC GTG CCA AAT GCG GCC GAG AAG ACC ACC TAC | 201 |
| Glu Arg Ile Gln Leu Pro Val Pro Asn Ala Ala Glu Lys Thr Thr Tyr |     |
| 15 20 25                                                        |     |
| AAC CAC CCG CAT GGG                                             | 216 |
| Asn His Pro His Gly                                             |     |
| 30                                                              |     |

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

(v1) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(ix) FEATURE:

- ```
(A) NAME/KEY: other  
(B) LOCATION: complement(2..103)  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 98  
                        region 148..249  
                        id HSB379F042  
                        est
```

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: 32..73  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.8  
seg FIVLSMWLCCGFE/IL

(81) SEQUENCE DESCRIPTION: SEQ ID NO: 259:

AACTTGGTGA CTCTAGGTGA CTGGTCGACA G ATG TTC ATT GTA CTA TCA ATG 52  
Met Phe Ile Val Leu Ser Met  
-10

TGG CTT TGC TGT GGG TTT GAA ATT TTG CAA ACT AAG AGT TGG GTG GCA 100  
Trp Leu Cys Cys Gly Phe Glu Ile Leu Gln Thr Lys Ser Trp Val Ala  
-5 1 5

GGG 103  
Gly  
10

## (2) INFORMATION FOR SEQ ID NO: 260:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 351 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: cDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(184..281)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
region 2..99  
id T07232  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(103..170)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
region 113..180  
id T07232  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 42..106
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 29..84  
id AA099117  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 280..324
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.0  
seq VVILSSXVPLAAM/GV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 260:

```
AGAACCCCCC AAAATGCCCC ATTCTCGGG GCCACGCCCT GGGAGCCTG CAACCGCCCC 60
CACAAATTCT AGCAGTGCCA AGAAGAAGGA TAAAGAGTT CAAGGTAAGC AGTGTGAGGA 120
TCTCTTTAAG GAACATGTTT TTCTTCTTTC ATTACGTGCT TTTGGAGGAA GAAAAAACA 180
GGCCAGAGAA GGGG3CCTGT GCTTTACTT CCTTGTAGTC ACACCTGTGG GGAATCTGGG 240
TCTGGCCATC CCAGCCCTGB NCGGAGGGCT GTGTCAGGA ATG GTG GTG GTG ATT 294
Met Val Val Val Ile
```

-15

```

TTG AGC AGT GYA GTT CCC TTG GCA GCC ATG GGG GTC ATG GGC TGT GTC   342
Leu Ser Ser Xaa Val Pro Leu Ala Ala Met Gly Val Met Gly Cys Val
-10                -5                1                5

CGG GTG TGG                               351
Arg Val Trp

```

## (2) INFORMATION FOR SEQ ID NO: 261:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 201 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 16..62
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
region 463..509  
id AAC69619  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 1..45
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq AECSSLLHPSVRG/SI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 261:

```

ATG TTG GCA GAA TGC AGT TCC TTA CTG CAT CCA TCA GTT AGA GGC TCG   48
Met Leu Ala Glu Cys Ser Ser Leu Leu His Pro Ser Val Arg Gly Ser
-15                -10                -5                1

ATC CCA GAG GCC ACC TGC CGT GTC CTG CCA TGT GGC CCT CTC CAC AAC   96
Ile Pro Glu Ala Thr Cys Arg Val Leu Pro Cys Gly Pro Leu His Asn
          5                10                15

ATG GCA GTT TGC TCT TGC AAG GCT AGC AGG ACC TTC TAC TGC AAC TTC   144
Met Ala Val Cys Ser Cys Lys Ala Ser Arg Ser Phe Tyr Cys Asn Phe
          20                25                30

AGA TCT CTC CGA CTT GCT GTC TCT GAC TTC TTG ATT CTT TTC CAA AAG   192
Arg Ser Leu Arg Leu Ala Val Ser Asp Phe Leu Ile Leu Phe Gln Lys
          35                40                45

GGG TTA GGG                               201

```

Gly Leu Gly  
50

## (2) INFORMATION FOR SEQ ID NO: 262:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 146 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 76..141
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 90  
region 50..115  
id R25850  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 89..141
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 96  
region 49..101  
id N44651  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 85..141
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 91  
region 38..94  
id N31513  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 54..98
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq MARLLGLCAWARK/SV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 262:

AACTCTGTCA COTCCGCTGG AAGGAGTGGG ACCCAGACTT GCTGGTCTGA TCC ATG 56  
Met  
-15

CAG ATG GCC AGG CTC TTA GGC CTC TGT GGC TGG GCA CGG AAG TCG GTG 104

Gln Met Ala Arg Leu Leu Gly Leu Cys Ala Trp Ala Arg Lys Ser Val  
 -10 -5 1

CGG ATG GCC AGC TCC AGG ATG ACC CGC CGG GAC CCG CCA AGG  
 Arg Met Ala Ser Ser Arg Met Thr Arg Arg Asp Pro Pro Arg  
 5 10 15

146

## (2) INFORMATION FOR SEQ ID NO: 263:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 231 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(44..83)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 97  
 region 313..352  
 id R56475  
 est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(73..226)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 95  
 region 136..289  
 id TC5392  
 est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(73..226)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 93  
 region 161..314  
 id HUM030E12A  
 est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(72..226)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
 region 161..315  
 id HUM016H07A  
 est

## (ix) FEATURE:

- (A) NAME/KEY: other

-- (B) LOCATION: complement(131..226)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 97  
 region 168..213  
 id H08767  
 est

## (ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: complement(39..77)  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 92  
 region 326..364  
 id H08767  
 est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 91..219  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.8  
 seq LISVLYLIPKTLT/TN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 263:

```

AACAAAAGGA GAGTTTTATA ATTCACCTTA AAAGGAGATT TGATGGTAAA GTTTAAAGAT   60
TAAATATTT TGTTCTTCAA TTACAGAGCG ATG ACC CCA CAG TAT CTG CCT CAC   114
                               Met Thr Pro Gln Tyr Leu Pro His
                               -40

GGT GGA AAA TAC CAA GTT CTT GGA GAT TAC TCT TTG GCA GTG GTC TTC   162
Gly Gly Lys Tyr Gln Val Leu Gly Asp Tyr Ser Leu Ala Val Val Phe
-35                -30                -25                -20

CCC CTG CAC TTT TCT GAT CTA ATT TCT GTT TTA TAC CTT ATA CCC AAA   210
Pro Leu His Phe Ser Asp Leu Ile Ser Val Leu Tyr Leu Ile Pro Lys
-15                -10                -5

ACA CTT ACT ACC AAC AGC CGG   231
Thr Leu Thr Thr Asn Ser Arg
1

```

## (2) INFORMATION FOR SEQ ID NO: 264:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 361 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Spleen

## (ix) FEATURE:



(A) NAME/KEY: other  
(B) LOCATION: 53..342  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 19..308  
id C19012  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 123..349  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 112..338  
id AA058608  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 22..83  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 91  
region 12..73  
id AA058608  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 103..331  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 92  
region 87..315  
id N42002  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 19..87  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 97  
region 1..69  
id R13667  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 139..361  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 25..247  
id AA1310C9  
est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 17..85  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.7  
seq FLPLPKRAAFACRG/CQ

(ix) SEQUENCE DESCRIPTION: SEQ ID NO: 264:

```

AAGGGGGCGT GGGGCC ATG GTG GTC TTG CGG GCG GGG AAG AAG ACC TTT CTC   52
Met Val Val Leu Arg Ala Gly Lys Lys Thr Phe Leu
      -20                      -15

CCC CCT CTM WGC CGC GCC TTC GCC TGC CGC GGC TGT CAA CTC GCT CCG   100
Pro Pro Leu Xaa Arg Ala Phe Ala Cys Arg Gly Cys Gln Leu Ala Pro
      -10                      -5                      1                      5

GAG CGC GGC GCC GAG CGC AGG GAT ACA GCG CCC AGC GGG GTC TCA AGA   148
Glu Arg Gly Ala Glu Arg Arg Asp Thr Ala Pro Ser Gly Val Ser Arg
      10                      15                      20

TTC TGC CCT CCA AGA AAG TCT TGC CAT GAT TGG ATA GGA CCC CCA GAT   196
Phe Cys Pro Pro Arg Lys Ser Cys His Asp Trp Ile Gly Pro Pro Asp
      25                      30                      35

AAA TAT TCA AAC CTT CGA CCT GTT CAC TTT TAC ATA CCT GAA AAT GAA   244
Lys Tyr Ser Asn Leu Arg Pro Val His Phe Tyr Ile Pro Glu Asn Glu
      40                      45                      50

TCT CCA TTG GAA CAA AAG CTT AGA AAA TTA AGA CAA GAA ACA CAA GAA   292
Ser Pro Leu Glu Gln Lys Leu Arg Lys Leu Arg Gln Glu Thr Gln Glu
      55                      60                      65

TGG AAT CAA CAG TTC TGG GCA AAC CAG AAT TTG ACT TTT AGT AAG GAA   340
Trp Asn Gln Gln Phe Trp Ala Asn Gln Asn Leu Thr Phe Ser Lys Glu
      70                      75                      80                      85

AAA GAA GAA TTT ATT CAC TCA   361
Lys Glu Glu Phe Ile His Ser
      90

```

## (2) INFORMATION FOR SEQ ID NO: 265:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 11..113
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 99  
region 2..104  
id N76875  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide

(B) LOCATION: 15..74  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.6  
 seq AHLCSDSLPEQQ/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 265:

```

AAGAGAGAAC CGCC ATG AAG AGA GAA GGG GGT GCC GCC CAC CTC TGC TCC      50
      Met Lys Arg Glu Gly Gly Ala Ala His Leu Cys Ser
      -20                      -15                      -10

GAC ASC CTC CCG GAG TCC CAG CAG CAA GAC GGC AAC CAC GCA CCC AAC      93
Asp Ser Leu Pro Glu Ser Gln Gln Asp Gly Asn His Ala Pro Asn
      -5                      1                      5

TTC TCC AGC CAC GGC                                          113
Phe Ser Ser His Gly
      10
  
```

(2) INFORMATION FOR SEQ ID NO: 266:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 342 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: other  
 (B) LOCATION: 255..343  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 93  
 region 12..100  
 id AA026923  
 est

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 205..327  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.6  
 seq PYSLAACFGSSQG/GV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 266:

```

ACCGAGAAGC CCTCACAGAT GCAGATGACT TTGGCCTACA GTTCCCGGTG GACCTGGATG      60
TGAGGGTGAA GGCTGTGCTG CTGGGAGCCA CATTCTCAT TGACTACATG TTCTTTGAGA      120
ACCGAGGAGG CCTGGGCCC TCTGCCATCA CCAGTTAGAG GCCACCATGG TGTGAGGAGA      180
  
```

```

CCATCACCTG GACCAGAACT CCAG ATG CTC ACC TGC CCT GGC CCC TCC TCT      231
               Met Val Thr Cys Pro Gly Pro Ser Ser
               -40                               -35

GGG CAG CCC CTT TCC TCC ATG TAC ACT GCA GGG GAC AGA AGG GGG GCC      279
Gly Gln Pro Leu Ser Ser Met Tyr Thr Ala Gly Asp Arg Arg Gly Ala
               -30                               -25                               -20

CCA TCC CTA CCC TAC TCC CTG GCC GCC TGC CCC TGT GGT TCC CAA GGA      327
Pro Ser Leu Pro Tyr Ser Leu Ala Ala Cys Pro Cys Gly Ser Gln Gly
               -15                               -10                               -5

GGG GTA TGT ATG AGA                                          342
Gly Val Cys Met Arg
  1                               5

```

## (2) INFORMATION FOR SEQ ID NO: 267:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 420 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(1..300)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..300  
id H13499  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(1..268)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 100  
region 1..268  
id W40371  
est

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: complement(1..93)
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 1..93  
id H04223  
est

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide

- (B) LOCATION: 109..162  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
 seq ALEVIVTLSETAA/AM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 267:

```

AAATCCCTCG TTGAGATTGC AGATACTGTT CCAAAGTATT TGGCTCCTCA CTTGGAAGCA    60
ACTCTACAGC TAAGTCTAAA GTTGTGTGGA GACACTAGCC TCAACAAT ATG CAA CGC    117
                                     Met Gln Arg
CAG CTT GCC CTT GAA GTG ATC GTC ACC CTC TCT GAG ACT GCA GCT GCT    165
Gln Leu Ala Leu Glu Val Ile Val Thr Leu Ser Glu Thr Ala Ala Ala
-15                               -10                               -5                               1
ATG TTA AGA AAA CAT ACC AAT ATT GTT GCA CAG ACT ATT CCT CAG ATG    213
Met Leu Arg Lys His Thr Asn Ile Val Ala Gln Thr Ile Pro Gln Met
                    5                               10                               15
TTA GCA ATG ATG GTT GAT TTG GAA GAA GAT GAG GAC TGG GCA AAT GCA    261
Leu Ala Met Met Val Asp Leu Glu Glu Asp Glu Asp Trp Ala Asn Ala
                20                               25                               30
GAT GAA CTA GAA GAT GAT GAT TTT GAC AGC AAT GCA GTT GCA GGC GAG    309
Asp Glu Leu Glu Asp Asp Asp Phe Asp Ser Asn Ala Val Ala Gly Glu
                35                               40                               45
AGT GCT CTA GAT CGA ATG GCT TGC GGA CTT GGT GGA AAG CTC GTT CTG    357
Ser Ala Leu Asp Arg Met Ala Cys Gly Leu Gly Gly Lys Leu Val Leu
                50                               55                               60                               65
CCG ATG ATC AAG GAA CAC ATT ATG CAA ATG CTT CAA AAT CGT AAG CTG    405
Pro Met Ile Lys Glu His Ile Met Gln Met Leu Gln Asn Arg Lys Leu
                70                               75                               80
TGT CCT TCA ATG CTA
Cys Pro Ser Met Leu
                85

```

(2) INFORMATION FOR SEQ ID NO: 268:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 392 base pairs  
 (B) TYPE: NUCLEIC ACID  
 (C) STRANDEDNESS: DOUBLE  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: CDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: other  
 (B) LOCATION: 177..348  
 (C) IDENTIFICATION METHOD: blastn

-- (D) OTHER INFORMATION: identity 100  
region 266..437  
id N32722  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 52..175  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 94  
region 142..265  
id N32722  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 3..41  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 1..39  
id N32722  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 36..387  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 96  
region 12..363  
id W32042  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 177..348  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 134..305  
id R55254  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 99..175  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 100  
region 57..133  
id R55254  
est

(ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 44..102  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 91  
region 1..53  
id R55254  
est

(ix) FEATURE:

(A) NAME/KEY: other

.. (B) LOCATION: 356..387  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 90  
 region 315..346  
 id R55254  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: 177..334  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 100  
 region 149..306  
 id W37647  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: 38..175  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 94  
 region 11..148  
 id W37647  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: 38..174  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 95  
 region 8..144  
 id R50622  
 est

(ix) FEATURE:  
 (A) NAME/KEY: other  
 (B) LOCATION: 174..295  
 (C) IDENTIFICATION METHOD: blastn  
 (D) OTHER INFORMATION: identity 99  
 region 143..264  
 id R50622  
 est

(ix) FEATURE:  
 (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 147..374  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.5  
 seq LASASELP LGSRP/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 268:

AACTTCCTGT GATCCCGGCG GTGACACCG CAACATGGCC CGTGAACGGA GCTGAAGTCG	60
ACGACTTCTC CTGGRARMCC CCGACTGAGG CGGAGACGAA GGTGCTGCAG GCGCGACGGG	120
ATGGGCAAGA TGGCATCTCC CGGCTC ATG GGC GAC TAT CTG CTG CGC GGT TAC	173
Met Gly Asp Tyr Leu Leu Arg Gly Tyr	
-75 -70	

CGC ATG CTG GGC GAG ACG TGT GCG GAC TGC GGG ACG ATC CTC CTC CAA	221
Arg Met Leu Gly Glu Thr Cys Ala Asp Cys Gly Thr Ile Leu Leu Gln	
-65 -60 -55	
GAC AAA CAG CGG AAA ATC TAC TGC GTG GCT TGT CAG GAA CTC GAC TCA	269
Asp Lys Gln Arg Lys Ile Tyr Cys Val Ala Cys Gln Glu Leu Asp Ser	
-50 -45 -40	
GAC GTG GAT AAA GAT AAT CCC GCT CTG AAT GCC CAG GCT GCC CTC TCC	317
Asp Val Asp Lys Asp Asn Pro Ala Leu Asn Ala Gln Ala Ala Leu Ser	
-35 -30 -25 -20	
CAA GCT CGG GAG CAC CAG CTG GCC TCA GCC TCA GAG CTC CCC CTG GGC	365
Gln Ala Arg Glu His Gln Leu Ala Ser Ala Ser Glu Leu Pro Leu Gly	
-15 -10 -5	
TCT CGA CCT GCG CCC CAA CCC CAC GGG	392
Ser Arg Pro Ala Pro Gln Pro His Gly	
1 5	

## (2) INFORMATION FOR SEQ ID NO: 269:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 234 base pairs
- (B) TYPE: NUCLEIC ACID
- (C) STRANDEDNESS: DOUBLE
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 61..232
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 100  
region 1..172  
id HSC1R  
vrt

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 72..232
- (C) IDENTIFICATION METHOD: fasta
- (D) OTHER INFORMATION: identity 100  
region 24..184  
id HUMC1R  
vrt

## (ix) FEATURE:

- (A) NAME/KEY: other
- (B) LOCATION: 109..232
- (C) IDENTIFICATION METHOD: blastn
- (D) OTHER INFORMATION: identity 92  
region 1..124  
id T74375



est

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 98..141  
(C) IDENTIFICATION METHOD: blastn  
(D) OTHER INFORMATION: identity 93  
region 1..44  
id T64778  
est

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: 112..156  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 8.1  
seq LLYLLVPALFCRA/GG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 269:

```
AACTCCACAG AAAACCCCTCC CCTCCCTGCT GTGCATGACG CGGGCTCCCT CTGCACACAG      60
TGACACGAAGA CGCTGTCGGG AGAGCCCAGG ATTCAACACG GGCCTTGAGA A ATG TGG      117
                                         Met Trp
                                         -15

CTC TTG TAC CTC CTG GTG CCG GCC CTG TTC TGC AGG GCA GGA GGC TCC      165
Leu Leu Tyr Leu Leu Val Pro Ala Leu Phe Cys Arg Ala Gly Gly Ser
      -10                      -5                      1

ATT CCC ATC CCT CAG AAG TTA TTT GGG GAG GTG ACT TCC CCT CTG TTC      213
Ile Pro Ile Pro Gln Lys Leu Phe Gly Glu Val Thr Ser Pro Leu Phe
      5                      10                      15

CCC AAG CCT TAC CCC AAC ACG      234
Pro Lys Pro Tyr Pro Asn Thr
      20                      25
```

## (2) INFORMATION FOR SEQ ID NO: 270:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 302 base pairs  
(B) TYPE: NUCLEIC ACID  
(C) STRANDEDNESS: DOUBLE  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: CDNA

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: other  
(B) LOCATION: 87..300  
(C) IDENTIFICATION METHOD: fasta  
(D) OTHER INFORMATION: identity 98  
region 1..214

239

id HSCALICIN  
vrt

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: 78..251
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq LAAVSPLVRS LIS/SN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 270:

```

AAATCTGATC CCACAGGCCT GAGAAAGTCT GCTCTCCAGW ACCTGCTGCT GATCTGTTTC    60
AGCCGACAAG AGGCACC ATG AAA TTG GAA TTC ACG GAG AAA AAC BAC RAT    110
           Met Lys Leu Glu Phe Thr Glu Lys Asn Xaa Xaa
           -55                               -50

AGC TTC GTG CTG CAR AAC CTG AAC AGA CAG AGG AAA CGC AAA GAG TAC    158
Ser Phe Val Leu Gln Asn Leu Asn Arg Gln Arg Lys Arg Lys Glu Tyr
          -45                               -40                               -35

TGG GAC ATG GCC CTG AGT GTG GAC AAC CAC GTC TTC TTT GCA CAT CGC    206
Trp Asp Met Ala Leu Ser Val Asp Asn His Val Phe Phe Ala His Arg
          -30                               -25                               -20

AAT GTG CTG GCT GCT GTC TCC CCA CTG GTG AGG AGC CTC ATC TCC AGC    254
Asn Val Leu Ala Ala Val Ser Pro Leu Val Arg Ser Leu Ile Ser Ser
          -15                               -10                               -5                               1

AAT GAC ATG AAG ACC GCT GAT GAG CTT TTC ATC ACC ATT GAC ACC AAG    302
Asn Asp Met Lys Thr Ala Asp Glu Leu Phe Ile Thr Ile Asp Thr Lys
           5                               10                               15

```

## (2) INFORMATION FOR SEQ ID NO: 271:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 13.2  
seq LLLLSTLVIPSAA/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 271:

Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val

```

-30          -25          -20          -15
Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro
      -10          -5          1
Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu
      5          10          15
Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu
      20          25          30
Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly
      35          40          45          50
Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly
      55          60          65
Asn Asn Thr Leu Ser Ser Xaa Leu Gln Ile Asp Xaa Met Thr Asp Asn
      70          75          80
Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala
      85          90          95

```

## (2) INFORMATION FOR SEQ ID NO: 272:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 12  
seq VLVLCVLLLQAQG/GY

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 272:

```

Met Ala Pro Gln Thr Leu Leu Pro Val Leu Val Leu Cys Val Leu Leu
-20          -15          -10
Leu Gln Ala Gln Gly Gly Tyr Arg Asp Lys Met Arg Met Gln Arg Ile
-5          1          5          10
Lys Val Cys Glu Lys Arg Pro Ser Ile Asp Leu Cys Ile His His Arg
      15          20          25

```

## (2) INFORMATION FOR SEQ ID NO: 273:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 101 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 11  
seq SLVLLCLTCSYA/FM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 273:

```

Met Trp Thr Leu Lys Ser Ser Leu Val Leu Leu Leu Cys Leu Thr Cys
      -15                      -10                      -5

Ser Tyr Ala Phe Met Phe Ser Ser Leu Arg Gln Lys Thr Ser Glu Pro
      1                      5                      10

Gln Gly Lys Val Gln Tyr Gly Glu His Phe Arg Ile Arg Gln Asn Leu
      15                      20                      25

Pro Glu His Thr Gln Gly Trp Leu Gly Ser Lys Trp Leu Trp Leu Leu
      30                      35                      40                      45

Xaa Val Val Val Pro Phe Val Ile Leu Gln Cys Gln Arg Asp Ser Glu
      50                      55                      60

Lys Asn Lys Glu Gln Ser Pro Pro Gly Leu Arg Gly Gly Gln Leu His
      65                      70                      75

Ser Pro Leu Lys Lys
      80

```

## (2) INFORMATION FOR SEQ ID NO: 274:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 115 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 10.6  
seq LLLPLLWGGSQ/EK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 274:

```

Met Leu Pro Leu Leu Leu Leu Pro Leu Leu Trp Gly Gly Ser Leu Gln
-15                      -10                      -5

Glu Lys Pro Val Tyr Glu Leu Gln Val Gln Lys Ser Val Thr Val Gln
1                      5                      10                      15

Glu Gly Leu Cys Val Leu Val Pro Cys Ser Phe Ser Tyr Pro Trp Arg
20                      25                      30

Ser Trp Tyr Ser Ser Pro Pro Leu Tyr Val Tyr Trp Phe Arg Asp Gly
35                      40                      45

Glu Ile Pro Tyr Tyr Ala Glu Val Val Ala Thr Asn Asn Pro Asp Arg
50                      55                      60

Arg Val Lys Pro Glu Thr Gln Gly Arg Phe Arg Leu Leu Gly Asp Val
65                      70                      75                      80

Gln Lys Lys Asn Cys Ser Leu Ser Ile Gly Asp Xaa Arg Met Glu Asp
85                      90                      95

Thr Gly Gly

```

(2) INFORMATION FOR SEQ ID NO: 275:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.4  
seq LLLLLCGPSQDQC/RP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 275:

```

Met Glu Thr Gly Ala Leu Arg Arg Pro Gln Leu Leu Pro Leu Leu Leu
-25                      -20                      -15

Leu Leu Cys Gly Pro Ser Gln Asp Gln Cys Arg Pro Val Leu Gln Asn
-10                      -5                      1                      5

Leu Leu Gln Ser Pro Gly Leu Thr Trp Ser Leu Glu Val Pro Thr Gly

```

Arg Glu Gly Lys Glu Gly Thr Met Arg Val Ser Pro Thr Ala Pro Arg  
 25 10 15 20 30 35

## (2) INFORMATION FOR SEQ ID NO: 276:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 93 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.1  
seq LVLTLCTLPLAVA/SA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 276:

Met Glu Arg Leu Val Leu Thr Leu Cys Thr Leu Pro Leu Ala Val Ala  
 -15 -10 -5

Ser Ala Gly Cys Ala Thr Thr Pro Ala Arg Asn Leu Ser Cys Tyr Gln  
 1 5 10 15

Cys Phe Lys Val Ser Ser Trp Thr Glu Cys Pro Pro Thr Trp Cys Ser  
 20 25 30

Pro Leu Asp Gln Val Cys Ile Ser Asn Glu Val Val Val Ser Phe Lys  
 35 40 45

Trp Ser Val Arg Val Leu Leu Ser Lys Arg Cys Ala Pro Arg Cys Pro  
 50 55 60

Asn Ser Gly  
 65

## (2) INFORMATION FOR SEQ ID NO: 277:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 120 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens



... (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -23...-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 9.8  
 seq FLLFFFLFLLTRG/SL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 277:

```

Met Met Leu Pro Gln Trp Leu Leu Leu Leu Phe Leu Leu Phe Phe Phe
      -20                      -15                      -10

Leu Phe Leu Leu Thr Arg Gly Ser Leu Ser Pro Thr Lys Tyr Asn Leu
      -5                      1                      5

Leu Glu Leu Lys Glu Xaa Xaa Xaa Gly Asn Gln Asp Cys Glu Thr Gly
      10                      15                      20                      25

Cys Cys Gln Arg Ala Pro Asp Asn Cys Glu Ser His Cys Ala Glu Lys
      30                      35                      40

Gly Ser Glu Gly Ser Leu Cys Gln Thr Gln Val Phe Phe Gly Gln Tyr
      45                      50                      55

Arg Ala Cys Pro Cys Leu Arg Asn Leu Thr Cys Ile Tyr Ser Lys Asn
      60                      65                      70

Glu Lys Trp Leu Ser Ile Ala Tyr Gly Arg Cys Gln Lys Ile Gly Arg
      75                      80                      85

Gln Lys Leu Ala Arg Lys Cys Ser
      90                      95
  
```

(2) INFORMATION FOR SEQ ID NO: 278:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -22...-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 9.3  
 seq LVVFCLALQLVPG/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 278:



Met Lys Pro Val Leu Pro Leu Gln Xaa Leu Val Val Phe Cys Leu Ala  
           -20                          -15                          -10

Leu Gln Leu Val Pro Gly Ser Pro Lys Gln Leu Gly  
       -5                                  1                          5

## (2) INFORMATION FOR SEQ ID NO: 279:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -35..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.5  
seq LFFSLFSAPLASA/VR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 279:

Met Phe Arg Gln Arg Gln Glu Thr Ala Gln Arg Ser Thr Gln Ser Cys  
   -35                          -30                          -25                          -20

Arg Cys Pro Arg Asp Gly Leu Phe Phe Ser Leu Phe Ser Ala Pro Leu  
                           -15                          -10                          -5

Ala Ser Ala Val Arg Ala Ala Xaa  
                           1                          5

## (2) INFORMATION FOR SEQ ID NO: 280:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.4  
seq RLLALPLALVLC/FE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 280:

Met Gly Ser Ser Ala Cys Glu Ile Ala Val Gly Thr Lys Arg Leu Leu  
 -25 -20 -15  
 Leu Ala Leu Pro Leu Ala Leu Val Leu Gly Phe Glu Gly Ser Ser Val  
 -10 -5 1 5  
 Pro Pro Arg Asn Phe  
 10

## (2) INFORMATION FOR SEQ ID NO: 281:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.4  
seq SLLFICFFGESFC/IC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 281:

Met Ser Asn Gln Arg Leu Pro Leu Ile Phe Ser Leu Leu Phe Ile Cys  
 -20 -15 -10  
 Phe Phe Gly Glu Ser Phe Cys Ile Cys Asp Gly Thr Val Trp Thr Xaa  
 -5 1 5  
 Val Xaa Trp Glu Ile Leu Pro Glu Glu Val His Tyr Trp Lys Val Lys  
 10 15 20 25  
 Gly Ser Pro Ser His Cys Leu Arg  
 30

## (2) INFORMATION FOR SEQ ID NO: 282:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -16..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 9.2  
seq FLSFLLALLSLNC/IP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 282:

Met Leu Trp Phe Leu Ser Phe Leu Leu Ala Leu Leu Ser Leu Asn Cys  
-15 -10 -5

Ile Pro Ile Gly  
1

## (2) INFORMATION FOR SEQ ID NO: 283:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -24..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 9.2  
seq ICCVIVLISLSWT/SP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 283:

Met Leu Xaa Ile Ser Leu Glu Ile Xaa Ser Phe Ile Cys Cys Val Ile  
-20 -15 -10

Val Leu Ile Ser Leu Ser Trp Thr Ser Pro Phe Thr Gly Val Tyr Leu  
-5 1 5

Ile Gly Leu Ile Ile Glu Pro Gly  
10 15

## (2) INFORMATION FOR SEQ ID NO: 284:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids  
(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -19..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 9.2  
seq ILFILTFFSHTFC/SR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 284:

Met Val Phe Arg Asn Cys Ile Leu Phe Ile Leu Thr Phe Phe Ser His  
-15 -10 -5

Thr Phe Cys Ser Arg Gln Asn Lys Ala Gln Pro Trp  
1 5

(2) INFORMATION FOR SEQ ID NO: 285:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -13..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 9.1  
seq MLAA CPLSPGCQS/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 285:

Met Leu Ala Ala Cys Pro Leu Ser Pro Gly Cys Gln Ser Ala Pro Ser  
-10 -5 1

Thr Trp Asn His Phe Pro Pro Glu Arg Ile Thr Thr Gly Ala Gly Ser  
5 10 15

Leu Leu Lys Pro Gly Gly Gly Leu Trp Pro Arg Thr Val Ser Leu Pro  
20 25 30 35

Ser Pro Ala

## (2) INFORMATION FOR SEQ ID NO: 286:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 76 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1  
seq FLTLITHCTVSWA/QS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 286:

Met Ala Trp Ser Pro Leu Phe Leu Thr Leu Ile Thr His Cys Thr Val  
                    -15                    -10                    -5

Ser Trp Ala Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Glu Ala  
                    1                            5                            10

Pro Arg Gln Arg Val Thr Ile Ser Cys Phe Gly Ser Ser Ser Asn Ile  
            15                            20                            25

Gly Arg Asn Ala Val Asn Trp Tyr Gln Gln Leu Pro Gly Arg Ser Pro  
            30                            35                            40                            45

Arg Leu Leu Ile Phe Tyr Asn Asn Leu Pro Ala Ser  
                    50                            55

## (2) INFORMATION FOR SEQ ID NO: 287:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -13..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.1  
seq LVSLCSWSPPPLTS/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 287:

Met Leu Lys Ser Val Leu Val Ser Leu Cys Ser Trp Ser Pro Pro Leu  
           -15                          -10                          -5  
 Thr Ser Ser Pro Arg  
                   1

(2) INFORMATION FOR SEQ ID NO: 288:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 44 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -22...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9  
seq FILAALSLSTTFS/LQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 288:

Met Thr Ser Lys Xaa Ile Leu Val Ser Phe Ile Leu Ala Ala Leu Ser  
           -20                          -15                          -10  
 Leu Ser Thr Thr Phe Ser Leu Gln Pro Tyr Gln Gln Lys Val Leu Leu  
           -5                          1                          5                          10  
 Val Ser Phe Asp Gly Phe Arg Trp Asp Tyr Leu Tyr  
                   15                          20

(2) INFORMATION FOR SEQ ID NO: 289:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20...-1

(D) OTHER INFORMATION: score 3.9

seq LAVXLGLATAVSA/GP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 289:

Met Lys Ser Leu Ser Leu Xaa Leu Ala Val Xaa Leu Gly Leu Ala Thr  
-20 -15 -10 -5

Ala Val Ser Ala Gly Pro Ala Trp  
1

(2) INFORMATION FOR SEQ ID NO: 290:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig peptide

(B) LOCATION: -21...-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 8.8

seq LLWALLEMQSLWP/QL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 290:

Met Trp Ala Met Glu Ser Gly His Leu Leu Trp Ala Leu Leu Phe Met  
-20 -15 -10

Gln Ser<sup>1</sup> Leu Trp Pro Gln Leu Thr Asp Gly Ala Thr Arg Val Tyr Tyr  
-5 1 5 10

Leu Gly Ile Arg Asp Val Gln Trp Asn Tyr Ala Pro Lys Gly Arg Asn  
15 20 25

Val Ile Thr Asn Gln Pro Leu Asp Ser Asp Ile Val Ala Ser Ser Phe  
30 35 40

Leu Lys Ser Asp Lys Asn Arg Ile Gly Gly Thr Thr Arg Arg Pro Trp  
45 50 55

(2) INFORMATION FOR SEQ ID NO: 291:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 49 amino acids

(3) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -20..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 9.8  
seq LLVMGSLPSASWS/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 291:

Met Ala Gln Thr Trp Ala Xaa Leu Leu Val Met Gly Ser Leu Pro Ser  
-20 -15 -10 -5

Ala Ser Trp Ser Leu Pro Cys Leu Ser Trp Glu Ser Leu Leu Lys Ala  
1 5 10

Ala Ala Cys Ser Glu Leu Asp Gly Arg Asn Val Gly Asn Thr Pro Thr  
15 20 25

Arg

(2) INFORMATION FOR SEQ ID NO: 292:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -22..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 8.7  
seq LITLLYVWPVINA/CQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 292:

Met Lys Cys Gly Phe Leu Ala Tyr Leu Leu Ile Thr Leu Leu Tyr Val  
-20 -15 -10

Trp Pro Val Ile Asn Ala Cys Gln  
-5 1



## (2) INFORMATION FOR SEQ ID NO: 293:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 36 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: -25..-1
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 8.5  
seq LKVLLLPLAPAAA/QD
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 293:

```

Met Arg Lys Pro Ala Ala Gly Phe Leu Pro Ser Leu Leu Lys Val Leu
-25                -20                -15                -10

Leu Leu Pro Leu Ala Pro Ala Ala Ala Gln Asp Ser Thr Gln Ala Ser
                -5                1                5

Thr Pro Gly Arg
                10

```

## (2) INFORMATION FOR SEQ ID NO: 294:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 54 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Spleen
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: -17..-1
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 8.4  
seq LLFLTSVV?FVLA/PR
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 294:

```

Met Arg Gln Ser Leu Leu Phe Leu Thr Ser Val Val Pro Phe Val Leu
-15                -10                -5

Ala Pro Arg Pro Pro Asp Asp Pro Gly Phe Gly Pro His Gln Arg Leu

```

1                      5                      10                      15  
 Glu Lys Leu Asp Ser Leu Leu Ser Asp Tyr Asp Ile Leu Ser Leu Ser  
                     20                      25                      30  
 Asn Ile Gln Gln Gln Xaa  
                     35

## (2) INFORMATION FOR SEQ ID NO: 295:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -25..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.1  
seq SVLLGLLALMATA/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 295:

Met Glu Leu Ser Gln Met Ser Glu Leu Met Gly Leu Ser Val Leu Leu  
 -25                      -20                      -15                      -10  
 Gly Leu Leu Ala Leu Met Ala Thr Ala Ala Val Ala Arg Gly Trp Leu  
                     -5                      1                      5  
 Arg Ala Gly Glu Val Arg  
                     10

## (2) INFORMATION FOR SEQ ID NO: 296:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 103 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -65..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

Gly Leu Gly Phe Ile Ile Ser Ser Arg Thr Arg Ser Glu Leu Pro His  
                   -45                                  -40                                  -35  
 Leu Val Ser Trp His Ser Trp Val Gly Ala Leu Thr Leu Leu Ala Thr  
                   -30                                  -25                                  -20  
 Ala Val Gln Ala Leu Cys Gly Leu Cys Leu Leu Cys Pro Arg Ala Ala  
                   -15                                  -10                                  -5  
 Arg Val Ser Arg Val Ala Arg Leu Lys Leu Tyr His Leu Thr Cys Gly  
   1                                  5                                  10                                  15  
 Leu Val Val Tyr Leu Met Ala Thr Val Thr Val Leu Leu Gly Met Tyr  
                   20                                  25                                  30  
 Ser Val Trp Phe  
                   35

## (2) INFORMATION FOR SEQ ID NO: 298:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 100 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -57..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq LLHRLASFHRVWS/FP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 298:

Met Leu Arg Phe Pro Thr Cys Phe Pro Ser Xaa Arg Val Xaa Gly Xaa  
                   -55                                  -50                                  -45  
 Lys Gln Leu Pro Gln Glu Ile Ile Xaa Leu Val Trp Ser Pro Xaa Arg  
                   -40                                  -35                                  -30  
 Asp Xaa Ile Xaa Leu Ala Asn Thr Ala Gly Glu Val Leu Leu His Arg  
                   -25                                  -20                                  -15                                  -10  
 Leu Ala Ser Phe His Arg Val Trp Ser Phe Pro Pro Asn Glu Asn Thr  
                   -5                                  1                                  5  
 Gly Xaa Glu Val Thr Cys Leu Ala Trp Arg Pro Asp Gly Lys Leu Leu  
                   10                                  15                                  20  
 Ala Phe Ala Leu Ala Asp Thr Lys Lys Ile Val Leu Cys Asp Val Glu  
                   25                                  30                                  35

Lys Pro Glu Ser  
40

(2) INFORMATION FOR SEQ ID NO: 299:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 130 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -42..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq LALVVALVAERFA/RR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 299:

```

Met Phe Met Val Leu Glu Val Val Val Ser Arg Val Thr Ser Ser Leu
  -40                      -35                      -30

Ala Met Leu Ser Asp Ser Phe His Met Leu Ser Asp Val Leu Ala Leu
  -25                      -20                      -15

Val Val Ala Leu Val Ala Glu Arg Phe Ala Arg Arg Thr His Ala Thr
  -10                      -5                      1                      5

Gln Lys Asn Thr Phe Gly Trp Ile Arg Ala Glu Val Met Gly Ala Leu
      10                      15                      20

Val Asn Ala Ile Phe Leu Thr Gly Leu Cys Phe Ala Ile Leu Leu Glu
      25                      30                      35

Ala Ile Glu Arg Phe Ile Glu Pro His Glu Met Gln Gln Pro Leu Val
      40                      45                      50

Val Xaa Trp Gly Arg Ala Trp Xaa Ala Ala Gly Gln Arg Ala Gly Ala
      55                      60                      65                      70

Leu Pro Leu Pro Pro Ser Gln Arg Leu Gln Pro Gly Leu Arg Pro Arg
      75                      80                      85
Pro Trp

```

(2) INFORMATION FOR SEQ ID NO: 300:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
- (B) TYPE: AMINO ACID

.. (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -39..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.5  
seq LLLLLGLIVLVNI/GI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 300:

```

Met Glu Asn Gln Leu Trp His Asn Thr Val Arg Cys Cys Asn Gln Tyr
      -35                      -30                      -25

Gln Glu Ser Pro His Asp Ala Glu Asp Ile Leu Leu Leu Leu Gly
      -20                      -15                      -10

Leu Ile Val Leu Val Asn Ile Gly Ile Asn Val Ala Thr Met Met Trp
      -5                      1                      5

His Gly Leu Gln Asn Ala Leu Asp Lys Met Ile Asp Trp Ala Thr Gln
      10                      15                      20                      25

Lys Ile Ala Val Phe Phe Ala Val Phe Val Ala Ala Ala Ala Arg
      30                      35                      40

```

(2) INFORMATION FOR SEQ ID NO: 301:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 52 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -18..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.5  
seq ITLLTLSPNSVCC/CP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 301:

```

Met Leu Ser Xaa Lys Ile Thr Leu Leu Thr Leu Ser Pro Asn Ser Val
      -15                      -10                      -5

```

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -18..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.4  
seq GNLVLCVLAISLA/SM

Met Glu Gly Pro Arg Gly Trp Leu Val Leu Cys Val Leu Ala Ile Ser  
-15 -10 -5

Leu Ala Ser Met Val Thr Glu Asp Leu Cys Arg Ala Pro Asp Gly Lys  
1 5 10

Lys Gly Glu Ala Gly Xaa Pro Gly Arg Arg Gly Arg Pro Gly Leu Lys  
15 20 25 30

Gly Glu Gln Arg

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig peptide

(B) LOCATION: -20..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 7.3  
 seq LAVFMLLAQLVSG/NW

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 303:

```

Met Lys Ser Leu Leu Phe Thr Leu Ala Val Phe Met Leu Leu Ala Gln
-20          -15          -10          -5

Leu Val Ser Gly Asn Trp Tyr Val Lys Lys Cys Leu Asn Asp Val Gly
          1          5          10

Ile Cys Lys Lys Lys Cys Lys Pro Glu Glu Met His Val Lys Asn Gly
          15          20          25

Trp Ala Met Cys Gly Lys Gln Arg Asp Cys Cys Val Pro Ala Asp Arg
          30          35          40

Arg Ala Asn Tyr Pro Val Phe Cys Val Gln Thr Lys Thr Thr Arg Ile
          45          50          55          60

Ser Thr Val Thr Ala Thr Thr Ala Thr Thr Thr Leu Met Met Thr Thr
          65          70          75

Ala Ser Met Ser Ser Met Ala Pro Thr Arg Phe Ser
          80          85
  
```

(2) INFORMATION FOR SEQ ID NO: 304:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -16..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 7.3  
 seq LILLFSLISIVC/MI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 304:

```

Met Leu Lys Leu Ile Leu Leu Phe Ser Leu Leu Ile Ser Ile Val Cys
-15          -10          -5

Met Ile
1
  
```

## (2) INFORMATION FOR SEQ ID NO: 305:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1  
seq LASLQWSLTAWC/GS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 305:

Met Thr Pro Trp Cys Leu Ala Cys Leu Gly Arg Arg Pro Leu Ala Ser  
-25 -20 -15

Leu Gln Trp Ser Leu Thr Leu Ala Trp Cys Gly Ser Gly Ser His Trp  
-10 -5 1 5

Thr Glu

## (2) INFORMATION FOR SEQ ID NO: 306:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9  
seq LWVLLLCALVVVTL/LV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 306:

Met Thr Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val  
-25 -20 -15

Leu Leu Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro



```
(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 35 amino acids
      (B) TYPE: AMINO ACID
      (C) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:
      (A) ORGANISM: Homo Sapiens
      (F) TISSUE TYPE: Testis

(ix) FEATURE:
      (A) NAME/KEY: sig_peptide
      (B) LOCATION: -20...-1
      (C) IDENTIFICATION METHOD: Von Heijne matrix
      (D) OTHER INFORMATION:  score 6.9
                               seq LFCATLSCMPATS/AP
```

```

Met Thr Gly Asn Asn Arg Asp Leu Phe Cys Ala Thr Leu Ser Cys Met
-20                      -15                      -10                      -5

Pro Ala Thr Ser Ala Pro His Met Lys Leu Pro Asp Ile Ser Phe His
          1                      5                      10

Leu Pro Gly
          15

```

```
(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 37 amino acids
      (B) TYPE: AMINO ACID
      (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:
      (A) ORGANISM: Homo Sapiens
      (F) TISSUE TYPE: Ovary

(ix) FEATURE:
      (A) NAME/KEY: sig_peptide
      (B) LOCATION: -26..-1
      (C) IDENTIFICATION METHOD: Von Heijne matrix
```

(D) OTHER INFORMATION: score 6.9  
seq LWVLLLCALVVTL/LV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 308:

```

Met Thr Met Arg His Asn Trp Thr Pro Asp Leu Ser Pro Leu Trp Val
-25                      -20                      -15

Leu Leu Leu Cys Ala His Val Val Thr Leu Leu Val Arg Ala Thr Pro
-10                      -5                      1                      5

Val Ser Gln Pro Thr
10

```

(2) INFORMATION FOR SEQ ID NO: 309:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.9  
seq LYLLGMLVPGGLG/YD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 309:

```

Met Lys Pro Leu Leu Glu Thr Leu Tyr Leu Leu Gly Met Leu Val Pro
-20                      -15                      -10                      -5

Gly Gly Leu Gly Tyr Asp Arg Ser Leu Ala Gln His Arg Gln Glu Ile
1                      5                      10

Val Asp Lys Ser Val Ser Pro Trp Ser Leu Glu Thr Tyr Ser Tyr Asn
15                      20                      25

Ile Tyr His Pro Met Gly Glu Ile Tyr Glu Trp Met Arg Glu Ile Ser
30                      35                      40

Glu Lys Tyr Lys Glu Val Val Thr Gln His Phe Leu Gly Val Thr Tyr
45                      50                      55                      60

Glu Thr Gln Pro Ala
65

```

(2) INFORMATION FOR SEQ ID NO: 310:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -65..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq LLFLISLAHLSQ/WT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 310:

Met Asn Gln Ala Asp Pro Arg Leu Arg Ala Val Cys Leu Trp Thr Leu  
-65 -60 -55 -50  
Thr Ser Ala Ala Met Ser Arg Gly Asp Asn Cys Thr Asp Leu Leu Ala  
-45 -40 -35  
Leu Gly Ile Pro Ser Ile Thr Gln Ala Trp Gly Leu Trp Val Leu Leu  
-30 -25 -20  
Gly Ala Val Thr Leu Leu Phe Leu Ile Ser Leu Ala Ala His Leu Ser  
-15 -10 -5  
Gln Trp Thr Arg Gly Arg Ser Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO: 311:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -42..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq LLSILSSLTMTVIC/RH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 311:

Met His Arg Gln Ile Ser Phe Leu Leu Arg Lys Pro Arg Lys Asn  
           -40                          -35                          -30

Trp Phe Cys Gln Asn His Val Asn Leu Arg Lys Arg Tyr Leu Leu Ser  
      -25                          -20                          -15

Ile Leu Ser Ser Leu Thr Met Val Ile Cys Arg His Gly  
      -10                          -5                          1

## (2) INFORMATION FOR SEQ ID NO: 312:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq ALSAXTFVSLHA/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 312:

Met Lys Gln Trp Leu Cys Trp Val Leu Arg Leu Glu Gly Arg Gln Gly  
           -40                          -35                          -30

Leu Gly Val Gly Glu Pro Arg Gly Leu Arg Leu Cys Leu Gly Ala Leu  
      -25                          -20                          -15

Ser Ala Xaa Thr Phe Val Ser Phe Leu His Ala Ala Pro His Ser His  
      -10                          -5                          1                          5

Pro Ala Leu Gly

## (2) INFORMATION FOR SEQ ID NO: 313:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -66..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.8  
seq LLFFLFPILFIRS/QH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 313:

```

Met Arg Leu Gly Leu Cys Phe Trp Val Pro His Arg Gly Glu Met Ser
-65                               -60                               -55

Phe Ser Ser His Tyr Ser Arg Gly Thr Trp Tyr Gln Trp Asp Leu Ser
-50                               -45                               -40                               -35

Leu Leu Met Leu Thr Leu Ile Ser Trp Phe Arg Trp Cys Leu Pro Ala
                               -30                               -25                               -20

Val Ser Thr Val Glu Leu Leu Phe Phe Leu Phe Pro Ile Leu Phe Ile
-15                               -10                               -5

Arg Ser Gln His Arg
1

```

## (2) INFORMATION FOR SEQ ID NO: 314:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 108 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -101..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.6  
seq IIIIVITIT3ACSA/CI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 314:

```

Met Asp Phe Trp Glu Glu Tyr Arg Arg Gly Asp Val Pro Phe Ser Trp
-100                               -95                               -90

Cys Pro Ile Arg Ser Tyr Leu Met Ser Val Cys Pro Val Thr Gly Lys
-85                               -80                               -75                               -70

Val Asn Leu Asn His Leu Val Lys Val Ala Ser Ala Arg Phe Leu His
                               -65                               -60                               -55

Gln Val Thr Ile Phe Pro Phe Leu Tyr Ser Val Lys Ala Asn Tyr Cys
-50                               -45                               -40

```

- (A) LENGTH: 92 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5  
seq LALGSAGLLWCLA/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 316:

Met Val Phe Ala Thr Ile Gly Phe Ser Leu Lys Ser Gly Leu Ala Leu  
-25 -20 -15

Gly Ser Ala Gly Leu Leu Trp Cys Leu Ala Gly Phe Phe Gly Tyr Asp  
-10 -5 1 5

Thr Gln Gln Pro Thr Ala Pro Asn Ala Ile Glu Gly Tyr Arg Val Met  
10 15 20

Ser Ser Phe Gly Val Gly Ala Leu Phe Ala Ala Cys Thr Ile Cys Leu  
25 30 35

Leu Ala Xaa Lys Leu Asn Lys Gln Thr Thr Leu Lys Met Ala Asp Asp  
40 45 50

Leu Ala Gln Arg Arg Gln Gln Ala Asp Leu Ala Pro  
55 60 65

(2) INFORMATION FOR SEQ ID NO: 317:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq VLLLSGSVSVGVGC/CA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 317:

Met Val Leu Leu Leu Ser Gly Ser Val Ser Val Gly Val Cys Cys Ala  
                   -10                  -5                  1  
 Tyr Leu Cys Ile Ser Ile Ser Lys Thr Pro Thr Ala Cys Ala Leu Tyr  
                   5                  10                  15  
 Gly Leu Tyr Leu Pro Phe Phe  
           20                  25

## (2) INFORMATION FOR SEQ ID NO: 318:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq GLCXLGXVXNVFA/GS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 318:

Met Cys Ser Gln Lys Arg Ala Val Ser Asn Gln Gly Leu Met Asp Leu  
                   -25                  -20                  -15  
 Gly Leu Cys Xaa Leu Cys Xaa Val Xaa Asn Val Phe Ala Gly Ser Met  
                   -10                  -5                  1  
 Pro Gly Lys Ser His Cys His Ser Pro Phe Ser Ile Asn Gln Gly Arg  
           5                  10                  15

## (2) INFORMATION FOR SEQ ID NO: 319:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq LIVLTLHSPSCDT/AQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 319:

```

Met Leu Ile Val Leu Thr Leu His Ser Pro Ser Cys Asp Thr Ala Gln
      -10                      -5                      1

Glu Glu Met Gly Arg Val Pro Thr Thr Pro Lys Cys Arg Trp Lys Leu
      5                      10                      15

Gly Leu Ser Met Cys Ser Leu Leu Thr Pro Gly
      20                      25

```

(2) INFORMATION FOR SEQ ID NO: 320:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 124 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: -62..-1
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 6.4  
seq SVLWLGALGLTIQ/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 320:

```

Met Thr Arg Leu Cys Leu Pro Arg Pro Glu Ala Arg Glu Asp Pro Ile
      -60                      -55                      -50

Pro Val Pro Pro Arg Gly Leu Gly Ala Gly Glu Gly Ser Gly Ser Pro
      -45                      -40                      -35

Val Arg Pro Pro Val Ser Thr Trp Gly Pro Ser Trp Ala Gln Leu Leu
      -30                      -25                      -20                      -15

Asp Ser Val Leu Trp Leu Gly Ala Leu Gly Leu Thr Ile Gln Ala Val
      -10                      -5                      1

Phe Ser Thr Thr Gly Pro Ala Leu Leu Leu Leu Leu Val Ser Phe Leu
      5                      10                      15

Thr Phe Asp Leu Leu His Arg Pro Ala Val Thr Leu Cys His Ser Ala
      20                      25                      30

```



## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -15..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.4  
seq LTCFLFLSLISTYP/SC

```

Met Val Leu Thr Cys Leu Phe Leu Ser Leu Ile Ser Thr Tyr Pro Ser
-15          -10          -5          1

Cys Ile Thr Leu Phe Leu Ser Lys Ile Pro Asn Pro Leu Ser Ser Leu
          5          10          15

Pro Ser Leu
          20

```

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -18..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.3  
seq FSSFLQLLLSSST/NP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 322:

Met Leu Ile Pro Val Phe Ser Phe Ser Leu Gln Leu Leu Ser Ser Ser  
                   -15                  -10                  -5  
 Ser Thr Asn Pro Val Asn Ser Thr Phe Gln Met Pro Phe Glu Ser Ser  
                   1                  5                  10  
 His Xaa Thr Thr Arg  
                   15

## (2) INFORMATION FOR SEQ ID NO: 323:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -47..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.3  
seq LLLLESVSGLLP/RT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 323:

Met Ala Ala Ala Xaa Leu Ser Gly Pro Ser Ala Gly Ser Ala Ala Gly  
                   -45                  -40                  -35  
 Val Pro Gly Gly Thr Gly Gly Leu Ser Ala Val Ser Ser Gly Pro Arg  
                   -30                  -25                  -20  
 Leu Arg Leu Leu Leu Leu Glu Ser Val Ser Gly Leu Leu Gln Pro Arg  
                   -15                  -10                  -5                  1  
 Thr Gly Ser Ala Val Ala Pro Val His Pro Pro Asn Arg Ser Ala Arg  
                   5                  10                  15

## (2) INFORMATION FOR SEQ ID NO: 324:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi). ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2  
seq NWLFLEVFETFCNC/FF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 324:

```

Met His Asn Trp Leu Phe Leu Phe Val Phe Thr Phe Cys Asn Cys Phe
-15                -10                -5                1

Phe Lys Asn Asn Gly
                    5

```

## (2) INFORMATION FOR SEQ ID NO: 325:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -17..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.2  
seq CFYFLSTALGSQA/DS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 325:

```

Met His Val Glu Cys Phe Tyr Phe Leu Ser Thr Ala Leu Gly Ser Gln
-15                -10                -5

Ala Asp Ser Trp Val Ser Gly Leu Gln Gln Ala Gly Leu Leu Pro Ala
 1                5                10                15

Ile Gly Tyr Arg

```

## (2) INFORMATION FOR SEQ ID NO: 326:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 59 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -19..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.1  
seq LALLWSLPASDLG/RS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 326:

```
Met Ser Pro Gly Ser Ala Leu Ala Leu Leu Trp Ser Leu Pro Ala Ser
      -15                -10                -5
Asp Leu Gly Arg Ser Val Ile Ala Gly Leu Trp Pro His Thr Gly Val
      1                5                10
Leu Ile His Leu Glu Thr Ser Gln Ser Phe Leu Gln Gly Gln Leu Thr
      15                20                25
Lys Ser Ile Phe Pro Leu Cys Cys Thr Ser Leu
      30                35                40
```

(2) INFORMATION FOR SEQ ID NO: 327:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -13..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.1  
seq MALALGSIPSSIA/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 327:

```
Met Ala Leu Ala Leu Gly Ser Ile Pro Ser Ser Ile Ala Ser Ser Trp
      -10                -5                1
Val His Val Ser His Phe Cys Pro Cys Leu Leu His Thr Thr Leu Pro
      5                10                15
Gln Ser Thr Pro Lys
```

20

## (2) INFORMATION FOR SEQ ID NO: 328:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.1  
seq FLFCTLFSLVVHP/SH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 328:

```

Met Leu Ala Phe Leu Phe Cys Thr Leu Phe Ser Leu Val Val His Pro
-15                      -10                      -5

Ser His Ile Asp Leu Lys Cys Ser Phe Tyr
 1              5              10

```

## (2) INFORMATION FOR SEQ ID NO: 329:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 106 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -36..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6  
seq LLYTLQTISSLSG/CF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 329:

```

Met Ala Gln Met Pro Leu Thr Gly Ser Tyr Gln Asp Leu Glu Tyr Phe
-35                      -30                      -25

```

```

Leu Glu Cys Met Phe Leu His Leu Leu Tyr Thr Leu Gln Thr Ile Ser
-20          -15          -10          -5

Ser Leu Ser Gly Cys Phe Lys Gln Phe Phe Phe Gln Leu Asn Cys Phe
          1          5          10

Cys Trp Gly Glu Ile Leu Trp His Ser Ser Phe Leu His Ser Gly Ser
          15          20          25

Cys Leu Leu Val Leu Leu Ile Lys Lys Lys Lys Ile Tyr Leu Gln Ser
          30          35          40

Xaa Xaa Ile Tyr Thr Gly Tyr Xaa Ile Asp Xaa Xaa Xaa Leu Xaa Xaa
45          50          55          60

Phe Ser Ile Pro Leu Ser Phe Ile Gln Phe
          65          70

```

## (2) INFORMATION FOR SEQ ID NO: 330:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6  
seq LLMGLWVRTVLQG/KE

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 330:

```

Met Ala Leu Leu Met Gly Leu Trp Val Arg Thr Val Leu Gln Gly Lys
-15          -10          -5          1

Glu Ala Ser Gly
          5

```

## (2) INFORMATION FOR SEQ ID NO: 331:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -19..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6  
seq LAILIXSLKLTIG/IQ

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 331:

Met Ile Asn His Leu Tyr Leu Ala Ile Leu Ile Xaa Ser Leu Lys Leu  
-15 -10 -5

Thr Ile Gly Ile Gln Lys Arg Phe Gly Pro Pro  
1 5

## (2) INFORMATION FOR SEQ ID NO: 332:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -50..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6  
seq LLYLCSFPLPGTS/LF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 332:

Met Gly Arg Gln Gly Thr Leu Glu Ile Glu Gly Ile Leu Cys Val Ile  
-50 -45 -40 -35

Thr Trp Leu Glu Ala Asn Leu Gly Lys Gln Lys Asp Glu Asn His Tyr  
-30 -25 -20

Tyr Lys Lys Leu Ser Leu Leu Tyr Leu Cys Ser Phe Pro Leu Pro Gly  
-15 -10 -5

Thr Ser Leu Phe Leu Leu Cys Ser Phe Ser Tyr Leu Thr Gln Arg Leu  
1 5 10

Ser Gln Gly Gly Gly  
15

## (2) INFORMATION FOR SEQ ID NO: 333:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -39..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq SAWWCVLLEWSQG/AS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 333:

Met Glu Leu Thr Asn Lys Gln Thr Gly Thr Asp Arg His Glu Gln Val  
                  -35                  -30                  -25

Leu Arg Arg Val Lys Gln Asp Lys Arg Ile Ser Ala Trp Trp Cys Val  
                  -20                  -15                  -10

Leu Leu Glu Trp Ser Gln Gly Ala Ser Leu Arg Arg Gln His Arg Gly  
                  -5                          1                          5

Glu Thr Ser Pro Lys Ser Gly Glu Arg Leu Ser Arg Gln Arg Glu Gln  
  10                          15                          20                          25

Gln Lys Pro Gln Met Ser Asp Lys Ser Leu  
                          30                          35

## (2) INFORMATION FOR SEQ ID NO: 334:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 70 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq VLGLLFSTISDTWA/PA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 334:



Met Ala Lys Arg Gln Asn Pro Thr Ser Val Leu Gly Leu Leu Phe Ser  
 -20 -15 -10

Ile Ser Asp Thr Trp Ala Pro Ala Val Ser Ser Trp Lys Ala Glu Ala  
 -5 1 5 10

Lys Asp Gly Ala Asp Gln Glu Asp Ala Arg Xaa Xaa Ser Gln Arg Ser  
 15 20 25

Pro Xaa Ser Thr Ala Gly Ser Gln Glu Pro Tyr Phe Trp Phe Val Trp  
 30 35 40

Val Glu Gly Glu Gly Arg  
 45

## (2) INFORMATION FOR SEQ ID NO: 335:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -25...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.9  
seq FCLSLQIFRVSLA/LA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 335:

Met Asn Val Leu Pro Phe Ser Tyr Tyr Tyr Ile Leu Phe Cys Leu Ser  
 -25 -20 -15 -10

Leu Gln Ile Phe Arg Val Ser Leu Ala Leu Ala Xaa Thr His Glu Val  
 -5 1 5

Pro Val Ser Thr His Thr Asn Xaa Leu His  
 10 15

## (2) INFORMATION FOR SEQ ID NO: 336:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -29..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.9  
seq FSYISXFLSPVCG/CS

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 336:

```

Met Lys Cys Leu Lys Val Asn Pro Phe Leu Phe Leu Val Phe Asn Phe
      -25                      -20                      -15

Phe Ser Tyr Ile Ser Xaa Phe Leu Ser Pro Val Cys Gly Cys Ser Val
      -10                      -5                      1

Cys Asn Leu Lys His Trp Glu Asn Glu Leu Leu Phe Pro Ser Pro His
      5                      10                      15

Phe Leu Pro Tyr Lys Phe Xaa Phe Leu Phe
      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 337:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -33..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.8  
seq XLCLGMALCPRQA/TR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 337:

```

Met Ser Trp Thr Val Pro Val Val Arg Ala Ser Gln Arg Val Ser Ser
      -30                      -25                      -20

Val Gly Ala Asn Xaa Leu Cys Leu Gly Met Ala Leu Cys Pro Arg Gln
      -15                      -10                      -5

Ala Thr Arg Ile Pro Leu Asn Gly Thr Trp Leu Phe Thr Pro Val Ser
      1                      5                      10                      15

```

Lys Met Ala

## (2) INFORMATION FOR SEQ ID NO: 338:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.8  
seq FLXLMTLTTHVHS/SA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 338:

Met Gly Phe Leu Xaa Leu Met Thr Leu Thr Thr His Val His Ser Ser  
-15                      -10                      -5                      1

Ala Lys Pro Asn Gly  
5

## (2) INFORMATION FOR SEQ ID NO: 339:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7  
seq RVLLLAQLFLGSG/KT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 339:

Met Leu Phe Arg Val Leu Leu Leu Ala Gln Leu Phe Leu Gly Ser Gly  
-15                      -10                      -5

Lys Thr Leu Arg Thr Pro  
 1 5

## (2) INFORMATION FOR SEQ ID NO: 340:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -32..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7  
seq SLPLSTSA?PLRG/LR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 340:

Met Arg Val Pro Glu Asp Leu Ala Ser Lys Ile Leu Leu Pro Gly Cys  
 -30 -25 -20

Ala Pro Gly Ser Leu Pro Leu Ser Thr Ser Ala Pro Pro Leu Arg Gly  
 -15 -10 -5

Leu Arg Leu Lys Glu His Pro Gly Arg Gly Pro Ser Ser Pro Lys Ala  
 1 5 10 15

Ala Cys Pro Glu Thr Pro Ala  
 20

## (2) INFORMATION FOR SEQ ID NO: 341:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -31..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.7  
seq SDLCQCQILARA/HD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 341:

Met Phe Pro His Xaa Glu Thr Gln Val Lys Cys Phe Trp Gln Gly Leu  
 -30 -25 -20  
 Arg Arg Ser Asp Leu Cys Leu Cys Gln Cys Ile Leu Ala Arg Ala His  
 -15 -10 -5 1  
 Asp Gly Asp Leu Tyr Leu Phe Phe  
 5

## (2) INFORMATION FOR SEQ ID NO: 342:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 64 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.6  
seq LAVFMXLAQLVSG/NW

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 342:

Met Lys Ser Leu Leu Phe Thr Leu Ala Val Phe Met Xaa Leu Ala Gln  
 -20 -15 -10 -5  
 Leu Val Ser Gly Asn Trp Tyr Val Lys Lys Cys Leu Asn Xaa Phe Gly  
 1 5 10  
 Ile Cys Lys Xaa Lys Cys Lys Pro Glu Glu Met His Val Lys Asn Gly  
 15 20 25  
 Trp Xaa Met Cys Gly Lys Gln Arg Asp Cys Cys Val Pro Ala Asn Gly  
 30 35 40

## (2) INFORMATION FOR SEQ ID NO: 343:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -23..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.6  
seq LLNVACCIPFSSS/LF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 343:

```

Met His Leu Tyr Ser Cys Ser Cys Met Arg Leu Leu Asn Val Ala Cys
      -20                      -15                      -10

Cys Ile Pro Phe Ser Ser Ser Leu Phe Pro His Ile Leu Phe Lys Ser
      -5                      1                      5

Leu Asn Tyr Ser Leu Thr Ser Phe Leu Lys Ala Val Arg Gly Arg Trp
10                      15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 344:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -16..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.5  
seq PLVLSPLSYQCSS/QG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 344:

```

Met Arg Ala Pro Leu Val Leu Ser Pro Leu Ser Tyr Gln Cys Ser Ser
-15                      -10                      -5

Gln Gly His Ile Trp
1                      5

```

## (2) INFORMATION FOR SEQ ID NO: 345:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 amino acids  
(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -36..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.5  
seq FTSMCILFHCLLS/FQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 345:

Met Gln Val Pro His Leu Arg Val Trp Thr Gln Val Xaa Asp Thr Phe  
-35 -30 -25

Ile Gly Tyr Arg Asn Leu Gly Phe Thr Ser Met Cys Ile Leu Phe His  
-20 -15 -10 -5

Cys Leu Leu Ser Phe Gln Arg  
1

(2) INFORMATION FOR SEQ ID NO: 346:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 46 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig peptide

(B) LOCATION: -36..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.4  
seq LWLMHQSEFQKSNS/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 346:

Met Gln Lys Leu Met Ala Val Pro Met Ile Thr Arg Ala Gln Gly Gly  
-35 -30 -25

Asp Thr Cys Thr Arg Gln Ile Leu Trp Leu Met His Gln Ser Phe Gln  
-20 -15 -10 -5

Lys Ser Asn Ser Ser Ser Thr Ser Tyr Cys Ser Ala Gln Gly  
1 5 10

## (2) INFORMATION FOR SEQ ID NO: 347:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -45..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4  
seq AHRSLCLWPACLC/AR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 347:

Met Cys Xaa Ala Gly Phe Xaa Asp His Pro Arg Ala Ala Arg His Ala  
-45 -40 -35 -30

Arg Thr Ser Arg His Pro Leu Pro Trp Val Cys Val Ser Gln Xaa Pro  
-25 -20 -15

Ala His Arg Ser Leu Cys Leu Trp Pro Ala Cys Leu Cys Ala Arg Val  
-10 -5 1

Leu Pro Pro Ala Pro Gly  
5

## (2) INFORMATION FOR SEQ ID NO: 348:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4  
seq ILVSFILAALSLS/TT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 348:



Met Thr Ser Lys Phe Ile Leu Val Ser Phe Ile Leu Ala Ala Leu Ser  
-15 -10 -5

Leu Ser Thr Thr Ile Gly  
1

(2) INFORMATION FOR SEQ ID NO: 349:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.4  
seq LLIFILTVHHTPS/LP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 349:

Met His Leu Leu Ile Phe Ile Leu Thr Val His His Thr Pro Ser Leu  
-15 -10 -5 1

Pro Ser

(2) INFORMATION FOR SEQ ID NO: 350:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq SSLMVQLISQVYS/CM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 350:

Met Leu Ser Ser Ser Leu Met Val Gln Leu Ile Ser Gln Val Tyr Ser  
 -15 -10 -5

Cys Met Arg Arg  
 1

(2) INFORMATION FOR SEQ ID NO: 351:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq FSYILCMLFCLFS/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 351:

Met Phe Ser Tyr Ile Leu Cys Met Leu Phe Cys Leu Phe Ser Gln Asp  
 -10 -5 1

Lys Phe Leu Glu Val Thr Leu Leu Cys Glu Arg Tyr Met Leu  
 5 10 15

(2) INFORMATION FOR SEQ ID NO: 352:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2  
seq VTILAFSLLVLSSES/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 352:

```

Met Leu Phe Leu Tyr Tyr Val Thr Leu Ala Phe Ser Leu Leu Val Leu
      -15                      -10                      -5
Ser Glu Ser Ala Val Leu Lys Arg Arg Glu Ile Phe Xaa Thr Gly Leu
      1                      5                      10
Gly Cys Val Thr Gly Leu Gly Cys Val Thr Gly Leu Arg
      15                      20                      25

```

## (2) INFORMATION FOR SEQ ID NO: 353:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.2  
seq LLSGLWLSSVKEC/DD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 353:

```

Met Leu Leu Ser Gly Leu Trp Leu Ser Ser Val Lys Glu Cys Asp Asp
      -10                      -5                      1
Trp Arg Ala Asp Gly Cys Leu Pro Ser Ile Val His Pro Leu Arg
      5                      10                      15

```

## (2) INFORMATION FOR SEQ ID NO: 354:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.2  
seq VFCTSWLMSSSSP/SI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 354:

```
Met Val Ala Phe Ser Val Phe Cys Phe Ser Trp Leu Met Ser Ser Ser
      -15              -10              -5

Ser Pro Ser Ile Phe Trp Ser His Phe Tyr Ser Pro Phe Lys Asp Leu
      1              5              10

Ser Lys Met Tyr Asn Tyr Val Ser Pro
      15              20
```

(2) INFORMATION FOR SEQ ID NO: 355:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 42 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -16..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.1  
seq LALGIGPPGCLQG/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 355:

```
Met Val Pro Leu Ala Leu Gly Ile Gly Pro Pro Gly Cys Leu Gln Gly
      -15              -10              -5

Ser Pro Ser Gln Trp Leu Val Arg Ala Pro Gly Ala Gln Leu Ser Pro
      1              5              10              15

Ile Gly Val Ala Thr Glu Arg Glu Gln Arg
      20              25
```

(2) INFORMATION FOR SEQ ID NO: 356:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -32..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1  
seq LLWFCTAMRPGGA/GL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 356:

Met Asn Leu Cys Met Gly Val Leu Leu Lys Val Gly Thr Ser Arg Arg  
-30 -25 -20  
Cys Leu Cys Leu Leu Trp Phe Cys Thr Ala Met Arg Pro Gly Gly Ala  
-15 -10 -5  
Gly Leu Pro Asn Ala Thr Pro Glu Trp  
1 5

## (2) INFORMATION FOR SEQ ID NO: 357:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.1  
seq SLAKSLFLRVARG/LG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 357:

Met Ser Leu Ala Lys Ser Leu Phe Leu Arg Val Ala Arg Gly Leu Gly  
-10 -5 1

## (2) INFORMATION FOR SEQ ID NO: 358:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 94 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -18..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5.1  
seq FLPSATLLLSAES/FF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 358:

Met Arg Leu Pro Pro Phe Leu Pro Ser Ala Thr Leu Leu Leu Ser Ala  
-15 -10 -5  
Glu Ser Phe Phe Arg Ser Val Ser Glu Tyr Pro Ser Leu Pro Ser Pro  
1 5 10  
Ser Ala Gly Gly Pro Gly Cys Val Ser Gly Lys Trp Gly Ser Gly Ser  
15 20 25 30  
Asn Gly Pro Leu Ser Ser Leu Ser Cys Ser Leu Cys Arg Lys Pro Leu  
35 40 45  
Leu His Ser Thr Ala Leu Ser Ser Ser Arg Pro Phe Phe Ser Pro Gly  
50 55 60  
Phe Pro Cys Gln Ile Ser Pro Arg Ser Gly Leu His Pro Leu  
65 70 75

## (2) INFORMATION FOR SEQ ID NO: 359:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -49..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 5  
seq PLLLLLREELVTG/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 359:

Met Ser Asp Arg Lys Arg Thr Lys Phe Ser Tyr Val Gln Leu Pro Cys  
-45 -40 -35  
Pro Ile Ser Leu Leu Pro Arg Ser Phe Lys Arg Gly Gln Ile Pro Gly

-30                      -25                      -20  
 Pro Ser Ala Pro Pro Leu Leu Leu Leu Arg Glu Glu Leu Val Thr  
       -15                      -10                      -5  
 Gly Ala Val  
       1

## (2) INFORMATION FOR SEQ ID NO: 360:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -41...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq FCFFPAFLVXVXS/QP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 360:

Met Thr Pro Leu Gly Ser Gly Pro Pro Arg Glu Ala Ser Ile Ala Gln  
       -40                      -35                      -30  
 Val Arg Gly Phe Ser Arg Thr Phe Phe Arg Val Ala Phe Cys Phe Phe  
       -25                      -20                      -15                      -10  
 Pro Ala Phe Leu Val Xaa Val Xaa Ser Gln Pro Ser Gly Phe Ser Thr  
                      -5                      1                      5  
 Thr Glu Thr Leu Cys Ala Gln Asp Phe Ser Asp Val Ile Phe Leu Arg  
       10                      15                      20  
 Arg Ala Asp Thr Arg Arg Trp Lys Lys Lys Gln Leu Arg Arg Arg  
       25                      30                      35

## (2) INFORMATION FOR SEQ ID NO: 361:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq CSALFPLLSLLSC/KE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 361:

```

Met Arg Cys Ser Ala Leu Phe Pro Leu Leu Ser Leu Leu Ser Cys Lys
-15          -10          -5          :
Glu Arg Xaa Trp Cys Leu Ser Thr Leu Glu Asp Ala Ala Thr Xaa Arg
          5          10          15
His Leu Gly Ser Arg Glu Gln Pro Ser Gly Asp Ala Glu Pro Val Glu
          20          25          30
Val Trp
          35

```

(2) INFORMATION FOR SEQ ID NO: 362:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5  
seq IISLLKLCSFCFI/KD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 362:

```

Met Leu Tyr Asp Gln Tyr Tyr Leu Ile Ile Ser Leu Leu Lys Leu Cys
-20          -15          -10
Ser Phe Cys Phe Ile Lys Asp Phe Lys Ala Ser Asn Ile Thr Leu Val
-5          1          5          10
Val Ile Leu

```

(2) INFORMATION FOR SEQ ID NO: 363:



- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 75 amino acids  
    (B) TYPE: AMINO ACID  
    (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:  
    (A) ORGANISM: Homo Sapiens  
    (F) TISSUE TYPE: Uterus
- (ix) FEATURE:  
    (A) NAME/KEY: sig\_peptide  
    (B) LOCATION: -65..-1  
    (C) IDENTIFICATION METHOD: Von Heijne matrix  
    (D) OTHER INFORMATION: score 5  
                                seq LCSFLSLRFCTLS/FM
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 363:

Met	Ala	Asn	Cys	Phe	Leu	Ser	His	Lys	Ser	Gln	Thr	Ile	Leu	Ile	Ser	
-65					-60					-55					-50	
Lys	Pro	Ala	Leu	Thr	Gln	Ser	His	Phe	Thr	Ser	Pro	Ala	Gly	Leu	Phe	
				-45					-40					-35		
Leu	Thr	Val	Glu	Lys	Ser	His	Leu	Leu	Thr	Arg	Leu	Phe	Phe	His	Trp	
			-30					-25					-20			
Leu	Ser	Leu	Val	Leu	Cys	Ser	Phe	Leu	Ser	Leu	Arg	Phe	Cys	Thr	Leu	
	-15						-10					-5				
Ser	Phe	Met	Cys	Ser	Phe	Ala	Leu	Phe	His	Leu						
	1				5					10						

(2) INFORMATION FOR SEQ ID NO: 364:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 93 amino acids  
    (B) TYPE: AMINO ACID  
    (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:  
    (A) ORGANISM: Homo Sapiens  
    (F) TISSUE TYPE: Testis
- (ix) FEATURE:  
    (A) NAME/KEY: sig\_peptide  
    (B) LOCATION: -18..-1  
    (C) IDENTIFICATION METHOD: Von Heijne matrix  
    (D) OTHER INFORMATION: score 5  
                                seq LTYLLEFLPDWAAV/FE
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 364:

```

Met His Gly Ala Gly Leu Thr Tyr Leu Leu Phe Leu Pro Asp Trp Ala
      -15              -10              -5

Ala Val Phe Glu Leu Tyr Asn Cys Glu Asp Glu Arg Cys Tyr Leu Asp
      1              5              10

Leu Ala Arg Leu Arg Gly Val His Tyr Ile Thr Trp Arg Arg Gln Asn
      15              20              25              30

Lys Val Phe Pro Gln Asp Lys Gly His His Pro Thr Leu Gly Glu His
      35              40              45

Pro Lys Phe Thr Asn Tyr Ser Phe Asp Val Glu Glu Phe Met Tyr Leu
      50              55              60

Val Leu Gln Ala Ala Asp His Val Leu Gln His Pro Gly
      65              70              75

```

## (2) INFORMATION FOR SEQ ID NO: 365:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.9  
seq CLSATLAFSGSFL/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 365:

```

Met Cys Cys Leu Ser Ala Thr Leu Ala Phe Ser Gly Ser Phe Leu Ala
-15              -10              -5              1

Pro His Leu Ile Phe Cys Cys Phe Ser His Leu Asn Val Ile Ile Leu
      5              10              15

Leu Ser Ser Leu Ser Pro Ile His Gly
      20              25

```

## (2) INFORMATION FOR SEQ ID NO: 366:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -39..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.9  
seq SGLRGLLLQEALG/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 366:

```

Met Ala Glu Leu Asp Leu Met Ala Pro Gly Pro Leu Pro Arg Ala Thr
      -35                      -30                      -25

Ala Gln Pro Pro Ala Pro Leu Ser Pro Asp Ser Gly Leu Arg Gly Leu
      -20                      -15                      -10

Leu Leu Gln Glu Ala Leu Gly Ala Val Pro Asp Pro Arg
      -5                      1                      5

```

(2) INFORMATION FOR SEQ ID NO: 367:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 56 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -28..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.9  
seq FLVACPLFGVCLX/FF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 367:

```

Met Thr Leu Thr His Gly Asn Asn Ile Leu His Leu Ala Asn Phe Phe
      -25                      -20                      -15

Leu Val Ala Cys Pro Leu Phe Gly Val Cys Leu Xaa Phe Phe Ile Leu
      -10                      -5                      1

Arg Phe Arg Leu Tyr Ile Gln Gly Pro Asn Val Thr Gln Val Ile Leu
      5                      10                      15                      20

His Leu Ser Gln Gly Thr Leu Ser
      25

```

(2) INFORMATION FOR SEQ ID NO: 368:

- ```
(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 16 amino acids
    (B) TYPE: AMINO ACID
    (C) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:
    (A) ORGANISM: Homo Sapiens
    (F) TISSUE TYPE: Ovary

(ix) FEATURE:
    (A) NAME/KEY: sig_peptide
    (B) LOCATION: -14..-1
    (C) IDENTIFICATION METHOD: Von Heijne matrix
    (D) OTHER INFORMATION:   score 4.9
                               seq VLRWLPWPRGSHS/DS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 368:
```

Met Val Leu Arg Trp Leu Pro Trp Pro Arg Gly Ser His Ser Asp Ser  
-10 -5 1

(2) INFORMATION FOR SEQ ID NO: 369:

- ```

(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 39 amino acids
    (B) TYPE: AMINO ACID
    (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:
    (A) ORGANISM: Homo Sapiens
    (F) TISSUE TYPE: Uterus

(ix) FEATURE:
    (A) NAME/KEY: sig_peptide
    (B) LOCATION: -24..-1
    (C) IDENTIFICATION METHOD: Von Heijne matrix
    (D) OTHER INFORMATION: score 4.8
                           seq FSFLGTLFHKSNS/ED

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 369:

```

Met Lys Ala Arg Leu Ser Gly Asn Leu Ile Cys Phe Ser Phe Leu Gly  
-20 -15 -10

Thr Leu Phe His Lys Ser Asn Ser Glu Asp Ser Ser Val Gly Lys Gly  
-5 1 5

Asn Trp Lys Lys Lys Asn Lys

10

15

## (2) INFORMATION FOR SEQ ID NO: 370:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq VCLVPQTPSLCLG/KG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 370:

```

Met Ser His Val Cys Leu Val Pro Gln Thr Pro Ser Leu Cys Leu Gly
-15                      -10                      -5

Lys Gly Thr Pro Arg Ser Arg Ser Ala Pro Phe Gln Ser Ser Gly Pro
 1              5              10              15

His Arg Leu Cys Ala
                20

```

## (2) INFORMATION FOR SEQ ID NO: 371:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq VLTSVNLFING/SV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 371:

300

Met Tyr Pro Ala Ser Phe Val Phe Lys Ile Pro Ser Thr Ala Tyr Val  
                   -25                  -20                  -15

Val Leu Thr Ser Val Asn Leu Phe Ile Gly Ile Asn Gly Ser Val Ala  
                   -10                  -5                  1

Thr Phe Val Leu Glu Leu Phe Thr Asp Asn Lys Leu Asn Asn Ile Asn  
           5                  10                  15

Asp Ile Leu Lys Ser Val Phe Leu Ile Phe Pro His Phe Cys Leu Gly  
   20                  25                  30                  35

Arg Gly Gln Thr

## (2) INFORMATION FOR SEQ ID NO: 372:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq RSSLWVTAPLVSA/CP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 372:

Met Ser Ser Ser Arg Lys Asp His Leu Gly Ala Xaa Ala Gln Ser Pro  
   -30                  -25                  -20                  -15

Ser Arg Ser Ser Leu Trp Val Thr Ala Pro Leu Val Ser Ala Cys Pro  
                   -10                  -5                  1

Thr Cys Ser Pro Ala Thr His Pro Thr Gly  
           5                  10

## (2) INFORMATION FOR SEQ ID NO: 373:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 56 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -19..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.8  
 seq ATYLVQSSACCPA/IV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 373:

Met Ala Ser Pro Ala Ala Ala Thr Tyr Leu Val Gln Ser Ser Ala Cys  
                   -15                  -10                  -5  
 Cys Pro Ala Ile Val Arg His Leu Cys Gln Xaa Tyr Arg Ser Met Pro  
                   1                          5                          10  
 Val Phe Leu Asp Pro Ala Xaa Ile Ala Thr Leu Glu Gly Ile Ser Trp  
           15                          20                          25  
 Arg Leu Pro Ser Ala Pro Ser Asp  
       30                          35

(2) INFORMATION FOR SEQ ID NO: 374:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 69 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -61..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.7  
 seq LLPCNLHXSWLHS/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 374:

Met Asn Ala Ala Ile Asn Thr Gly Pro Ala Pro Ala Val Thr Lys Thr  
       -60                          -55                          -50  
 Glu Thr Glu Val Gln Asn Pro Asp Val Leu Trp Asp Leu Asp Ile Pro  
       -45                          -40                          -35                          -30  
 Glu Ala Arg Ser His Ala Asp Gln Asp Ser Asn Pro Xaa Ala Glu Ala  
           -25                          -20                          -15  
 Leu Leu Pro Cys Asn Leu His Xaa Ser Trp Leu His Ser Ser Pro Arg  
       -10                          -5                          1

Pro Asp Pro His Ser  
5

(2) INFORMATION FOR SEQ ID NO: 375:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 61 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq GIFLVIFCSESFS/LL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 375:

```

Met Ile Asn Leu Leu Val Gly Asn Cys Ile Tyr Leu Leu Gly Ala Ile
      -40                -35                -30

Arg Ala Ser Cys Met Cys Arg Xaa Met Ser Phe Ala Lys Phe Gly Ile
      -25                -20                -15

Phe Leu Val Ile Phe Cys Ser Glu Ser Phe Ser Leu Leu Leu Trp Asn
      -10                -5                1                5

Phe Ser Ser Ile Tyr Val Lys Thr Phe Trp Pro Val Gly
              10                15

```

(2) INFORMATION FOR SEQ ID NO: 376:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq LRFLLRDPGCLLA/QP



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 376:

Met Leu Cys Cys Gly Pro Leu Arg Phe Leu Leu Arg Asp Pro Gly Cys  
                           -15                          -10                          -5

Leu Leu Ala Gln Pro Glu Leu Ala Phe Trp Gly Pro Ala Ser Phe Ile  
                           1  5  10

Ser Gly Gly Leu Val Val Val Ser Glu Thr Pro His Pro Ser Phe Pro  
       15  20  25

Leu Asp Pro Pro  
       30

## (2) INFORMATION FOR SEQ ID NO: 377:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.7  
seq ILLRMTVLPTLWT/RR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 377:

Met Arg Lys Thr Ser Phe Ile Leu Leu Arg Met Thr Val Leu Pro Thr  
                           -15                          -10                          -5

Leu Trp Thr Arg Arg Arg Val Gln Leu Val  
                           1  5

## (2) INFORMATION FOR SEQ ID NO: 378:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq VRVGLVLVXRALC/LX

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 379:

```

Met Trp Trp Lys Pro Ala Pro Glu Glu Gly Val Arg Val Gly Leu Val
    -20                -15                -10

Leu Val Xaa Arg Ala Leu Cys Leu Xaa Val Leu Ser Arg Phe Met Phe
    -5                      1                      5

Xaa Asn Pro Gly Leu Gly Gly Met Gly
10                      15

```

## (2) INFORMATION FOR SEQ ID NO: 379:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq FNFLLGNSSCVYQ/RP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 379:

```

Met Phe Asn Phe Leu Leu Gly Asn Ser Ser Cys Val Tyr Gln Arg Pro
    -10                -5                1

Ile Arg Leu Lys Leu Ile Ile Phe Pro Ser Gly
    5                      10

```

## (2) INFORMATION FOR SEQ ID NO: 380:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 119 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -42..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq LDPAVSLSAPAFSA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 380:

```

Met Lys Arg Gly Ala Phe Ser Asn Leu Asn Asp Ser Gln Leu Ser Ala
  -40                      -35                      -30

Ser Phe Leu Gln Pro Ser Leu Gln Ala Asn Cys Pro Ala Leu Asp Pro
  -25                      -20                      -15

Ala Val Ser Leu Ser Ala Pro Ala Phe Ala Ser Ala Leu Arg Ser Met
  -10                      -5                      1                      5

Lys Ser Ser Gln Ala Ala Arg Lys Asp Asp Phe Leu Arg Ser Leu Ser
      10                      15                      20

Asp Gly Asp Ser Gly Thr Ser Glu His Ile Ser Ala Val Val Thr Ser
      25                      30                      35

Pro Arg Ile Ser Cys His Gly Ala Ala Ile Pro Xaa Ala Xaa Ala Xaa
      40                      45                      50

Xaa Xaa Gly Cys Ser Cys Xaa Thr Glu Arg Xaa Leu Xaa Xaa Pro Pro
      55                      60                      65                      70

Ser Leu Leu Ser Leu Glu Ala
      75

```

## (2) INFORMATION FOR SEQ ID NO: 381:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 80 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -24..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq FFIFCSLNTLLG/GV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 381:

```

Met Lys Ser Ala Lys Leu Gly Phe Leu Leu Arg Phe Phe Ile Phe Cys
      -20              -15              -10

Ser Leu Asn Thr Leu Leu Leu Gly Gly Val Asn Lys Ile Ala Glu Lys
      -5              1              5

Ile Cys Gly Asp Leu Lys Asp Pro Cys Lys Leu Asp Met Asn Phe Gly
      10              15              20

Ser Cys Tyr Glu Val His Phe Arg Tyr Phe Tyr Asn Arg Thr Ser Lys
      25              30              35              40

Arg Cys Glu Thr Phe Val Phe Ser Ser Cys Asn Gly Asn Leu Asn Gly
      45              50              55

```

## (2) INFORMATION FOR SEQ ID NO: 382:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq ILFPLHSVIGSHP/QC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 382:

```

Met Asp Ile Leu Phe Pro Leu His Ser Val Ile Gly Ser His Pro Gln
-15              -10              -5              1

Cys Leu Pro Glu Arg Xaa Thr Ala Arg Met Ile Lys Leu Lys Trp Gly
      5              10              15

Asn Gly Ser Gly Ser Asp Phe Gly
      20              25

```

## (2) INFORMATION FOR SEQ ID NO: 383:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -27..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq FGILILLSQRQWS/KN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 383:

```

Met Leu Lys Val Phe Arg Ala Xaa His Pro Lys Ile Cys His Phe Gly
  -25                      -20                      -15

Ile Leu Ile Leu Leu Ser Gln Arg Gln Trp Ser Lys Asn Arg Cys Arg
  -10                      -5                      1                      5

Glu Gly Cys Leu Thr Thr Leu Phe Leu Phe Glu Ala Glu His Lys Ser
          10                      15                      20

Ser Leu Val Lys
          25

```

(2) INFORMATION FOR SEQ ID NO: 384:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 45 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -34..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq LKWRKLAASWTLS/QE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 384:

```

Met Leu Val Arg Asn Ala Arg Arg Gly Ser Arg Gly Arg Ser Pro Trp
          -30                      -25                      -20

Trp Arg Ala Gly Cys Leu Xaa Trp Arg Lys Leu Ala Ala Ser Trp Thr
          -15                      -10                      -5

Leu Ser Gln Glu Ile Phe Arg Gly Ser Arg Lys Gly Ser
          1                      5                      10

```

## (2) INFORMATION FOR SEQ ID NO: 385:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 52 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq FTLGLGYPIPTRL/QP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 385:

Met Thr Lys Gly His His His Gln His Pro Leu His Pro His Pro Leu  
                  -25                  -20                  -15

Phe Thr Leu Gly Leu Gly Tyr Pro Ile Pro Thr Arg Leu Gln Pro Cys  
                  -10                  -5                  1

Thr Leu Ser Ser Asp Pro Leu Leu Asp Ile Thr Cys Ser Leu Arg Ser  
          5                  10                  15

Pro Ser Ser Gly  
          20

## (2) INFORMATION FOR SEQ ID NO: 386:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq RLHILFIVCLARG/KG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 386:

Met Thr Tyr His Xaa Ile Gln Phe Ser Glu Arg Leu His Ile Leu Phe  
 -20 -15 -10

Ile Val Cys Leu Ala Arg Gly Lys Gly  
 -5 1

(2) INFORMATION FOR SEQ ID NO: 387:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 74 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -46..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq LIYCGLSQPLTLG/VT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 387:

Met Ser Gln Phe Pro Leu Cys Ser Pro Pro Trp Lys Pro Leu Val Lys  
 -43 -40 -35

Val Ser Arg Asn Leu Lys Ile Arg Met Ser Ile Pro Trp Pro Leu Ser  
 -30 -25 -20 -15

Val Leu Ile Tyr Cys Gly Leu Ser Gln Pro Leu Thr Leu Gly Val Thr  
 -10 -5 1

Ser Pro Ser Phe Pro Gln Asn Ser Phe Phe Pro Trp Leu Pro Glu His  
 5 10 15

Pro Thr His Leu Val Ser Ser Thr Pro Gln  
 20 25

(2) INFORMATION FOR SEQ ID NO: 393:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 140 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 389:

```

Met Val Leu Thr Thr Leu Pro Leu Pro Ser Ala Asn Ser Pro Val Asn
  -35                -30                -25

Met Pro Thr Thr Gly Pro Asn Ser Leu Ser Tyr Ala Ser Ser Ala Leu
  -20                -15                -10

Ser Pro Cys Leu Xaa Ala Pro Lys Ser Pro Arg Leu Gly
  -5                  1                5

```

(2) INFORMATION FOR SEQ ID NO: 390:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq LLPTLPWLPSRL/LS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 390:

```

Met Gln Arg Asn Ala Thr Phe Ile His Leu Gln Leu Ala Ile Arg Pro
-30                -25                -20                -15

Ser Leu Leu Pro Thr Leu Pro Trp Leu Pro Ser Thr Arg Leu Leu Ser
          -10                -5                1

Pro Thr Pro Leu Gly Gln Leu Arg Gly Pro Pro Gly Xaa Gln Arg Ala
      5                10                15

Met Pro Thr Ala His Leu Arg
  20                25

```

(2) INFORMATION FOR SEQ ID NO: 391:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:



- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq ILFCFHSFHPPLFQ/DT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 391:

Met Asn Ile Leu Phe Cys Phe His Ser Phe His Pro Leu Phe Gln Asp  
 -15                      -10                      -5                      1

Thr Ile Glu Phe  
                     5

## (2) INFORMATION FOR SEQ ID NO: 392:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq FNFLFLVQLCILA/CD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 392:

Met Leu Thr Asn Arg Asn Tyr Phe Asn Phe Leu Phe Leu Val Gln Leu  
 -20                      -15                      -10                      -5

Cys Ile Leu Ala Cys Asp Asn Ala Tyr Leu Gln Ser Cys Pro Leu Thr  
                     1                      5                      10

Ser Lys Thr Pro Leu Leu Gln Thr His Ser Ala Leu Phe Tyr Asn Ser  
                     15                      20                      25

Thr Tyr Gly Ile Phe Leu Leu Gly Val  
                     30                      35

## (2) INFORMATION FOR SEQ ID NO: 393:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq ALCRFVGMQPCTA/QT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 393:

```

Met Lys Leu Asn Pro Gly Gln Val Pro Thr Trp Trp Glu Ala Leu Cys
-25                -20                -15

Arg Phe Val Gly Met Gln Pro Cys Thr Ala Gln Thr Gly Leu Leu Pro
-10                -5                1                5

His Gly Thr His Asn Thr Arg Glu Arg Gln Arg Asp Pro Ser Ala Gln
10                15                20

Lys Asn Thr Arg Arg Phe Ser Pro Val Gly
25                30

```

(2) INFORMATION FOR SEQ ID NO: 394:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -27...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq LCLNLCPCSSSLL/SP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 394:

```

Met Leu Ala Gly Phe Arg Arg Ser Ala Pro Ala Ser Gln Ser Leu Cys
-25                -20                -15

Leu Asn Leu Cys Pro Cys Ser Ser Ser Leu Leu Ser Pro Ala

```

-10

-5

1

## (2) INFORMATION FOR SEQ ID NO: 395:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq SFYLLFFLNDVPP/CP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 395:

Met Lys Glu Gly Ala Ser Phe Tyr Leu Leu Phe Phe Leu Asn Asp Val  
-15 -10 -5

Pro Pro Cys Pro Pro His Thr Pro Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO: 396:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq ETLLKLSSQSRT/NR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 396:

Met Gly Leu Glu Cys Cys Cys Pro Pro His Asn Leu Arg Val Tyr Ile  
-25 -20 -15

Glu Thr Leu Leu Leu Lys Leu Ser Ser Gln Ser Arg Thr Asn Arg Leu  
 -10 -5 1

## (2) INFORMATION FOR SEQ ID NO: 397:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq VLSIAASLLQCRL/AV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 397:

Met Gln Leu Cys Pro Phe Thr Ser Val Leu Ser Ile Ala Ala Ser Leu  
 -20 -15 -10

Leu Gln Cys Arg Leu Ala Val Val Thr Glu Thr Ile Trp Pro Pro Gln  
 -5 1 5 10

Xaa Trp

## (2) INFORMATION FOR SEQ ID NO: 398:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -44..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq QLLFKLNSTWCRA/LQ

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 398:

```

Met Asp Val Thr Cys Cys Phe Asp Ala Val Glu Gly Ser Asp Phe Arg
      -40              -35              -30

Val Cys Cys His Gly Cys Val Ser Trp Leu Cys Leu Gln Met Leu Gln
      -25              -20              -15

Leu Leu Phe Lys Leu Asn Ser Thr Trp Cys Arg Ala Leu Gln Ser Glu
      -10              -5              1

Thr Ser Leu Ala Ser Arg Arg Leu Trp Met Trp Val Ser His Leu Xaa
      5              10              15              20

Glu Phe Phe Thr Val Thr Pro Trp
              25

```

## (2) INFORMATION FOR SEQ ID NO: 399:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq HCFCFTLFSYSSS/FF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 399:

```

Met Arg Gln Gly Pro Gly Ala Pro Leu His Cys Phe Cys Phe Thr Leu
      -20              -15              -10

Phe Ser Tyr Ser Ser Ser Phe Phe Phe Phe
      -5              1

```

## (2) INFORMATION FOR SEQ ID NO: 400:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

- ```
(A) NAME/KEY: sig_peptide
(B) LOCATION: -15, -1
(C) IDENTIFICATION METHOD: Von Heijne matrix
(D) OTHER INFORMATION: score 4.2
                        seq ITLLGIWLTXLRLQ/FP
```

Met His Ile Thr Leu Leu Gly Ile Trp Leu Thr Xaa Arg Leu Gln Phe  
-15 -10 -5 1

Pro Arg Ser Gly Arg Ala Gly  
5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -17..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.2  
seg SWVCLLSAGTAFE/DY

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 401:

Met Leu Tyr Gly Ser Trp Val Cys Leu Leu Ser Ala Gly Thr Ala Phe  
-15 -10 -5

Glu Asp Tyr His Leu Gly Gly Thr  
1 5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(E) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -32..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq XXXXFLLGRRVVG/ES

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 402:

Met Leu Phe Phe Pro Leu Leu Ser Phe Arg Phe Leu Pro Ser Glu Ser  
-30 -25 -20

Leu Leu Lys Xaa Xaa Xaa Xaa Phe Leu Leu Gly Arg Arg Val Val Gly  
-15 -10 -5

Glu Ser Xaa Phe Ile Phe Thr Cys Gly Asn Leu Leu Leu Ile Trp Pro  
1 5 10 15

Tyr Gly

## (2) INFORMATION FOR SEQ ID NO: 403:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.2  
seq WAILGCWGTLRG/HL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 403:

Met Pro Val Trp Ala Ile Leu Gly Cys Trp Gly Thr Leu Ser Arg Gly  
-15 -10 -5

His Leu Pro Val Ser Leu Asp Pro Lys  
1 5

## (2) INFORMATION FOR SEQ ID NO: 404:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -38..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.1  
seq GILCGSLPGPSLC/PP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 404:

```
Met Gly Met Ser Gly Lys Lys His Phe Pro Leu Ser Trp Asp His Ile
    -35                      -30                      -25

Gln Gly Ser Thr Glu Ala Thr Ser Gln Gly Ile Leu Cys Gly Ser Leu
    -20                      -15                      -10

Pro Gly Pro Ser Leu Cys Pro Pro
    -5                      1
```

(2) INFORMATION FOR SEQ ID NO: 405:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 74 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -37..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4  
seq PLSLDCCGHS LCRA/CI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 405:

```
Met Ala Ser Lys Ile Leu Leu Asn Val Gln Glu Glu Val Thr Cys Pro
    -35                      -30                      -25

Ile Cys Leu Glu Leu Leu Thr Glu Pro Leu Ser Leu Asp Cys Gly His
    -20                      -15                      -10

Ser Leu Cys Arg Ala Cys Ile Thr Val Ser Asn Lys Glu Ala Val Thr
    -5                      1                      5                      10

Ser Met Gly Gly Lys Ser Ser Cys Pro Val Cys Gly Ile Ser Xaa Ser
```



15

20

25

Xaa Glu His Leu Gln Ala Asn Gln His Arg  
30 35

## (2) INFORMATION FOR SEQ ID NO: 406:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq YYMVCLFFRLIFS/EH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 406:

Met Tyr Tyr Met Val Cys Leu Phe Phe Arg Leu Ile Phe Ser Glu His  
-10 -5 1

Leu Pro Ile Ile Gly Thr Val Thr Ser His Lys Thr Gly Thr Gly  
5 10 15

## (2) INFORMATION FOR SEQ ID NO: 407:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -36..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq KLAGLWSPGLVPA/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 407:

Met Gly Ala Gly Gly Xaa Arg Glu Ile Arg Ala Ala Ala Ser Trp  
 -35 -30 -25

Leu Arg Ala Ala Glu His Ser Lys Leu Ala Gly Leu Trp Ser Pro Gly  
 -20 -15 -10 -5

Leu Val Pro Ala Ala Pro Arg Thr Glu Asn Tyr Thr Ile Gly Pro Leu  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO: 408:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -60..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq LVRRTLLVAALRA/WM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 408:

Met Gly Ser Lys Cys Cys Lys Gly Gly Pro Asp Glu Asp Ala Val Glu  
 -60 -55 -50 -45

Arg Gln Arg Arg Gln Lys Leu Leu Leu Ala Gln Leu His His Arg Lys  
 -40 -35 -30

Arg Val Lys Ala Ala Gly Gln Ile Gln Ala Trp Trp Arg Gly Val Leu  
 -25 -20 -15

Val Arg Arg Thr Leu Leu Val Ala Ala Leu Arg Ala Trp Met Ile Gln  
 -10 -5 1

Cys Trp Trp Arg Thr Leu Val Gln Arg Arg Ile Arg Gln Arg Arg Gln  
 5 10 15 20

## (2) INFORMATION FOR SEQ ID NO: 409:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -26..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4  
seq SIHSWQLL TSAQP/QQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 409:

Met Gln Gln Gly His Pro His Leu Ser Ala Gly Thr Leu Ser Ile His  
-25 -20 -15  
Ser Trp Gln Leu Leu Thr Ser Ala Gln Pro Gln Gln Ala Gly  
-10 -5 1

## (2) INFORMATION FOR SEQ ID NO: 410:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 58 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -49..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4  
seq ATCCLSL FQWCAV/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 410:

Met Ser Arg Tyr Glu Xaa Gly Ser Ser Leu Leu Pro Phe Pro Asp His  
-45 -40 -35  
Phe Ser Val Tyr Ser Phe Lys Xaa Xaa Ser Phe Phe Glu Ala Tyr Ser  
-30 -25 -20  
Ile Ser Asp Tyr Ala Thr Cys Cys Leu Ser Leu Phe Gln Trp Cys Ala  
-15 -10 -5  
Val Leu Arg Phe Leu Ser Leu Pro Leu Pro  
1 5

## (2) INFORMATION FOR SEQ ID NO: 411:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -24..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq LLLHHYLLLFITT/SR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 411:

Met Ile Tyr Phe Ile Lys Ile Asn Asn Lys Leu Leu Leu Leu His His  
-20 -15 -10

Tyr Leu Leu Leu Phe Ile Thr Thr Ser Arg Pro Thr Gly  
-5 1 5

(2) INFORMATION FOR SEQ ID NO: 412:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -27..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq LSWALCLSQSGYY/HP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 412:

Met Glu Leu Leu Tyr Leu Lys Val Lys Arg Gly Gln Lys Asp Leu Ser  
-25 -20 -15

Trp Ala Leu Cys Leu Ser Gln Ser Gly Tyr Tyr His Pro Ser His Pro  
-10 -5 1 5

His Trp

## (2) INFORMATION FOR SEQ ID NO: 413:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq TLAVTLSALGATG/LF

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 413:

Met Thr Leu Ala Val Thr Leu Ser Ala Leu Gly Ala Thr Gly Leu Phe  
                  -10                  -5                  1  
Lys Glu Ala Cys Asp Leu Thr Phe Leu Asn Ile Gly Gln Ile Thr Ser  
          5                          10                          15  
Xaa Leu Lys Gln Ser Gly Gly Pro Gln  
      20                          25

## (2) INFORMATION FOR SEQ ID NO: 414:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 72 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -41..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq CRCLITLPRSCR/P/ST

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 414:

Met Leu Gly Pro Pro Leu Gln Pro Gly Ser His Gly Lys Val Leu Ala  
      -40                  -35                  -30  
Pro Gln Gly Ser Ser Gly Leu Thr Pro Pro Phe Pro Cys Arg Cys Leu

```

-25          -20          -15          -10
Ile Thr Leu Pro Arg Ser Cys Arg Pro Ser Thr Ser Val Pro Arg Xaa
          -5                      1                      5
Ala Ser Thr Arg Ser Ser Gln Arg Pro Xaa Ser Ser Cys Trp Arg Ser
          10                      15                      20
Ser Cys Ser Thr Thr Ala Thr Met
          25                      30

```

(A) LENGTH: 48 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -27..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.8  
seq QLXLILVHFPAYS/VE

```
Met Gly Asn Val Cys Ser Cys Cys Leu Arg Ala Arg Tyr Gln Gln Leu
      -25              -20              -15

Xaa Leu Ile Leu Val His Phe Pro Ala Tyr Ser Val Glu Asp Gln Arg
      -10              -5              1              5

Val Asp Pro Gly Val Pro Gly Glu Ser Thr Val Cys His His Asn Arg
      10              15              20
```

(A) LENGTH: 41 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

1) ORGANISM: Homo Sapiens  
2) TISSUE TYPE: Spleen

(11). FUTURE:

```

Met Leu Tyr Gly Leu Gly Ser Gly Pro Arg Cys Val Ile Ser Cys Ile
  -20                      -15                -10

His Gly Val Trp Cys Glu Glu Gly Asp Gly Ser Leu Pro Arg Leu His
  -5                      1                      5                10

Val Ala Leu Met Ile Pro Ala Leu Gly
      15                20

```

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(E) TISSUE TYPE: Testis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 417:

(2) INFORMATION FOR SEQ ID NO: 418:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -13..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq MLFLVLFYSAIFL/FT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 418:

Met Leu Phe Leu Val Leu Phe Tyr Ser Ala Ile Phe Leu Phe Thr Leu  
          -10                  -5                  1  
  
Thr Phe Phe  
          5

## (2) INFORMATION FOR SEQ ID NO: 419:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq VSLCVAALEFPLQA/YC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 419:

Met Val Ser Leu Cys Val Ala Ala Leu Phe Pro Leu Gln Ala Tyr Gly  
          -10                  -5                  1

## (2) INFORMATION FOR SEQ ID NO: 420:

## (i) SEQUENCE CHARACTERISTICS:



## (2) INFORMATION FOR SEQ ID NO: 422:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq MLPLFCSPWESGG/RT

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 422:

```

Met Met Leu Pro Leu Phe Cys Ser Pro Trp Glu Ser Gly Gly Arg Thr
      -10                      -5                      1
Val Lys Gln Ser Glu Gly Xaa Cys Xaa Phe Gln Ala Pro His Gly
      5                      10                      15

```

## (2) INFORMATION FOR SEQ ID NO: 423:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq KLLSDLSVDSARC/KP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 423:

```

Met Ala Lys Leu Leu Ser Asp Leu Ser Val Asp Ser Ala Arg Cys Lys
-15          -10          -5                      1
Pro Gly Asn Asn Leu Thr Lys Ser Leu Leu Asn Ile His Asp Lys Gln
      5                      10                      15

```

Leu Gln His Asp Pro Arg  
20

(2) INFORMATION FOR SEQ ID NO: 424:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.7  
seq VCVGHLLPARVST/RS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 424:

Met Cys Gly Tyr Trp Val Cys Trp Gly His Leu Leu Pro Ala Arg Val  
-15 -10 -5  
Ser Thr Arg Ser Ser Glu Gln Pro Arg Val Thr Pro Arg Asp Glu Asp  
1 5 10  
Ala Met Met Ser Ala Ser Leu Leu Thr Trp Arg Tyr Val Thr Phe Met  
15 20 25 30  
Val Pro Met Pro Leu Ser Pro Cys Arg Ser Val Trp Val Cys Phe Arg  
35 40 45  
Gln Lys Ile Leu Glu Tyr Val Xaa Ala  
50 55

(2) INFORMATION FOR SEQ ID NO: 425:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -24..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.7  
 seq AILGLSTFLNLLS/IN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 425:

```
Met Lys Leu Ser Cys Ala Gly Cys Ala Asp Thr Ala Ile Leu Gly Leu
      -20              -15              -10

Ser Thr Phe Leu Asn Leu Leu Ser Ile Asn Leu Leu Gly Met Ile Ser
      -5              1              5

Phe Ser
      10
```

(2) INFORMATION FOR SEQ ID NO: 426:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 58 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Spleen
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: -21..-1
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 3.7  
seq FSLGSCPAGPLSA/CV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 426:

```
Met Ile Pro Phe Ser Gly Thr Val Phe Ser Leu Gly Ser Cys Pro Ala
      -20              -15              -10

Gly Pro Leu Ser Ala Cys Val Pro Asp His Gly Ser Leu Gln Tyr Pro
      -5              1              5              10

Leu Thr Ile Tyr Gln Gln Asp Cys Xaa Thr His Xaa Cys Pro Arg Cys
      15              20              25

Leu Ser Leu Pro Leu Gln His Pro Arg Gln
      30              35
```

(2) INFORMATION FOR SEQ ID NO: 427:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 97 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -35..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.7  
seq PAVSLSAPAFASA/LR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 427:

Met Ile Pro Ser Ser Gln Pro Arg Phe Xaa Asn Pro Ala Cys Lys Gln  
-35 -30 -25 -20

Thr Val Leu Leu Xaa Asp Pro Ala Val Ser Leu Ser Ala Pro Ala Phe  
-15 -10 -5

Ala Ser Ala Leu Arg Ser Met Xaa Ser Ser Gln Ala Ala Arg Lys Asp  
1 5 10

Asp Phe Leu Arg Ser Leu Ser Asp Gly Asp Ser Gly Thr Ser Glu His  
15 20 25

Ile Ser Ala Val Val Thr Ser Pro Arg Ile Ser Cys His Gly Ala Ala  
30 35 40 45

Ile Pro Thr Ala Arg Ala Leu Cys Leu Xaa Cys Ser Cys Cys Thr Glu  
50 55 60

Arg

(2) INFORMATION FOR SEQ ID NO: 428:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -15..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.6  
seq PTFLLISDSFLTS/QP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 428:

Met Ala Pro Thr Phe Leu Leu Ile Ser Asp Ser Phe Leu Thr Ser Gln  
 -15                                -10                                -5                                1  
 Pro Ser Phe Phe Phe Phe  
                               5

## (2) INFORMATION FOR SEQ ID NO: 429:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
seq LSLGKIQWCLS/EN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 429:

Met Ile Ser Leu Ile Val Leu Ser Leu Leu Gly Ile Lys Ile Gln Trp  
                               -15                                -10                                -5  
 Cys Leu Ser Glu Asn Thr Leu Phe Cys Asp Ser Asp Tyr Leu Leu Ser  
                               1                                5                                10  
 Pro Lys Ala Pro Ile Glu Pro Leu Ser Phe Asn Leu Thr Thr Gln Gly  
                               15                                20                                25

## (2) INFORMATION FOR SEQ ID NO: 430:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6

seq LLYFNTFLPRKVA/RV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 430:

Met Ala Cys Asp Ser Phe Leu Lys Asp Ala Leu Pro Gln Glu Leu Ser  
                   -40                  -35                  -30

Gln Leu Xaa Phe Leu Phe Pro Leu Val Asp Met Arg Glu Asp Leu Leu  
                   -25                  -20                  -15

Tyr Phe Asn Thr Phe Leu Pro Arg Lys Val Ala Arg Val  
                   -10                  -5                  1

## (2) INFORMATION FOR SEQ ID NO: 431:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -53..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
                                   seq FLILHFFPQQIRK/KI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 431:

Met Leu Leu Leu Asn Glu Asn Leu Lys Ala Glu Ile Gln Lys Asn Glu  
                   -50                  -45                  -40

Ala Gln Gly Ser Cys Ile Leu Phe Leu Phe Cys Phe Glu Ser Gln Asn  
                   -35                  -30                  -25

Met Arg Ser Lys Ser Ile Phe Pro Phe Leu Ile Leu His Phe Phe Pro  
                   -20                  -15                  -10

Gln Gln Ile Arg Lys Lys Ile Val Val Leu Leu Leu Gly Leu Asn Ser  
                   -5                  1                  5                  10

Gln Lys Ala Gly  
                   15

## (2) INFORMATION FOR SEQ ID NO: 432:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 amino acids
- (B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -26..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.5  
seq LLPFTFLSLKAFL/QX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 432:

Met Ile Ser Lys Tyr Val His Tyr Ser Leu Thr Asp Leu Leu Leu Pro  
-25 -20 -15

Phe Thr Phe Leu Ser Leu Lys Ala Phe Leu Gln Xaa Arg Val Leu Met  
-10 -5 1 5

Ser Leu Pro Gln His Lys Pro Trp  
10

(2) INFORMATION FOR SEQ ID NO: 433:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -27..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.5  
seq CSLLSSFCALHFG/LK

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 433:

Met Ala Arg Thr Met Gly Val Pro Arg Ala Cys Lys Ala Phe Cys Ser  
-25 -20 -15

Leu Leu Ser Ser Phe Cys Ala Leu His Phe Gly Leu Lys Lys Gln Tyr  
-10 -5 1 5

Gly Thr Ser Tyr Leu His Ala Cys Ala Tyr Ala Ser Pro Leu Thr Trp  
10 15 20

(2) INFORMATION FOR SEQ ID NO: 434:

(A) LENGTH: 29 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -15..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.5  
seq LCFLLPHHRLOEA/RX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 434:

Met Ile Leu Cys Phe Leu Leu Pro His His Arg Leu Gln Glu Ala Arg  
-15 -10 -5 1

Xaa Ile Gln Val Leu Lys Xaa Leu Pro Arg Glu Lys Leu  
5 10

(2) INFORMATION FOR SEQ ID NO: 435:

(A) LENGTH: 83 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Spleen

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -26..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.5  
seq QCFVFCFSPKIYG/VI

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 435:

Met Gln Asp Tyr Val Ser His Ala Val Arg Arg His Cys Gln Cys Phe  
-25 -20 -15



```

Phe Val Cys Phe Ser Pro Lys Ile Tyr Gly Val Ile Thr Trp Thr Val
-10          -5          1          5

Leu Ile Thr Gly Ala Arg Val Leu Ser Glu Pro Gln Arg Leu Trp Val
          10          15          20

Arg Leu Asp Asp Ile Thr Ala Asn Ala Ala Cys Gly Tyr Arg Lys Gln
          25          30          35

Glu Pro Arg Lys Thr Phe Glu Asn Asn Trp Glu Asn Leu Tyr Thr Asp
          40          45          50

Trp Asn Trp
55

```

(A) LENGTH: 35 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -31..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 3.5  
seq VLLNLALSHFNNC/GL

Met Glu Phe Ala His Ala Ala Glu Cys Val Ser Phe Ala Leu Asn Glu  
-30 -25 -20

Thr His Val Leu Leu Asn Leu Ala Leu Ser His Phe Asn Asn Cys Gly  
-15 -10 -5 1

Leu Ala Val

(A) LENGTH: 48 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens

..(F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -30..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq LLAASWLPRDAPC/EA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 437:

```

Met Gly Asn Gln Gly Phe Pro Tyr Leu Ser Pro Ser Leu Ser Val Gln
-30          -25          -20          -15

Asp Leu Leu Ala Ala Ser Trp Leu Pro Arg Asp Ala Pro Cys Glu Ala
          -10          -5          1

Pro Pro Gly Leu Pro Ser Gln Thr Met Leu Cys Ala Pro Gly Pro Arg
      5          10          15

```

(2) INFORMATION FOR SEQ ID NO: 438:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -22..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq AQLASPLLPGATP/VA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 438:

```

Met Lys Tyr Gln Met Val Ser Gly Ser Ala Gln Leu Ala Ser Pro Leu
      -20          -15          -10

Leu Pro Gly Ala Thr Pro Val Ala Gly Thr Ile Leu Lys Ser Leu Leu
      -5          1          5          10

Leu Arg Thr Val Lys Met Met Arg Val Tyr Gly
      15          20

```

(2) INFORMATION FOR SEQ ID NO: 439:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 amino acids

.. (B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

(ix) FEATURE:  
(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -21..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 12.7  
seq ILFLLSWGPLQG/QQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 439:

Met Gly Pro Ser Thr Pro Leu Leu Ile Leu Phe Leu Leu Ser Trp Ser  
-20 -15 -10

Gly Pro Leu Gln Gly Gln Gln His His Leu Val Glu Tyr Met Glu Arg  
-5 1 5 10

Arg His Gly

(2) INFORMATION FOR SEQ ID NO: 440:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:  
(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(ix) FEATURE:  
(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -20..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 7.4  
seq LVFCVGLLTMAKA/ES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 440:

Met Ala Ser Leu Gly His Ile Leu Val Phe Cys Val Gly Leu Leu Thr  
-20 -15 -10 -5

Met Ala Lys Ala Glu Ser Pro Lys Glu His Asp Pro Arg  
1 5

(2) INFORMATION FOR SEQ ID NO: 441:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: AMINO ACID
- (C) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -23..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.3  
seq ALSLLLVSGSLLP/GP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 441:

```

Met Ser Gly Ser Ser Leu Pro Ser Ala Leu Ala Leu Ser Leu Leu Leu
      -20                      -15                      -10

Val Ser Gly Ser Leu Leu Pro Gly Pro Gly Ala Ala Gln Asn Glu Pro
      -5                      1                      5

Arg Ile Val Thr Ser Glu Glu Val Ile Ile Arg Asp Ser Pro Val
  10                      15                      20

```

## (2) INFORMATION FOR SEQ ID NO: 442:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 65 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -57..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1  
seq VGLAVVSLGGSRG/SG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 442:

```

Met Met Glu Val Val Val Gly Asn Gly Val Val Ala Leu Arg Gly Ile
      -55                      -50                      -45

Pro Pro Arg Thr Ser Arg Lys Ser Ser Arg Lys Thr Arg Phe Cys Gly
      -40                      -35                      -30

```

Glu Arg Gly Ser Lys Gln Ser Gly Lys Cys Ser Pro Val Gly Leu Ala  
 -25 -20 -15 -10

Val Val Ser Leu Gly Gly Ser Arg Gly Ser Gly Lys Gly Leu Gly Arg  
 -5 1 5

Leu

(2) INFORMATION FOR SEQ ID NO: 443:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 58 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.7  
seq CFSLVLLLSIWT/TR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 443:

Met Ala Arg Cys Phe Ser Leu Val Leu Leu Thr Ser Ile Trp Thr  
 -15 -10 -5

Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala Glu Glu Leu Ser Ile  
 1 5 10 15

Gln Val Ser Cys Arg Xaa Met Gly Ile Thr Leu Val Ser Lys Lys Ala  
 20 25 30

Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys  
 35 40

(2) INFORMATION FOR SEQ ID NO: 444:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 136 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -21..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.6  
 seq CLSCLLIPLALWS/II

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 444:

Met Gly Ser Arg Lys Cys Gly Gly Cys Leu Ser Cys Leu Leu Ile Pro  
 -20 -15 -10

Leu Ala Leu Trp Ser Ile Ile Val Asn Ile Leu Leu Tyr Phe Pro Asn  
 -5 1 5 10

Gly Gln Thr Ser Tyr Ala Ser Ser Asn Lys Leu Thr Asn Tyr Val Trp  
 15 20 25

Tyr Phe Glu Gly Ile Cys Phe Ser Gly Ile Met Met Leu Ile Val Thr  
 30 35 40

Thr Val Leu Leu Val Leu Glu Asn Asn Asn Asn Tyr Lys Cys Cys Gln  
 45 50 55

Ser Glu Asn Cys Ser Lys Lys Tyr Val Thr Leu Leu Ser Ile Ile Phe  
 60 65 70 75

Ser Ser Leu Gly Ile Ala Phe Ser Gly Tyr Cys Leu Val Ile Ser Ala  
 80 85 90

Leu Gly Leu Val Gln Gly Pro Tyr Cys Arg Thr Leu Asp Gly Trp Glu  
 95 100 105

Tyr Ala Phe Glu Gly Thr Ala Gly  
 110 115

## (2) INFORMATION FOR SEQ ID NO: 445:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 82 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -21..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.6  
 seq CLSCLLIPLALWS/II

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 445:

Met Gly Ser Arg Lys Cys Gly Gly Cys Leu Ser Cys Leu Leu Ile Pro  
 -20 -15 -10

Leu Ala Leu Trp Ser Ile Ile Val Asn Ile Leu Leu Tyr Phe Pro Asn  
 -5 1 5 10

Gly Gln Thr Ser Tyr Ala Ser Ser Asn Lys Leu Thr Asn Tyr Val Trp  
 15 20 25

Tyr Phe Glu Gly Ile Cys Phe Ser Gly Ile Met Met Leu Ile Val Thr  
 30 35 40

Thr Val Leu Leu Val Leu Glu Asn Asn Asn Asn Tyr Lys Cys Cys Gln  
 45 50 55

Ser Gly  
 60

## (2) INFORMATION FOR SEQ ID NO: 446:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -17..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 5.3  
seq ILFGVSFVFLTHC/TI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 446:

Met Met Val Met Ile Leu Phe Gly Val Ser Phe Val Phe Leu Thr His  
 -15 -10 -5

Cys Thr Ile Gln Ser Ser Cys Gly  
 1 5

## (2) INFORMATION FOR SEQ ID NO: 447:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -19..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.4  
seq VLVSLP#PH#ALT/CC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 447:

```

Met Ser Asn Thr His Thr Val Leu Val Ser Leu Pro His Pro His Pro
      -15                      -10                      -5

Ala Leu Thr Cys Cys His Leu Gly Xaa Pro His Pro Val Arg Ala Pro
      1                      5                      10

Arg Pro
      15

```

## (2) INFORMATION FOR SEQ ID NO: 448:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide  
(B) LOCATION: -106..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.1  
seq IITLACVPMTSFT/RN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 448:

```

Met Xaa Val Tyr Arg Leu Gln Thr Gln Glu Lys Pro Asn Thr Thr Val
-105                      -100                      -95

Gln Val Pro Ala Phe Leu Gln Glu Leu Val Asp Arg Asp Asn Ser Lys
-90                      -85                      -80                      -75

Phe Glu Glu Trp Cys Ile Glu Met Ala Glu Met Arg Xaa Lys Val Trp
      -70                      -65                      -60

Ile Lys Glu Lys Gln Asn Thr Lys Arg Leu Arg Ser Cys Thr Lys Gly
      -55                      -50                      -45

Tyr Leu Leu Glu Leu Ser Pro Met Ser Leu Ser Leu Trp Asn Gly Cys

```



-40                      -35                      -30  
 Lys Ser Gly Trp Met Asn Gln Gln Xaa Pro Asn Leu Leu Ile Ile Thr  
 -25                      -20                      -15  
 Leu Ala Cys Val Pro Met Thr Ser Phe Thr Arg Asn Lys Ile Ser Ile  
 -10                      -5                      1                      5  
 Met Lys Arg Ile Ser Glu Tyr Ala Ala Asp Ile Phe Tyr  
                     10                      15

## (2) INFORMATION FOR SEQ ID NO: 449:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.9  
seq LIAVVIIILLIFT/SV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 449:

Met Phe Pro Val Leu Gly Trp Ile Leu Ile Ala Val Val Ile Ile Ile  
 -20                      -15                      -10  
 Leu Leu Ile Phe Thr Ser Val Thr Arg Cys Leu  
 -5                      1                      5

## (2) INFORMATION FOR SEQ ID NO: 450:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 65 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 3.9  
seq SVCLCPCLNKGQS/EN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 450:

```

Met Phe Ser Cys Cys Ile Ser Val Cys Leu Cys Pro Cys Leu Asn Lys
      -15              -10              -5
Gly Gln Ser Glu Asn Leu Ser Arg Asp Cys Gly His Trp Leu Asn Pro
      1              5              10
His His Arg Arg Leu Trp Pro Phe Gly Arg Arg His Pro Gln Asp Cys
      15              20              25
Gly Leu Phe Gln Asp Ser Gln Xaa Tyr Gly Glu Ser Lys Asp Trp Asn
      30              35              40              45
Gly

```

## (2) INFORMATION FOR SEQ ID NO: 451:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 88 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -29..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq LTYLLLLSPIKYP/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 451:

```

Met Arg Leu Cys Leu Ile Met Tyr Cys Ser Phe Gly Thr Leu Ser His
      -25              -20              -15
Leu Thr Tyr Leu Leu Leu Ser Pro Ile Lys Tyr Pro Leu Asp Leu
      -10              -5              1
Asp Phe Leu Tyr Pro Ile Phe Ser Thr Val Tyr Lys Arg Tyr Ile Val
      5              10              15
Thr Val Asn Phe Cys Ile Ser Cys Ser Glu Ser Phe Leu Leu Ser Asp
      20              25              30              35
Leu Ile Ala Leu Phe Leu Ile Arg Glu Leu Gln Leu Leu Gln His Thr
      40              45              50
Val Ser Val Val Gln Pro Pro Thr

```

## (2) INFORMATION FOR SEQ ID NO: 452:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -26..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 10.5  
seq LLLALLLPVQVSS/FV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 452:

Met Gly Lys Gly Met Val Ala Met Leu Ile Leu Gly Leu Leu Leu Leu  
-25 -20 -15

Ala Leu Leu Leu Pro Val Gln Val Ser Ser Phe Val Pro Leu Thr Ser  
-10 -5 1 5

Met Pro Glu Ala Thr Ala Ala Glu Thr Thr Lys Pro Ser Asn Gly  
10 15 20

## (2) INFORMATION FOR SEQ ID NO: 453:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 9.5  
seq LLVLFVLLANVQG/PG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 453:

(A) LENGTH: 44 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -18..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 8.3  
seq LLSLSSLPLVLLG/WE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 455:

Met Glu Thr Gly Arg Leu Leu Ser Leu Ser Ser Leu Pro Leu Val Leu  
-15 -10 -5

Leu Gly Trp Glu Tyr Ser Ser Gln Thr Leu Asn Leu Val Pro Ser Thr  
1 5 10

Ser Ile Leu Ser Phe Val Pro Phe Ile Pro Arg Val  
15 20 25

(2) INFORMATION FOR SEQ ID NO: 456:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 59 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -19..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 8.2  
seq QVLALVLVAALWG/GT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 456:

Met Ala Ala Ser Leu Gly Gln Val Leu Ala Leu Val Leu Val Ala Ala  
-15 -10 -5

Leu Trp Gly Gly Thr Gln Pro Leu Leu Lys Arg Ala Ser Ala Gly Leu  
1 5 10

Gln Arg Val His Glu Pro Thr Trp Ala Gln Gln Leu Gln Glu Met  
15 20 25

Lys Thr Leu Phe Leu Asn Thr Glu Tyr Leu Met  
30 35 40

## (2) INFORMATION FOR SEQ ID NO: 457:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 84 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -59..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.1  
seq FLLGISNLSQVRA/SN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 457:

```

Met His Ile Lys Ser Ile Ile Leu Glu Gly Phe Lys Ser Tyr Ala Gln
      -55                      -50                      -45

Arg Thr Glu Val Asn Gly Phe Asp Pro Leu Phe Asn Ala Ile Thr Gly
      -40                      -35                      -30

Leu Asn Gly Ser Gly Lys Ser Asn Ile Leu Asp Ser Ile Cys Phe Leu
      -25                      -20                      -15

Leu Gly Ile Ser Asn Leu Ser Gln Val Arg Ala Ser Asn Leu Gln Asp
      -10                      -5                      1                      5

Leu Val Tyr Lys Asn Gly Gln Ala Gly Ile Thr Lys Ala Ser Val Ser
      10                      15                      20

Ile Xaa Phe Asp
      25

```

## (2) INFORMATION FOR SEQ ID NO: 458:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide

- (B) LOCATION: -19...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8  
seq WGFLCVLFTAVHP/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 458:

```

Met Ser Pro Ser Pro Arg Trp Gly Phe Leu Cys Val Leu Phe Thr Ala
      -15                -10                -5

Val His Pro Ala Pro Ser Thr Ala Pro Val Gln Asp Lys Cys Pro Val
      1                5                10

Asn Thr Trp Glu Ala Met Xaa Xaa Val Leu Pro Ala Ala Pro Ala Asn
    15                20                25

Arg Pro Pro Thr Gln Ala Phe Pro Ser Ala Ser Thr Ala Thr Gly
    30                35                40

```

(2) INFORMATION FOR SEQ ID NO: 459:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 65 amino acids
  - (B) TYPE: AMINO ACID
  - (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Homo Sapiens
  - (F) TISSUE TYPE: Testis

- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: -24...-1
  - (C) IDENTIFICATION METHOD: Von Heijne matrix
  - (D) OTHER INFORMATION: score 7.8  
seq FLLCLCIAYWAST/AV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 459:

```

Met Cys Ser Leu Leu Tyr Pro Leu Val Thr Phe Phe Leu Leu Cys Leu
      -20                -15                -10

Cys Ile Ala Tyr Trp Ala Ser Thr Ala Val Phe Leu Ser Thr Ser Asn
      -5                1                5

Glu Ala Val Tyr Lys Ile Phe Asp Asp Ser Pro Cys Pro Phe Thr Ala
    10                15                20

Lys Thr Cys Asn Pro Glu Thr Phe Pro Ser Ser Asn Glu Pro Arg His
    25                30                35                40

Gly

```

## (2) INFORMATION FOR SEQ ID NO: 460:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 51 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.6  
seq FLFFSTLFSSIFT/EA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 460:

Met Leu Pro Phe Leu Phe Phe Ser Thr Leu Phe Ser Ser Ile Phe Thr  
-15 -10 -5

Glu Ala Gln Lys Gln Tyr Trp Val Cys Asn Ser Ser Asp Ala Ser Ile  
1 5 10 15

His Thr Pro Thr Val Ile Lys Cys Asn Thr Gln Phe Gln Leu Met Leu  
20 25 30

Thr Pro Gly  
35

## (2) INFORMATION FOR SEQ ID NO: 461:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.5  
seq VALNLILVPCCAA/WC

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 461:

Met Val Ala Leu Asn Leu Ile Leu Val Pro Cys Cys Ala Ala Trp Cys



```

Met Ala Ala Arg Gly Val Ile Ala Pro Val Gly Glu Ser Leu Arg Tyr
      -50                      -45                      -40

Ala Glu Tyr Leu Gln Pro Ser Ala Lys Arg Pro Asp Ala Asp Val Asp
      -35                      -30                      -25

Gln Gln Arg Leu Val Arg Ser Leu Ile Ala Val Gly Leu Gly Val Ala
      -20                      -15                      -10

Ala Leu Ala Phe Ala Gly Arg Tyr Ala Phe Arg Ile Trp Lys Pro Leu
      -5                      1                      5                      10

Glu Gln Val Ile Thr Glu Thr Ala Lys Lys Ile Ser Thr Pro Ser Phe
      15                      20                      25

Ser Ser Tyr Tyr Lys Gly Gly Phe Glu Gln Lys Met Ser Arg Arg Glu
      30                      35                      40

Ala Gly Leu Ile Leu Gly Val Ser Pro Ser Ala Gly Lys Ala Lys Ile

```

45

50

55

Arg Thr Ala His Arg Arg Val Met Ile  
60 65

## (2) INFORMATION FOR SEQ ID NO: 463:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 7.1  
seq KLKLLSLLR?SLC/IP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 463:

Met Ile Lys Leu Lys Leu Leu Ser Leu Leu Arg Pro Ser Leu Cys Ile  
-15 -10 -5 1

Pro Gln Leu Leu Arg Thr Asn Ala Thr Leu Leu Phe Thr Ile Ala Ser  
5 10 15

Cys Asn Leu Gln Ile Pro Ala Ser Pro Arg Arg  
20 25

## (2) INFORMATION FOR SEQ ID NO: 464:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.5  
seq GLCVLQLTTAVTS/AF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 464:

Met Pro Ser Val Asn Ser Ala Gly Leu Cys Val Leu Gln Leu Thr Thr  
-20                               -15                               -10                               -5  
Ala Val Thr Ser Ala Phe Leu Leu Ala Lys Val Asn Pro Phe Glu Xaa  
                                  1                               5                               10  
Phe Leu Ser Arg Gly Phe Trp Leu Cys Ala  
          15                               20

(2) INFORMATION FOR SEQ ID NO: 465:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -20..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 6.4  
seq ALFLLVSYMIRS/GT

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 465:

Met Met Leu Gly Leu His Phe Ala Leu Phe Leu Leu Val Ser Xaa Tyr  
-20                               -15                               -10                               -5  
Met Ile Arg Ser Gly Thr Gly Asn Lys Ile Glu Glu Gly Gly Arg  
                                  1                               5                               10

(2) INFORMATION FOR SEQ ID NO: 466:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 48 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -13..1

(C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 6.4  
 seq MALLLSVLRVLIG/GF

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 466:

```
Met Ala Leu Leu Leu Ser Val Leu Arg Val Leu Leu Gly Gly Phe Phe
      -10                -5                1
Ala Leu Val Gly Leu Ala Lys Leu Ser Glu Glu Ile Ser Ala Pro Val
      5                10                15
Ser Glu Arg Met Asn Ala Leu Phe Val Xaa Phe Ala Glu Val Leu Gly
      20                25                30                35
```

(2) INFORMATION FOR SEQ ID NO: 467:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -17..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 6.2  
 seq LWLSLVAAHWGEA/VL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 467:

```
Met Leu Lys Ser Leu Trp Leu Ser Leu Val Ala Trp His Trp Gly Glu
      -15                -10                -5
Ala Val Leu Leu Ser Pro His Leu Pro Ala Ala Ala Glu Trp Pro Arg
      1                5                10                15
Ala Ala Cys Asp Ser Gly Gly Glu Pro
      20
```

(2) INFORMATION FOR SEQ ID NO: 468:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Ovary

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -15...-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6.1  
seq IVTWLXSEFSSA/EE

Met Gly Ile Val Thr Trp Leu Leu Xaa Ser Phe Met Ser Ser Ala Glu  
-15 -10 -5 1

Glu Ser Val Ser Ala Arg Thr Arg  
5

(A) LENGTH: 37 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Uterus

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -23..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 6  
seq IGLMFLMLGCAIP/IY

Met Ala Gly Ile Lys Ala Leu Ile Ser Leu Ser Phe Gly Gly Ala Ile  
-25 -20 -15

Gly Leu Met Phe Leu Met Leu Gly Cys Ala Leu Pro Ile Tyr Asn Lys  
-10 -5 1

Tyr Trp Pro Pro Gly  
5

(A) LENGTH: 34 amino acids  
(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Ovary

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -26..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.8  
seq VKLVTLSPVPSLA/SS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 470:

Met Lys Lys Gln Lys His Gln Lys Leu Trp Cys Ile Ser Val Lys Leu  
-25 -20 -15

Val Thr Leu Ser Val Pro Thr Ser Leu Ala Ser Ser Leu Thr Ser Pro  
-10 -5 1 5

Thr Gly

(2) INFORMATION FOR SEQ ID NO: 471:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 78 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -40..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 5.8  
seq VLFALFVAFLLRG/KL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 471:

Met Asp Gly Ile Pro Met Ser Met Lys Asn Glu Met Pro Ile Ser Gln  
-40 -35 -30 -25

Leu Leu Met Ile Ile Ala Pro Ser Leu Gly Phe Val Leu Phe Ala Leu  
-20 -15 -10

Phe Val Ala Phe Leu Leu Arg Gly Lys Leu Met Clu Thr Tyr Cys Ser  
-5 1 5

Gln Lys His Thr Arg Leu Asp Tyr Ile Gly Asp Ser Lys Asn Val Leu

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -19..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 8  
 seq WGFLOVLFTAVHP/AP

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 473:

Met Ser Pro Ser Pro Arg Trp Gly Phe Leu Cys Val Leu Phe Thr Ala  
                           -15                          -10                          -5  
 Val His Pro Ala Pro Ser Thr Ala Pro Val Gln Asp Lys Cys Pro Val  
                           1                                          5                                          10  
 Asn Thr Trp Glu Ala Met Gln Ala Ser Ser Gln Gln Leu Leu Gln Thr  
           15                                          20                                          25  
 Asp Pro Met  
           30

(2) INFORMATION FOR SEQ ID NO: 474:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 93 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -76..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 5.3  
 seq IILASASFSPNFT/QV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 474:

Met Thr Ser Gln Pro Val Pro Asn Glu Thr Ile Ile Val Leu Pro Ser  
           -75                                          -70                                          -65  
 Asn Val Ile Asn Phe Ser Gln Ala Glu Lys Pro Glu Pro Thr Asn Gln  
           -60                                          -55                                          -50                                          -45  
 Gly Gln Asp Ser Leu Lys Lys His Leu His Ala Glu Xaa Lys Val Ile  
                           -40                                          -35                                          -30  
 Gly Thr Ile Gln Ile Leu Cys Gly Met Met Val Leu Ser Leu Gly Ile  
           -25                                          -20                                          -15



Ile Leu Ala Ser Ala Ser Phe Ser Pro Asn Phe Thr Gln Val Thr Ser  
           -10                      -5                      1  
 Thr Leu Leu Asn Ser Ala Tyr Pro Phe Ile Gly Pro Gly  
       5                      10                      15

## (2) INFORMATION FOR SEQ ID NO: 475:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 127 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -91..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.8  
seq IILRLPWLNRSQT/VV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 475:

Met Arg Ala Leu Glu Asn Asp Phe Phe Asn Ser Pro Pro Arg Lys Thr  
       -90                      -85                      -80  
 Val Arg Phe Gly Gly Thr Val Thr Glu Val Leu Leu Lys Tyr Lys Lys  
       -75                      -70                      -65                      -60  
 Gly Glu Thr Asn Asp Phe Glu Leu Leu Lys Asn Gln Leu Leu Asp Pro  
                       -55                      -50                      -45  
 Asp Ile Lys Asp Asp Gln Ile Ile Asn Trp Leu Leu Glu Phe Arg Ser  
                       -40                      -35                      -30  
 Ser Val Met Tyr Leu Thr Lys Asp Phe Glu Gln Leu Ile Ser Ile Ile  
                       -25                      -20                      -15  
 Leu Arg Leu Pro Trp Leu Asn Arg Ser Gln Thr Val Val Glu Glu Tyr  
       -10                      -5                      1                      5  
 Leu Ala Phe Leu Gly Asn Leu Val Ser Ala Glu Thr Val Phe Leu Arg  
                       10                      15                      20  
 Pro Cys Leu Ser Met Ile Ala Ser His Phe Xaa Pro Pro Glu Leu  
                       25                      30                      35

## (2) INFORMATION FOR SEQ ID NO: 476:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 amino acids

(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -31..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.8  
seq WAFSCGTWLPSPRA/EW

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 476:

Met Val Phe Pro Ala Lys Arg Phe Cys Leu Val Pro Ser Met Glu Gly  
-30 -25 -20

Val Arg Trp Ala Phe Ser Cys Gly Thr Trp Leu Pro Ser Arg Ala Glu  
-15 -10 -5 1

Trp Leu Leu Ala Val Arg Ser Ile Gln Pro Glu Glu Lys Glu Arg Ile  
5 10 15

Gly Gln Phe Val Phe Ala Arg Asp Gly  
20 25

(2) INFORMATION FOR SEQ ID NO: 477:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 93 amino acids

(B) TYPE: AMINO ACID

(D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: -82..-1

(C) IDENTIFICATION METHOD: Von Heijne matrix

(D) OTHER INFORMATION: score 4.7  
seq LTCLADLFHSIAT/XX

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 477:

Met Asn Cys Phe Gln Gly Thr Asn Ala Ser Ala Leu Glu Lys Asp Ile  
-80 -75 -70

Gly Pro Glu Gln Phe Pro Ile Asn Glu His Tyr Phe Gly Leu Val Asn  
-65 -60 -55

Phe Gly Asn Thr Cys Tyr Cys Asn Ser Val Leu Gln Ala Leu Tyr Ser  
 -50 -45 -40 -35  
 Cys Arg Pro Phe Arg Glu Asn Val Leu Ala Tyr Lys Ala Gln Gln Lys  
 -30 -25 -20  
 Lys Lys Glu Asn Leu Leu Thr Cys Leu Ala Asp Leu Phe His Ser Ile  
 -15 -10 -5  
 Ala Thr Xaa Lys Lys Lys Val Xaa Ser Ser His Leu Gly  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO: 473:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 55 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -16..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq ALRVRXXXFGTRA/CR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 478:

Met Ala Ala Ala Leu Arg Val Arg Xaa Xaa Xaa Phe Gly Thr Arg Ala  
 -15 -10 -5  
 Cys Arg Arg His Gly Leu Pro His Arg Ala Xaa Trp Leu Arg Asn Arg  
 1 5 10 15  
 Val Xaa Asp Arg Tyr Phe Arg Ile Gln Glu Val Leu Lys Xaa Ala Arg  
 20 25 30  
 His Phe Arg Gly Arg Lys Arg  
 35

## (2) INFORMATION FOR SEQ ID NO: 479:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (v1) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -15..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq LLTHNLLSSHVRG/VG

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 479:

```

Met Lys Leu Leu Thr His Asn Leu Leu Ser Ser His Val Arg Gly Val
-15          -10          -5          1

Gly Ser Arg Gly Phe Pro Leu Arg Leu Gln Ala Thr Glu Val Arg Ile
          5          10          15

Cys Pro Val Glu Phe Asn Pro Asn Phe Val Ala Arg Arg
          20          25          30

```

## (2) INFORMATION FOR SEQ ID NO: 480:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 61 amino acids  
(B) TYPE: AMINO ACID  
(D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
(F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
(B) LOCATION: -41..-1  
(C) IDENTIFICATION METHOD: Von Heijne matrix  
(D) OTHER INFORMATION: score 4.6  
seq VSAGSLLLPAQA/EX

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 480:

```

Met Gly Xaa Phe Ser Arg Arg Thr Phe Cys Gly Arg Ser Gly Arg Ser
-40          -35          -30

Cys Arg Gly Gln Leu Val Gln Val Ser Arg Pro Glu Val Ser Ala Gly
-25          -20          -15          -10

Ser Leu Leu Leu Pro Ala Pro Gln Ala Glu Xaa His Ser Ser Xaa Xaa
          -5          1          5

Leu Tyr Pro Arg Pro Lys Ser Leu Leu Pro Lys Met Gly
          10          15          20

```

## (2) INFORMATION FOR SEQ ID NO: 481:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 79 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -55..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.6  
seq CALSLPDAFGASG/GR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 481:

Met Glu Gly Gly Val Arg Leu Asp Leu Ser Ala Cys Gly Glu Thr Ser  
-55 -50 -45 -40  
Gly Val Ala Val Ser Glu Leu Pro Ala Ser Glu Thr Ala Ala Leu Val  
-35 -30 -25  
Pro Glu Gly His Gly Pro Gly Leu Arg Ala Cys Ala Leu Ser Leu Pro  
-20 -15 -10  
Asp Ala Pro Gly Ala Ser Gly Gly Arg His His Leu Ile Leu Val Pro  
-5 1 5  
Gly Gln Gln His Thr Gly Leu Pro Ala Ser His Val His Pro Gln  
10 15 20

## (2) INFORMATION FOR SEQ ID NO: 482:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.5  
seq TLLSFAALTAAFS/VL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 482:

Met Thr Leu Leu Ser Phe Ala Ala Leu Thr Ala Ala Phe Ser Val Leu  
 -10 -5 1

Pro Lys

## (2) INFORMATION FOR SEQ ID NO: 483:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 78 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -21...-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.4  
seq GLSKLQFAPFSSA/LD

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 483:

Met Ala Ala Ala Thr Gly Asp Pro Gly Leu Ser Lys Leu Gln Phe Ala  
 -20 -15 -10

Pro Phe Ser Ser Ala Leu Asp Val Gly Phe Trp His Glu Leu Thr Gln  
 -5 1 5 10

Lys Lys Leu Asn Glu Tyr Arg Leu Asp Glu Ala Pro Lys Asp Ile Lys  
 15 20 25

Gly Tyr Tyr Tyr Asn Gly Asp Ser Ala Gly Xaa Pro Ala Arg Leu Thr  
 30 35 40

Leu Glu Phe Ser Ala Phe Asp Met Ser Ala Pro Thr Pro Ser  
 45 50 55

## (2) INFORMATION FOR SEQ ID NO: 484:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens

(F) TISSUE TYPE: Spleen

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -27..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.4  
 seq LSKSLLLVPXKLS/LL

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 484:

Met Phe Thr Ser Thr Gly Ser Ser Gly Leu Tyr Lys Ala Pro Leu Ser  
 -25 -20 -15  
 Lys Ser Leu Leu Leu Val Pro Ser Xaa Leu Ser Leu Leu Xaa Ala Gln  
 -10 -5 1 5

(2) INFORMATION FOR SEQ ID NO: 485:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -40..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.3  
 seq ITLVSAAPGXVIC/EM

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 485:

Met Thr Ser Met Thr Gln Ser Leu Arg Glu Val Ile Lys Ala Met Thr  
 -40 -35 -30 -25  
 Lys Ala Arg Asn Phe Glu Arg Val Leu Gly Lys Ile Thr Leu Val Ser  
 -20 -15 -10  
 Ala Ala Pro Gly Lys Val Ile Cys Glu Met Lys Val Glu Glu Glu His  
 -5 1 5  
 Thr Asn Ala Ile Gly Thr Leu His Gly Gly Leu Thr Ala Thr Leu Val  
 10 15 20  
 Asp Asn Ile Ser Thr Met Ala Leu Leu Cys Thr Glu Arg Gly Ala Pro  
 25 30 35 40  
 Gly Val Ser

## (2) INFORMATION FOR SEQ ID NO: 486:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -73..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.3  
seq DIILSGLVPGST7/LH

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 486:

Met Ala Asp Phe Gly Ile Ser Ala Gly Gln Phe Val Ala Val Val Trp  
                  -70                  -65                  -60  
Asp Lys Ser Ser Pro Val Glu Ala Leu Lys Gly Leu Val Asp Lys Leu  
                  -55                  -50                  -45  
Gln Ala Leu Thr Gly Asn Glu Gly Arg Val Ser Val Glu Asn Ile Lys  
                  -40                  -35                  -30  
Gln Leu Leu Gln Ser Ala His Lys Glu Ser Ser Xaa Asp Ile Ile Leu  
                  -25                  -20                  -15                  -10  
Ser Gly Leu Val Pro Gly Ser Thr Thr Leu His Ser Ala Glu Ile Leu  
                  -5                                  1                                  5  
Ala Glu Ile Ala Arg Val  
                  10

## (2) INFORMATION FOR SEQ ID NO: 487:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix



(D) OTHER INFORMATION: score 4.2  
seq GILLGLLLLGHLT/VR

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 487:

Met Gly Ile Leu Leu Gly Leu Leu Leu Gly His Leu Thr Val Arg  
-10 -5 1

(2) INFORMATION FOR SEQ ID NO: 488:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 144 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (E) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -19..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1  
seq LLLGQRCSLKVSG/QE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 488:

Met Phe Leu Thr Val Lys Leu Leu Leu Gly Gln Arg Cys Ser Leu Lys  
-15 -10 -5

Val Ser Gly Gln Glu Ser Val Ala Thr Leu Lys Arg Leu Val Ser Arg  
1 5 10

Arg Leu Lys Val Pro Glu Glu Gln Gln His Leu Leu Phe Arg Gly Gln  
15 20 25

Leu Leu Glu Asp Asp Lys His Leu Ser Asp Tyr Cys Ile Gly Pro Asn  
30 35 40 45

Ala Ser Ile Asn Val Ile Met Gln Pro Leu Glu Lys Met Ala Leu Lys  
50 55 60

Glu Ala His Gln Pro Gln Thr Gln Pro Leu Trp His Gln Leu Gly Leu  
65 70 75

Val Leu Ala Lys His Phe Glu Pro Gln Asp Ala Lys Ala Val Leu Gln  
80 85 90

Leu Leu Arg Phe Glu His Glu Glu Arg Leu Gln Lys Ile Ser Leu Glu  
95 100 105

His Leu Gln Gln Leu Ala Gln Tyr Leu Leu Ala Glu Glu Leu Thr Trp  
110 115 120 125

## (2) INFORMATION FOR SEQ ID NO: 489:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 106 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -32..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4.1  
seq RLLSSLLLTMSNN/NP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 489:

Met Asn Val Ile Asp His Val Arg Asp Met Ala Ala Ala Gly Leu His  
-30 -25 -20

Ser Asn Val Arg Leu Leu Ser Ser Leu Leu Leu Thr Met Ser Asn Asn  
-15 -10 -5

Asn Pro Glu Leu Phe Ser Pro Pro Gln Lys Tyr Gln Leu Leu Val Tyr  
1 5 10 15

His Ala Asp Ser Leu Phe His Asp Lys Glu Tyr Arg Asn Ala Val Ser  
20 25 30

Lys Tyr Thr Met Ala Leu Gln Gln Lys Lys Ala Leu Ser Lys Thr Ser  
35 40 45

Lys Val Arg Pro Ser Thr Gly Asn Ser Ala Ser Thr Pro Gln Ser Gln  
50 55 60

Cys Leu Pro Ser Glu Ile Glu Val Lys Tyr  
65 70

## (2) INFORMATION FOR SEQ ID NO: 490:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 59 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -41...-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 4.1  
 seq RVLCPILKXAAAAP/KR

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 490:

Met Gly Thr Pro Ser Leu Ser Ile Leu Leu Ile Gly Ala Pro Glu Ser  
 -40 -35 -30  
 Pro Ile Pro Tyr Phe Pro Tyr His Ser Gly Thr Gly Arg Val Leu Cys  
 -25 -20 -15 -10  
 Pro Leu Leu Xaa Ala Ala Ala Ala Pro Lys Arg Asp Val Pro Glu Thr  
 -5 1 5  
 Gly Leu Thr Arg Gln Leu Lys Arg His Pro Gly  
 10 15

## (2) INFORMATION FOR SEQ ID NO: 491:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 57 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -26...-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.9  
 seq HALFVLCCLLYAMS/HN

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 491:

Met Val Tyr His Ala Leu Asp Ser Pro Asp Asp Asp Tyr His Ala Leu  
 -25 -20 -15  
 Phe Val Leu Cys Leu Leu Tyr Ala Met Ser His Asn Lys Gly Met Asp  
 -10 -5 1 5  
 Pro Glu Lys Leu Glu Arg Ile Gln Leu Pro Val Pro Asn Ala Ala Glu  
 10 15 20  
 Lys Thr Thr Tyr Asn His Pro His Gly  
 25 30

## (2) INFORMATION FOR SEQ ID NO: 492:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -14..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq FIVLSMWLCCGFE/IL

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 492:

Met Phe Ile Val Leu Ser Met Trp Leu Cys Cys Gly Phe Glu Ile Leu  
                  -10                  -5                  1

Gln Thr Lys Ser Trp Val Ala Gly  
          5                  10

## (2) INFORMATION FOR SEQ ID NO: 493:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq VVILSSXVPLAAM/GV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 493:

Met Val Val Val Ile Leu Ser Ser Xaa Val Pro Leu Ala Ala Met Gly  
-15                  -10                  -5                  1

Val Met Gly Cys Val Arg Val Trp  
          5

## (2) INFORMATION FOR SEQ ID NO: 494:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 67 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq AECSSLLHPSVRG/SI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 494:

```

Met Leu Ala Glu Cys Ser Ser Leu Leu His Pro Ser Val Arg Gly Ser
-15              -10              -5              1

Ile Pro Glu Ala Thr Cys Arg Val Leu Pro Cys Gly Pro Leu His Asn
          5              10              15

Met Ala Val Cys Ser Cys Lys Ala Ser Arg Ser Phe Tyr Cys Asn Phe
      20              25              30

Arg Ser Leu Arg Leu Ala Val Ser Asp Phe Leu Ile Leu Phe Gln Lys
  35              40              45

Gly Leu Gly
  50

```

## (2) INFORMATION FOR SEQ ID NO: 495:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq MARLLGLCAWARK/SV

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 495:

Met Gln Met Ala Arg Leu Leu Gly Leu Cys Ala Trp Ala Arg Lys Ser  
-15                   -10                   -5                   1  
Val Arg Met Ala Ser Ser Arg Met Thr Arg Arg Asp Pro Pro Arg  
          5                   10                   15

(2) INFORMATION FOR SEQ ID NO: 496:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Uterus

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -43..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.8  
seq LISVLYLIPKTLT/TN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 496:

Met Thr Pro Gln Tyr Leu Pro His Gly Gly Lys Tyr Gln Val Leu Gly  
          -40                   -35                   -30  
Asp Tyr Ser Leu Ala Val Val Phe Pro Leu His Phe Ser Asp Leu Ile  
          -25                   -20                   -15  
Ser Val Leu Tyr Leu Ile Pro Lys Thr Leu Thr Thr Asn Ser Arg  
          -10                   -5                   1

(2) INFORMATION FOR SEQ ID NO: 497:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 115 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -23..-1

- (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.7  
 seq FLPPFLXRAFACRG/CQ

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 497:

```

Met Val Val Leu Arg Ala Gly Lys Lys Thr Phe Leu Pro Pro Leu Xaa
      -20              -15              -10

Arg Ala Phe Ala Cys Arg Gly Cys Gln Leu Ala Pro Glu Arg Gly Ala
      -5              1              5

Glu Arg Arg Asp Thr Ala Pro Ser Gly Val Ser Arg Phe Cys Pro Pro
10              15              20              25

Arg Lys Ser Cys His Asp Trp Ile Gly Pro Pro Asp Lys Tyr Ser Asn
              30              35              40

Leu Arg Pro Val His Phe Tyr Ile Pro Glu Asn Glu Ser Pro Leu Glu
      45              50              55

Gln Lys Leu Arg Lys Leu Arg Gln Glu Thr Gln Glu Trp Asn Gln Gln
      60              65              70

Phe Trp Ala Asn Gln Asn Leu Thr Phe Ser Lys Glu Lys Glu Glu Phe
      75              80              85

Ile His Ser
      90
  
```

(2) INFORMATION FOR SEQ ID NO: 498:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 33 amino acids  
 (B) TYPE: AMINO ACID  
 (D) TOPOLOGY: LINEAR
- (ii) MOLECULE TYPE: PROTEIN
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Homo Sapiens  
 (F) TISSUE TYPE: Testis
- (ix) FEATURE:  
 (A) NAME/KEY: sig\_peptide  
 (B) LOCATION: -20..-1  
 (C) IDENTIFICATION METHOD: Von Heijne matrix  
 (D) OTHER INFORMATION: score 3.6  
 seq AHLCSDSLPEQQ/QD

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 498:

```

Met Lys Arg Glu Gly Gly Ala Ala His Leu Cys Ser Asp Ser Leu Pro
-20              -15              -10              -5

Glu Ser Gln Gln Gln Asp Gly Asn His Ala Pro Asn Phe Ser Ser His
      1              5              10
  
```

Gly

## (2) INFORMATION FOR SEQ ID NO: 499:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -41..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.6  
seq PYSLAACPCGSQG/GV

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 499:

Met Val Thr Cys Pro Gly Pro Ser Ser Gly Gln Pro Leu Ser Ser Met  
-40 -35 -30

Tyr Thr Ala Gly Asp Arg Arg Gly Ala Pro Ser Leu Pro Tyr Ser Leu  
-25 -20 -15 -10

Ala Ala Cys Pro Cys Gly Ser Gln Gly Gly Val Cys Met Arg  
-5 1 5

## (2) INFORMATION FOR SEQ ID NO: 500:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 104 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Spleen

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -18..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq ALEVIVTLSETAA/AM

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 500:



Met Gln Arg Gln Leu Ala Leu Glu Val Ile Val Thr Leu Ser Glu Thr  
 -15 -10 -5

Ala Ala Ala Met Leu Arg Lys His Thr Asn Ile Val Ala Gln Thr Ile  
 1 5 10

Pro Gln Met Leu Ala Met Met Val Asp Leu Glu Glu Asp Glu Asp Trp  
 15 20 25 30

Ala Asn Ala Asp Glu Leu Glu Asp Asp Asp Phe Asp Ser Asn Ala Val  
 35 40 45

Ala Gly Glu Ser Ala Leu Asp Arg Met Ala Cys Gly Leu Gly Gly Lys  
 50 55 60

Leu Val Leu Pro Met Ile Lys Glu His Ile Met Gln Met Leu Gln Asn  
 65 70 75

Arg Lys Leu Cys Pro Ser Met Leu  
 80 85

## (2) INFORMATION FOR SEQ ID NO: 501:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 82 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

## (ii) MOLECULE TYPE: PROTEIN

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

## (ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -76..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 3.5  
seq LASASELP LGSRP/AP

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 501:

Met Gly Asp Tyr Leu Leu Arg Gly Tyr Arg Met Leu Gly Glu Thr Cys  
 -75 -70 -65

Ala Asp Cys Gly Thr Ile Leu Leu Gln Asp Lys Gln Arg Lys Ile Tyr  
 -60 -55 -50 -45

Cys Val Ala Cys Gln Glu Leu Asp Ser Asp Val Asp Lys Asp Asn Pro  
 -40 -35 -30

Ala Leu Asn Ala Gln Ala Ala Leu Ser Gln Ala Arg Glu His Gln Leu  
 -25 -20 -15

Ala Ser Ala Ser Glu Leu Pro Leu Gly Ser Arg Pro Ala Pro Gln Pro  
 -10 -5 1

His Gly  
5

(2) INFORMATION FOR SEQ ID NO: 502:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 41 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Ovary

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -15..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 8.1  
seq LLYLLVPALFCRA/GG

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 502:

Met Trp Leu Leu Tyr Leu Leu Val Pro Ala Leu Phe Cys Arg Ala Gly  
-15                      -10                      -5                      1  
Gly Ser Ile Pro Ile Pro Gln Lys Leu Phe Gly Glu Val Thr Ser Pro  
                    5                      10                      15  
Leu Phe Pro Lys Pro Tyr Pro Asn Thr  
                    20                      25

(2) INFORMATION FOR SEQ ID NO: 503:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 75 amino acids
- (B) TYPE: AMINO ACID
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo Sapiens
- (F) TISSUE TYPE: Testis

(ix) FEATURE:

- (A) NAME/KEY: sig\_peptide
- (B) LOCATION: -58..-1
- (C) IDENTIFICATION METHOD: Von Heijne matrix
- (D) OTHER INFORMATION: score 4  
seq LAAVSPLVRSLS/SN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 503:

Met Lys Leu Glu Phe Thr Glu Lys Asn Xaa Xaa Ser Phe Val Leu Gln  
-55 -50 -45

Asn Leu Asn Arg Gln Arg Lys Arg Lys Glu Tyr Trp Asp Met Ala Leu  
-40 -35 -30

Ser Val Asp Asn His Val Phe Phe Ala His Arg Asn Val Leu Ala Ala  
-25 -20 -15

Val Ser Pro Leu Val Arg Ser Leu Ile Ser Ser Asn Asp Met Lys Thr  
-10 -5 1 5

Ala Asp Glu Leu Phe Ile Thr Ile Asp Thr Lys  
10 15

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/is 98/01231

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C12N15/10 C12N15/66 C12Q1/68  
 G01N33/50 C07K16/18 G01N33/53 A61K48/00 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                               | Relevant to claim No. |
|------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| E          | WO 98 46755 A (MCCARTHY SEAN A ;MILLENNIUM BIOTHERAPEUTICS INC (US)) 22 October 1998<br>see the claims<br>see page 7, paragraph 2; figure 5<br>see page 10, line 17 - line 26<br>see page 50, line 32 - page 80, line 15<br>SEQ. ID: 13 and 14<br>see page 107 - page 109<br>--- | 1-37                  |
| X          | HILLIER L ET AL: "Homo sapiens cDNA clone 728407 (AC No. AA397836)"<br>EMBL SEQUENCE DATABASE, 28 April 1997,<br>XP002083926<br>Heidelberg, Germany<br>see the whole document<br>---                                                                                             | 3-10,<br>15-34        |
| Y          | ---                                                                                                                                                                                                                                                                              | 35-37                 |
|            | ---<br>-/--                                                                                                                                                                                                                                                                      |                       |

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

11 November 1998

Date of mailing of the international search report

0 8. 02 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
 Fax (+31-70) 340-3016

Authorized officer

Oderwald, H

## INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/IB 98/01231

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                    | Relevant to claim No. |
|----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| Y        | LOCKHART D J ET AL: "EXPRESSION MONITORING BY HYBRIDIZATION TO HIGH-DENSITY OLIGONUCLEOTIDE ARRAYS" BIO/TECHNOLOGY, vol. 14, no. 13, December 1996, pages 1675-1680, XP002022521<br>see abstract; figures 1,2<br>see page 1678, paragraph 3 - page 1679, paragraph 2                                                  | 35-37                 |
| A        | ---<br>YOKOYAMA-KOBAYASHI M. ET AL.: "A signal sequence detection system using secreted protease activity as an indicator" GENE, vol. 163, 1995, pages 193-196, XP002053953<br>see abstract                                                                                                                           | 12,13                 |
| A        | ---<br>LIN Y ET AL: "INHIBITION OF NUCLEAR TRANSLOCATION OF TRANSCRIPTION FACTOR NF-KB BY A SYNTHETIC PEPTIDE CONTAINING A CELL MEMBRANE-PERMEABLE MOTIF AND NUCLEAR LOCALIZATION SEQUENCE" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 24, 16 June 1995, pages 14255-14258, XP002050723<br>see abstract; figure 1 | 14                    |
| A        | ---<br>WO 96 34981 A (GENSET (FR); MERENKOVA IRENA NICOLAEVNA; DUMAS MILNE EDWARDS JEAN) 7 November 1996<br>cited in the application                                                                                                                                                                                  |                       |
| A        | ---<br>KATO S. ET AL.: "Construction of a human full-length cDNA bank" GENE, vol. 150, 1994, pages 243-250, XP002081364<br>cited in the application                                                                                                                                                                   |                       |
| A        | ---<br>EP 0 625 572 A (KANAGAWA ACAD OF SCIENCE AND TECHNOL FOUNDATION (JP); KATO S; SEKINE S) 23 November 1994<br>cited in the application                                                                                                                                                                           |                       |
| A        | ---<br>CARNINCI P. ET AL.: "High-efficiency full-length cDNA cloning by biotinylated CAP trapper" GENOMICS, vol. 37, no. 3, 1 November 1996, pages 327-336, XP002081729<br>cited in the application                                                                                                                   |                       |
| A        | ---<br>WO 97 07198 A (GENETICS INSTITUTE INC (US); JACOBS K; MCCOY JM; KELLEHER K; CARLIN M) 27 February 1997<br>---                                                                                                                                                                                                  |                       |
|          | ---<br>-/--                                                                                                                                                                                                                                                                                                           |                       |

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 98/01231

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                          | Relevant to claim No. |
|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| A          | <p>TASHIRO K. ET AL.: "Signal sequence trap:<br/>a cloning strategy for secreted proteins<br/>and type I membrane proteins"<br/>SCIENCE,<br/>vol. 261, 30 July 1993, pages 600-603,<br/>XP000673204</p>     |                       |
| A          | <p>HEIJNE VON G.: "A new method for<br/>predicting signal sequence cleavage sites"<br/>NUCLEIC ACIDS RESEARCH,<br/>vol. 14, no. 11, 1986, pages 4683-4690,<br/>XP002053954<br/>cited in the application</p> |                       |

# INTERNATIONAL SEARCH REPORT

Intern. application No.  
PCT/IB 98/01231

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-37 partially (Invention 1. on continuation-sheet)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: Invention 1: 1-37 all partially

Nucleic acid comprising sequence as in Seq. ID:38, complementary sequence, fragments, hybridising sequences. Polypeptide comprising a signal peptide encoded by said nucleotide sequence. Vector encoding a fusion protein comprising said signal peptide. Method of directing the extracellular secretion of a polypeptide by means of said vector. Method of importing a polypeptide into a cell by means of said signal peptide. Method of making a cDNA encoding a secretory protein, partially encoded by said nucleotide sequence, corresponding cDNA. Polypeptide encoded by said nucleotide sequence, comprising sequence as in Seq. ID:271, method of making said polypeptide. Method of obtaining a promoter located upstream of said nucleotide sequence, promoter thereof.

2. Claims: Inventions 2-233: 1-37 all partially

Idem as subject 1 but limited to each of the DNA sequences as in Seq.ID:39-270, and corresponding polypeptides, where invention 2 is limited to Seq.ID:39 and 272, invention 3 is limited to Seq.ID:40 and 273,....., invention 233 is limited to Seq.ID:270 and 503.

For the sake of conciseness, the first subject matter is explicitly defined, the other subject matters are defined by analogy thereto.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01231

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|-------------------------------------------|---------------------|----------------------------|---------------------|
| WO 9846755 A                              | 22-10-1998          | AU 7137398 A               | 11-11-1998          |
| WO 9634981 A                              | 07-11-1996          | FR 2733765 A               | 08-11-1996          |
|                                           |                     | FR 2733762 A               | 08-11-1996          |
|                                           |                     | AU 5982996 A               | 21-11-1996          |
|                                           |                     | CA 2220045 A               | 07-11-1996          |
|                                           |                     | EP 0824598 A               | 25-02-1996          |
| EP 0625572 A                              | 23-11-1994          | JP 6153953 A               | 03-06-1994          |
|                                           |                     | WO 9408001 A               | 14-04-1994          |
|                                           |                     | US 5597713 A               | 28-01-1997          |
| WO 9707198 A                              | 27-02-1997          | US 5707829 A               | 13-01-1998          |
|                                           |                     | AU 6712396 A               | 18-02-1997          |
|                                           |                     | AU 6768596 A               | 12-03-1997          |
|                                           |                     | CA 2227220 A               | 06-02-1997          |
|                                           |                     | CA 2229208 A               | 27-02-1997          |
|                                           |                     | EP 0839196 A               | 06-05-1998          |
|                                           |                     | EP 0851875 A               | 08-07-1998          |
|                                           |                     | WO 9704097 A               | 06-02-1997          |